

## Contents

### iii Graphical Abstracts

#### Regular papers

- |                                                                                                                                                                                                        |     |                                                                                                                                                                                              |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| M.G. O'Shea, M.S. Samuel, C.M. Konik and M.K. Morrell (Canberra, Australia)                                                                                                                            | 1   | Fluorophore-assisted carbohydrate electrophoresis (FACE) of oligosaccharides: efficiency of labelling and high-resolution separation                                                         |
| R. Olsson, P. Rundström, B. Persson and T. Frejd (Lund, Sweden)                                                                                                                                        | 13  | Organotitanium-induced stereoselective alkylative <i>endo</i> -cleavage of benzyl pentopyranosides                                                                                           |
| G. Wulff, A. Steinert and O. Höller (Düsseldorf, Germany)                                                                                                                                              | 19  | Modification of amylose and investigation of its inclusion behaviour                                                                                                                         |
| O. Zegrocka, W. Abramski, Z. Urbańczyk-Lipkowska and M. Chmielewski (Warsaw, Poland)                                                                                                                   | 33  | Formation of $\beta$ -lactams fused to the pyranoid ring via the Mitsunobu reaction                                                                                                          |
| A. Blennow, A.M. Bay-Smidt, B. Wischmann, C.E. Olsen and B. Lindberg Møller (Copenhagen, Denmark)                                                                                                      | 45  | The degree of starch phosphorylation is related to the chain length distribution of the neutral and the phosphorylated chains of amylopectin                                                 |
| B. Grzeszczyk (Warsaw, Poland), O. Holst, S. Müller-Loennies (Borstel, Germany) and A. Zamojski (Warsaw, Poland)                                                                                       | 55  | Five monophosphates of methyl L-glycero- $\alpha$ -D-manno-heptopyranoside: synthesis, hydrolysis and migrations                                                                             |
| R. Uchida, A. Nasu, S. Tokutake, K. Kasai, K. Tobe and N. Yamaji (Chiba, Japan)                                                                                                                        | 69  | New enzymatic synthesis of 6 <sup>3</sup> -modified maltooligosaccharides and their inhibitory activities for human $\alpha$ -amylases                                                       |
| L. Sun and E.L. Chaikof (Atlanta, USA)                                                                                                                                                                 | 77  | The synthesis of neoglycophospholipid conjugates via reductive amination of $\omega$ -oxo-alkylglycosides and phosphatidylethanolamines                                                      |
| S. Ogawa, M. Ashiura and C. Uchida (Yokohama, Japan)                                                                                                                                                   | 83  | Synthesis of $\alpha$ -glucosidase inhibitors: kojibiose-type pseudodisaccharides and a related pseudotrisaccharide                                                                          |
| K.-S. Shin, H. Kiyohara, T. Matsumoto and H. Yamada (Tokyo, Japan)                                                                                                                                     | 97  | Rhamnogalacturonan II dimers cross-linked by borate diesters from the leaves of <i>Panax ginseng</i> C.A. Meyer are responsible for expression of their IL-6 production enhancing activities |
| M.R. Vignon and C. Gey (Grenoble, France)                                                                                                                                                              | 107 | Isolation, <sup>1</sup> H and <sup>13</sup> C NMR studies of (4-O-methyl-D-glucurono)-D-xylans from luffa fruit fibres, jute bast fibres and mucilage of quince tree seeds                   |
| D. Garozzo, G. Impallomeni, E. Spina and L. Sturiale (Catania, Italy)                                                                                                                                  | 113 | The structure of the exocellular polysaccharide from the cyanobacterium <i>Cyanospira capsulata</i>                                                                                          |
| M.T. Dueñas-Chasco (San Sebastian, Spain), M.A. Rodríguez-Carvajal, P. Tejero-Mateo, J.L. Espartero (Seville, Spain), A. Irastorza-Iribas (San Sebastian, Spain) and A.M. Gil-Serrano (Seville, Spain) | 125 | Structural analysis of the exopolysaccharides produced by <i>Lactobacillus</i> spp. G-77                                                                                                     |
| J. Müthing, U. Maurer (Bielefeld, Germany), U. Neumann (Hannover, Germany), B. Kniep (Dresden, Germany) and S. Weber-Schürholz (Bielefeld, Germany)                                                    | 135 | Glycosphingolipids of skeletal muscle: I. Subcellular distribution of neutral glycosphingolipids and gangliosides in rabbit skeletal muscle                                                  |
| J. Müthing, U. Maurer and S. Weber-Schürholz (Bielefeld, Germany)                                                                                                                                      | 147 | Glycosphingolipids of skeletal muscle: II. Modulation of Ca <sup>2+</sup> -flux in triad membranes by gangliosides                                                                           |

## Notes

- Y. Wang (Birmingham, UK), G.W.J. Fleet (Oxford, UK) and L. Zhao (Birmingham, UK) 159 1,1,3,3-Tetraisopropylidisiloxane-1,3-diyl (TIPS) migration facilitates one-pot formation of enantiomerically pure methyl 2,3-expoxy-D-xylonate
- C. Husson, L. Odier and Ph.J.A. Vottéro (Grenoble, France), 163 New conditions for the synthesis of *scyllo*-inositol starting from *myo*-inositol
- C. De Castro (Napoli, Italy), A. Evidente (Portici, Italy), R. Lanzetta (Napoli, Italy), P. Lavermicocca (Bari, Italy), E. Manzo, A. Molinaro and M. Parrilli (Napoli, Italy) 167 Presence of  $\beta$ -glycosyl linkages in caryophyllan: the main polysaccharide from the *Pseudomonas caryophylli* LPS fraction
- A. Gamian, E. Katzenellenbogen, E. Romanowska (Wrocław, Poland), H. Grosskurth and J. Dabrowski (Heidelberg, Germany) 173 Reinvestigation of the O-specific polysaccharides of *Hafnia alvei* lipopolysaccharides isolated from strains ATCC 13337 and 1187
- E. Vinogradov, B. Petersen and K. Bock (Copenhagen, Denmark) 177 Structural analysis of the intact polysaccharide mannan from *Saccharomyces cerevisiae* yeast using  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy at 750 MHz
- 185 Book review
- 187 News Release
- 189 Announcement
- iii Graphical Abstracts

## Regular papers

- T. Aree, J. Jacob, W. Saenger and H. Hoier (Berlin, Germany) 191 Crystal structure of  $\alpha$ -cyclodextrin-acetonitrile-hexahydrate
- K.N. Drew, J. Zajicek (Notre Dame, USA), G. Bondo (South Bend, USA), B. Bose and A.S. Serianni (Notre Dame, USA) 199  $^{13}\text{C}$ -labeled aldopentoses: detection and quantitation of cyclic and acyclic forms by heteronuclear 1D and 2D NMR spectroscopy
- N. Nishimura, M. Banno, A. Maki, Y. Nishiyama and I. Maeba (Nagoya, Japan) 211 Synthesis of 3- $\beta$ -D-ribofuranosylpyrazole-1-carboxamide
- Y. Takagi, N. Kobayashi, M.S. Chang, G.-J. Lim and T. Tsuchiya (Kawasaki, Japan) 217 Synthesis and antitumor activity of the 7-*O*-(2,6-dideoxy-2-fluoro- $\alpha$ -L-talopyranosyl)-dauno-mycinone derivatives modified at C-3' or C-4'
- H.C. Hansen and G. Magnusson (Lund, Sweden) 233 Synthesis of some amino and carboxy analogs of galabiose; evaluation as inhibitors of the pilus protein PapG<sub>96</sub> from *Escherichia coli*
- H.C. Hansen and G. Magnusson (Lund, Sweden) 243 Synthesis of some divalent *O*- and *S*-glycosides of galabiose and globotriose
- K.S. Rangappa, H. Manjunathaswamy, M.P. Raghavendra (Mysore, India) and D.C. Gowda (Washington, DC, USA) 253 Oxidation of threose-series pentoses and hexoses by sodium *N*-chloro-*p*-toluenesulfonamide
- E. Wiczorek and J. Thiem (Hamburg, Germany) 263 Preparation of modified glycosyl glycerol derivatives by glycal rearrangement
- Š. Karácsonyi, V. Pätoprstý and M. Kubačková (Bratislava, Slovak Republic) 271 Structural study on arabinogalactan-proteins from *Picea abies* L. Karst
- T. Terabayashi (Sagamihara, Japan) and Y. Kawanishi (Chiyoda-ku, Japan) 281 Naturally occurring ganglioside lactones in Minke whale brain
- O.M. Hanniffy (Galway, Ireland), A.S. Shashkov, S.N. Senchenkova (Moscow, Russian Federation), S.V. Tomshich, N.A. Komandrova, L.A. Romanenko (Vladivostok, Russian Federation), Y.A. Knirel (Moscow, Russian Federation) and A.V. Savage (Galway, Ireland) 291 Structure of a highly acidic O-specific polysaccharide of lipopolysaccharide of *Pseudoalteromonas haloplanktis* KMM 223 (44-1) containing L-iduronic acid and D-QuNHb4NHb

- S.E.C. Whitney, J.E. Brigham, A.H. Darke (Bedfordshire, UK), J.S.G. Reid (Stirling, UK) and M.J. Gidley (Bedfordshire, UK) 299 Structural aspects of the interaction of mannan-based polysaccharides with bacterial cellulose
- M.M. Rahman (Athens, USA), D.S. Stephens, C.M. Kahler (Atlanta, USA), J. Glushka and R.W. Carlson (Athens, USA) 311 The lipooligosaccharide (LOS) of *Neisseria meningitidis* Serogroup B Strain NMB contains L2, L3, and novel oligosaccharides, and lacks the lipid-A 4'-phosphate substituent
- R. Falshaw and R.H. Furneaux (Lower Hutt, New Zealand) 325 Structural analysis of carrageenans from the tetrasporic stages of the red algae, *Gigartina lanceata* and *Gigartina chapmanii* (Gigartinaceae, Rhodophyta)
- P. Cimino, M.M. Corsaro, C. De Castro (Napoli, Italy), A. Evidente (Portici, Italy), C.E. Marciano, M. Parrilli (Napoli, Italy), A. Sfalanga and G. Surico (Firenze, Italy) 333 Structural determination of the O-deacetylated O-chain of lipopolysaccharide from *Burkholderia* (*Pseudomonas*) *cepacia* strain PVFi-5A

## Notes

- E.B. Starikov, W. Saenger and T. Steiner (Berlin, Germany) 343 Quantum chemical calculations on the weak polar host–guest interactions in crystalline cyclomaltoheptaose ( $\beta$ -cyclodextrin)-but-2-yne-1,4-diol heptahydrate
- J. Wang and R.I. Hollingsworth (East Lansing, USA) 347 The identification of glycerol-3-yl 6-deoxy-6-C-sulfo- $\alpha$ -D-glucopyranoside (glyceryl  $\alpha$ -sulfoquinovoside) as a metabolite in *Rhizobium*, a non-photosynthetic organism
- H.B. Mereyala and S.R. Gurralla (Hyderabad, India) 351 A highly diastereoselective, practical synthesis of allyl, propargyl 2,3,4,6-tetra-O-acetyl- $\beta$ -D-glucopyranosides and allyl, propargyl heptaacetyl- $\beta$ -D-lactosides
- C. Crombez-Robert, M. Benazza, C. Fréhou and G. Demailly (Amiens, France) 355 Tin-mediated regioselective etherification and esterification of unprotected xylitol
- P.I. Kitov, C. Railton and D.R. Bundle (Alberta, Canada) 361 The synthesis of 16-mercaptohexadecanyl glycosides for biosensor applications
- C.E. Heath, M.C. Gillam, C.S. Callis, R.R. Patto and C.J. Abelt (Williamsburg, USA) 371 Ulose formation by selenoxide elimination
- M. Pajatsch, A. Böck (München, Germany) and W. Boos (Konstanz, Germany) 375 Enzymatic preparation of radiolabeled linear maltodextrins and cyclodextrins of high specific activity from [ $^{14}$ C] maltose using amylomaltase, cyclodextrin glucosyltransferase and cyclodextrinase

- I Author index
- III Subject index
- IX Volume contents