



The structure of *Cryptococcus neoformans* galactoxylomannan contains β -D-glucuronic acid

Christian Heiss^{a,*}, J. Stacey Klutts^b, Zhirui Wang^a, Tamara L. Doering^c, Parastoo Azadi^a

^a Analytical Services/Complex Carbohydrate Research Center—University of Georgia, 315 Riverbend Road, Athens, GA 30602, United States

^b University of Iowa Carver College of Medicine and VA Medical Center, 601 Highway 6 West, Iowa City, IA 52246, United States

^c Department of Molecular Microbiology, Washington University School of Medicine, 660 S. Euclid Avenue, St. Louis, MO 63110, United States

ARTICLE INFO

Article history:

Received 15 January 2009

Received in revised form 26 February 2009

Accepted 2 March 2009

Available online 10 March 2009

Keywords:

Cryptococcus neoformans

GalXM

CPS

Glucuronic acid

Polysaccharides, structure

ABSTRACT

The structure of galactoxylomannan, a capsular polysaccharide from the opportunistic yeast *Cryptococcus neoformans*, was re-examined by NMR spectroscopy and GC–MS. The residue that is 3-linked to the side chain galactose and was previously assigned as β -D-xylose [Vaishnav, V. V.; Bacon, B. E.; O'Neill, M.; Cherniak, R. *Carbohydr. Res.* **1998**, 306, 315–330] was determined to be β -D-glucuronic acid. A revised structure for this polymer is presented, along with a proposal that this compound be termed glucuronoxylomannogalactan (GXMGal).

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Cryptococcus neoformans is an opportunistic fungal pathogen of humans and other mammals. This organism causes serious diseases, including lung infections and fatal meningitis.¹ It presents a substantial therapeutic challenge, especially in the developing world.¹ The main virulence factor of *C. neoformans* is the carbohydrate capsule surrounding the cell. Strains of *C. neoformans* lacking this structure do not cause disease in animal models,² suggesting that the mechanisms of capsule synthesis are potential drug targets.

The cryptococcal capsule consists of two polysaccharides. Most of the capsule mass is composed of a glucuronoxylomannan (GXM) of at least 1000 kDa. NMR-based studies of this polymer determined that it has an α -(1→3)-mannan backbone substituted with GlcA and Xyl.³ The remainder of the capsule is composed of a smaller polymer that has been termed galactoxylomannan (GalXM).

The NMR spectra of intact GalXM are highly complex due to the structure of this polysaccharide and the non-stoichiometric glycosylation of some of its residues. Despite these challenges, Cherniak and colleagues proposed a structure for this polysaccharide a decade ago.⁴ According to that report, GalXM is a 100 kDa polysaccharide, consisting of an α -(1→6)-galactan backbone with galactomannan side chains that are further substituted with variable numbers of xylose residues (Fig 1A). In this structure determination,

the authors relied primarily upon degradation studies, using NMR spectroscopy to elucidate the structures of fragments from periodate oxidation and acetolysis. Xylose residues did not survive either of the depolymerization protocols and therefore were not observed in these experiments. An HMBC experiment on the intact polysaccharide confirmed the linkage of two xylose residues to mannose, but the identity of the substituent on O-3 of the side chain galactose (Gal^{III} in Fig. 1) could only be inferred by indirect evidence.

In the course of studying *C. neoformans* capsule synthesis, we identified a β -(1→2)-xylosyltransferase (Cxt1p) and generated a mutant strain (*cxt1Δ*) devoid of this activity. The *cxt1Δ* strain produces GalXM that lacks both of the mannose-linked xylose residues.⁵ This change simplifies the NMR spectra and aids in their interpretation through comparison with spectra of the wild-type polysaccharide. While performing such analyses, we discovered a discrepancy between our composition, linkage, and 2-D NMR data and the published GalXM structure. Here we propose a revised structure of this capsular polysaccharide.

2. Results and discussion

We initiated parallel structural analyses of GalXM isolated from cells with (wild type) and without (*cxt1Δ*) the gene encoding the xylosyltransferase Cxt1p. Monosaccharide composition analysis of GalXM by GC–MS of the per-O-TMS methyl glycoside derivatives showed the expected levels of xylose in the wild-type polysaccharide, but showed only traces of xylose in the polysaccharide from

* Corresponding author. Tel.: +1 706 542 3008; fax: +1 706 542 4412.

E-mail addresses: cheiss@ccrc.uga.edu, heiss8@bellsouth.net (C. Heiss).




Figure 1. Structure of GalXM: (A) previously published structure; (B) revised structure. Subscripts indicate the coefficient of substitution. Previously published data for the structure in panel A (4) indicate that these coefficients are between 0 and 1, but specific values were not provided. Values from our studies (5) for the revised structure are: wild type: $x = 0.7$, $y = 1$, $z = 0.3$; *cxt1Δ*: x and $y < 0.02$, $z = 0.4$. Superscripts are included for easier reference.

strains lacking Cxt1p activity (Fig. 2). Assuming that this xylose is part of the polymer, the amount found corresponds to a degree of

substitution of less than 2%. Likewise, methylation analysis of wild-type GalXM detected terminal xylose, but no terminal xylose was detected in mutant GalXM.⁵

Our composition and methylation analysis experiments indicated that the *cxt1Δ* mutation essentially abolished incorporation of xylose into the polymer. However, the ratio of 4-Gal to 3,4-Gal was the same in both wild type and mutant, indicating that a residue linked to O-3 of Gal^{III} was still present in the mutant at wild-type levels. The identification of this residue as xylose in the published structure raised the question of why we had not detected xylose in methylation analysis and why we had detected only a trace in the composition analysis of the mutant polysaccharide. We had previously attributed the lack of xylose in our GC–MS analysis to evaporation of the volatile partially permethylated xylitol acetate derivative,⁵ but our results from NMR spectroscopy have led us to re-examine this conclusion.

The COSY spectrum of wild-type GalXM showed three H-1/H-2 cross peaks in the region where xylose might be expected, that is, in the region between 4.7 and 4.4 ppm (H-1) and between 3.3 and 3.4 ppm (H-2). Although two separate complete spin systems consistent with xylose were identifiable, the third one appeared to be partially obscured by the first two (data not shown). Based on the published structure of GalXM, we had previously assumed all three of these residues to be xylose. 2-D NOESY demonstrated NOE contacts between H-1 of each of these residues and their linking sugars, providing us with the position of each residue on the GalXM side chain. Thus, Xyl^I is linked to O-2 of Man^I, Xyl^{II} to O-3 of Man^{II}, and the third residue to O-3 of Gal^{III}.

In the COSY spectrum of mutant GalXM the two xylose spin systems that had been completely assigned were missing, whereas the third residue, previously thought to be xylose, was still present. Figure 3 shows part of the COSY spectrum of mutant GalXM, marked to indicate the positions at which the xylose cross peaks are located in the wild-type spectrum. It is clear that the xylose spin systems are missing in the mutant; in particular the conspic-




Figure 2. A comparison of the GC–MS profiles of TMS methyl glycosides of GalXM. Lower trace, wild type; upper trace, *cxt1Δ*.




Figure 3. Partial gCOSY NMR spectrum of *cxt1Δ* GalXM. The GlcA spin system is connected by solid lines. The positions of the xylose peaks from the wild-type spectrum are marked by empty squares (cross peaks) and filled circles (diagonal peaks) and are connected by broken lines. Lower case Roman numerals correspond to the entry number in Table 1. Abbreviations include: e, equatorial; a, axial. The data were acquired at a spectrometer frequency of 500 MHz.

uous absence of the cross peaks between H-4 and H-5 and between H-5a and H-5e proves that the sugar residue in question is not xylose. According to NOESY, this residue is linked to O-3 of Gal^{III} (Fig. 4). The critical NOESY cross peak (x_1 -iv₃) overlaps with the Gal^{III} H-1/H-3 intraresidue cross peak (iv₁-₃). However, the increased linewidth and the center position of the resulting, combined cross peak, just between the two spin systems, indicate that it is made up of two overlapping signals.

We did observe a weak COSY cross peak near the position of H-1 of one of the xylose residues (Fig. 3), which we previously attributed to a Man-linked xylose residue present in low levels in the mutant GalXM (~10% of wild-type levels).⁵ However, the other resonances in this spin system are not consistent with that of xylose, suggesting that xylose is essentially absent in *cxt1Δ* GalXM. The coefficients given in Figure 1 for Xyl reflect this new information.

Having concluded that the substituent on O-3 of Gal^{III} was not xylose, we began to suspect that this residue might be glucuronic acid (GlcA). The chemical shifts of H-1, H-2, and H-3 of glucuronic acid are similar to those of xylose. Furthermore, H-3 and H-4 of GlcA have nearly identical chemical shifts, and H-5 of GlcA resonates at the same chemical shift as H-4 of xylose. Thus the GlcA spin system looks like that of xylose without the H-5 signals. We reasoned that if the residue were GlcA, then its chemical shifts should change with pD and the associated acid dissociation. To test our hypothesis, we acquired TOCSY spectra at two different pD values and discovered that the chemical shifts of the residue in question are indeed dependent on solution pD (Table 1). Furthermore, the carbon chemical shifts of this spin system, obtained from HSQC experiments of both wild-type and mutant GalXM, were also consistent with β-GlcA. (Table 1). Since carbonyl carbons are not seen by HSQC, and HMBC was not sensitive enough to detect C-6 of GlcA, we acquired a 100,000-scan, directly detected carbon spectrum on wild-type GalXM, and found a weak signal at 176 ppm, further indicating the presence of a uronic acid (data not shown). The low intensity of this signal is accounted for by its carbonyl




Figure 4. Partial NOESY NMR spectrum of *cxt1Δ* GalXM, demonstrating the linkage between GlcA and O-3 of Gal^{III}. Lower case Roman numerals correspond to the entry number in Table 1. The data were acquired at a spectrometer frequency of 600 MHz.

Table 1
Proton/carbon chemical shift assignment of wild-type GalXM^a

Entry	Residue		Chemical shift ^b (ppm)						NOE
			1	2	3	4	5	6	
i	→3)-α-Manp ^{II}	¹ H	5.22	4.25	4.03	3.81	4.04	3.69/3.74	iii-3
		¹³ C	102.6	68.6	79.1	66.0	70.5	61.5	
ii	α-Manp ^{II}	¹ H	5.16	4.07	ND	ND	ND	ND	
		¹³ C	103.1	ND	ND	ND	ND	ND	
iii	→2,3)-α-Manp ^I	¹ H	4.99	4.12	4.07	3.87	4.08	3.86/3.84	iv-4, v-4
		¹³ C	100.6	79.2	76.5	67.2	71.0	61.3	
iv	→3,4)-β-Galp ^{III}	¹ H	4.69	3.84	3.86	4.30	3.79	3.80/3.75	vii-3,4
		¹³ C	105.1	71.6	81.5	76.8	74.2	62.0	
v	→4)-β-Galp ^{III}	¹ H	4.66	3.64	3.75	4.09	3.77	3.89/3.80	vii-3,4
		¹³ C	105.4	72.3	73.1	78.4	75.8	62.0	
vi	→6)-α-Galp ^I	¹ H	4.98	3.85	3.89	4.04	4.20	3.93/3.71	vii-6
		¹³ C	99.2	69.3	70.5	70.5	69.8	67.6	
vii	→3,6)-α-Galp ^{II}	¹ H	5.01	4.06	4.01	4.29	4.20	3.93/3.71	vi-6
		¹³ C	99.1	68.3	80.9	70.1	69.8	67.6	
viii	β-Xylp ^{II}	¹ H	4.54	3.39	3.49	3.68	4.02/3.32		i-2,3
		¹³ C	101.9	73.9	76.6	70.1	66.1		
ix	β-Xylp ^I	¹ H	4.39	3.31	3.46	3.64	4.02/3.34		iii-2
		¹³ C	104.4	73.6	76.6	70.2	66.1		
x	β-GlcAp	pD 4.6	4.75	3.39	3.56	3.56	3.87		iv-3
		pD 7.4	4.70	3.37	3.53	3.56	3.68		
		pD 4.6	¹³ C	105.1	74.0	76.4	72.7	75.7	
		pD 7.4		105.0	74.3	76.5	72.5	76.4	

^a All chemical shifts were the same (±0.02 ppm for ¹H and ±0.2 ppm for ¹³C) at pD 4.6 and pD 7.4 except for those of the GlcA residue.

^b ND, not determined.

character and the non-stoichiometric presence of GlcA. Importantly, the proton chemical shift changes observed with varying pD agree with those published previously,⁶ demonstrating downfield movement upon protonation. A 1-D carbon spectrum to measure the chemical shift of the carboxyl carbon was only acquired at pD 7.4; the other GlcA carbon shifts are nearly the same at both pD values, although C-5 experiences a minor upfield shift upon protonation. This shift is opposite to the downfield movement of the proton chemical shifts, but such apparently counterintuitive behavior has been observed previously for carboxylic acids and amino acids,⁷ and has been explained by theory.⁸

Our experiments at different pD supported our hypothesis that the Gal^{III} substituent was GlcA. We next re-examined the gas chromatograms from monosaccharide composition analysis (Fig. 2) and detected a minor amount of GlcA in both wild type and mutant. Although the peaks were small, they were clearly identified as uronic acid peaks by their mass spectra. The spectrum of the signal at 18.17 min had an abundant ion at *m/z* 230 and the spectra of peaks at 24.93 and 25.23 min had significant ions at *m/z* 234; these masses are characteristic for uronic acids.^{9,10} The retention times and the relative peak intensities were also the same as those of a GlcA standard. The GlcA peaks appear small because the response factor of GlcA in GC–MS is low (0.38 or about 50% relative to those of the other sugars present), because of sub-stoichiometric substitution, and because the other sugars are beyond the linear range of the TMS composition analysis. We have previously observed overestimation of sugars present at high levels (unpublished results). Table 2 lists the experimentally determined compositions of both samples.

We noted that the mutant sample contained a significant amount of glucose, which was derived most likely from a lower molecular weight contamination. It is likely that the signal in the NMR spectrum that was previously attributed to Man-linked Xyl (see above) arises from this contamination. According to the chemical shift of its C-1 (104.0 ppm), this glucose is not free. However, there are no cross peaks to this signal in NOESY, indicating that this glucose has an NOE correlation time distinct from that of GalXM and is therefore not part of that polysaccharide. We also noted trace arabinose in each of the samples, which we attribute to contamination.

Table 2
Monosaccharide composition analysis of wild type and *cxt1Δ* GalXM

Glycose	Mol %	
	Wild type	<i>cxt1Δ</i>
Xylose	21	0.30
Glucuronic acid	1.5	2.0
Mannose	26	32
Galactose	51	63
Glucose	0.30	3.0
Arabinose	0.20	0.20

To further demonstrate the presence of GlcA in GalXM, we performed methylation analysis. GlcA is not detected in methylation analysis protocols using NaOH because the strongly basic conditions cause dehydration of GlcA by β-elimination, formation of a C-4–C-5 double bond, and further degradation. Although GlcA can be methylated without major degradation if dimsyl anion is used as base, it is necessary to reduce the carboxyl group of the resulting methyl ester to the alcohol; without reduction the ester would be hydrolyzed in the subsequent acid treatment, leading to a nonvolatile carboxylate. For these reasons we performed our analysis with Hakomori permethylation and prereduction of carboxyl groups. This study demonstrated the presence of (terminal) GlcA (Table 3). Prereduction of GlcA with Super-Deuteride[®] yields doubly ²H-labeled glucose; we detected this compound in GC–MS by extracting the ion at *m/z* = 207 (solid trace in Fig. 5); it was thus distinguished from contaminating Glc already present before prereduction, which has an ion at *m/z* = 205 (broken trace in Fig. 5). We did note that t-Xyl was low and 3-Man was high in the wild-type GalXM, which may indeed be due to evaporation of the volatile partially methylated xylitol acetate, but overall the results of the analysis are in excellent agreement with the structure proposed in Figure 1B.

Finally, we performed GC–MS analysis of the TMS (R)-(–)-2-butyl glycosides (not shown). This further confirmed the presence of GlcA and revealed that it is in the D-form.

The question arises of why GlcA was not detected in the previous structure determination of GalXM. Monosaccharide composition in

Table 3
Methylation analysis with prereduction and Hakomori permethylation of GalXM^a

Residue	Mol %	
	Wild type	<i>cxt1Δ</i>
t-Xyl	3.6	NF
t-Man	5.8	14.6
t-Glc	1.0	1.9
t-GlcA	2.1	4.6
t-Hexf	3.0	2.2
t-Gal	1.6	1.3
3-Glc	NF	0.5
4-Xyl	0.9	NF
3-Man	14.4	18.1
3-Gal	NF	1.7
4-Man	3.8	1.7
4-Gal	10.0	9.8
6-Glc	1.7	2.0
4-Glc	3.5	1.2
2,3-Man	14.0	NF
6-Gal	14.0	17.8
3,4-Gal	5.7	6.3
3,6-Glc	NF	0.3
3,6-Man	NF	0.2
4,6-Gal	NF	0.3
3,6-Gal	12.5	15.4
2,3,6-Man	2.0	NF
2,3,6-Gal	0.4	NF

^a NF, not found; Hexf, unidentified hexofuranose.




Figure 5. Extracted ion chromatograms of the methylation analysis with Hakomori permethylation and prereduction. Solid line: *m/z* 207; broken line: *m/z* 205.

that work was obtained from GC of per-O-acetylated aldononitriles, a method that is unable to detect uronic acids without prior carboxyl reduction;¹¹ this reduction was not performed.⁴ Likewise, methylation analysis detects GlcA only if performed with prereduction. It is therefore not surprising that this monosaccharide was not detected in the earlier study. For technical reasons mentioned above, the NMR identification of GlcA in this structural context is also challenging.

In conclusion, we propose that the residue that is (1→3)-linked to the side chain galactose of GalXM is β-D-glucuronic acid, instead of β-D-xylose as previously determined. We note that the presence of GlcA is consistent with earlier characterization of GalXM¹² and a later compositional study using high pH anion-exchange chromatography,¹³ although not with the structure that has been generally accepted.⁴

We further propose that this polymer be renamed. The original designation of this material as ‘galactoxylomannan’ was based on compositional analyses that indicated it contained galactose, xylose, and mannose. However, previous analysis has demonstrated that this polymer is a galactan, rather than a mannan,⁴ and we now show that it further includes GlcA. The structure shown in

Figure 1B is therefore a ‘glucuronoxylomannogalactan’, which could be abbreviated as GXMGal. This nomenclature accurately represents both the polymer composition and structure, and parallels the common terminology for the second capsule polysaccharide: glucuronoxylomannan abbreviated as GXM.

Our revision of structure explains the chromatographic properties of this polymer, namely, its binding to anion-exchange resin.^{4,5} It also has significant implications for studies of capsule polysaccharide biosynthesis. For example, the previous structure implied the action of up to three xylosyltransferases in GalXM (GXMGal) synthesis; our revision indicates that one of the activities should instead be a glucuronosyltransferase. Our results, consistent with the earlier analysis of this polysaccharide, indicate that the side branches of the polymer are not completely substituted (see coefficients in Fig. 1 legend). Biosynthetic explanations for this heterogeneity could include incomplete sequential modification of a previously assembled core structure, polymer assembly from heterogeneous subunits, or a combination of processes. Additional studies of the GXMGal biosynthetic pathway that characterize its enzymatic features and identify any precursor units will be required to elucidate this question.

Cryptococcal polysaccharides differ dramatically from those of the mammals afflicted by this pathogen. Complete determination of these polysaccharide structures and their biosynthetic pathways will advance our understanding of basic biochemical functions and may bring us nearer to developing effective antifungal therapies.

3. Experimental

3.1. Strains

Like the cells used for the original analysis of GalXM structure,⁴ the strains used in this work are serotype D ‘Cap67’ strains. These cells do not produce GXM, simplifying purification and analysis of GalXM. For text clarity, we refer below to Cap67 strains with normal Cxt1p activity as wild type, and to Cap67 strains lacking the gene encoding Cxt1p as *cxt1Δ*.

3.2. Purification of GalXM

Growth of cells and isolation of GalXM were carried out as described previously.⁵ The final gel filtration step was of particular importance for obtaining high-quality samples for analysis.

3.3. Monosaccharide composition analysis

A 200-μg aliquot of GalXM was heated for 16 h in 1 M methanolic HCl at 80 °C. The sample was then per-O-trimethylsilylated by treatment with Tri-Sil® (Pierce) at 80 °C for 20 min. GC–MS analysis of the TMS methyl glycosides was performed on an Agilent 6890N gas chromatograph interfaced to a 5975B mass-selective detector (MSD), using an Alltech EC-1 fused-silica capillary column (30 m × 0.25 mm ID).

3.4. Methylation analysis

A 600-μg aliquot of the GalXM sample was suspended in about 200 μL of DMSO and magnetically stirred for 5 d. The sample was then permethylated by the method of Anumula and Taylor.¹⁴ NaOH (200 μL) was added and, after 10 min, 100 μL of CH₃I was added, and the sample was stirred for 40 min. An additional 200 μL of base and 100 μL of CH₃I were then added and stirring was continued for 40 min. The reaction was worked up by addition of 2 mL of H₂O, removal of excess CH₃I by sparging with N₂, and CH₂Cl₂ extraction. The permethylated material was hydrolyzed

using 2 M TFA (2 h in a sealed tube at 121 °C), reduced with NaBD₄, and acetylated using Ac₂O/TFA. The resulting partially methylated alditol acetates (PMAAs) were analyzed on a Hewlett–Packard 5890 GC interfaced to a 5970 MSD; separation was performed on a 30-m Supelco 2330 bonded-phase fused-silica capillary column.

3.5. Methylation analysis with prereduction of carboxyl groups

For glycosyl linkage analysis, the sample was permethylated, reduced, re-permethylated, depolymerized, reduced, and acetylated; and the resultant PMAAs were analyzed by GC–MS as described by York et al.¹⁵ First, an aliquot of the sample was suspended in about 200 µL of DMSO and placed on a magnetic stirrer for 5 d, at which time 0.6 mL of potassium dimethylsilylate (3.6 M) was added to each sample. After 7 h of stirring at room temperature, the reaction mixture was cooled to 0 °C, excess CH₃I (0.7 mL) was added dropwise, and the tube was sealed. The incubation was then continued overnight at room temperature. After quenching the reaction with H₂O, removing excess CH₃I with N₂, and extracting with CH₂Cl₂, the permethylated samples were reduced with 200 µL of 1.0 M Super Deuteride® in THF under an N₂ atmosphere (105 min, 25 °C). The mixture was neutralized with three drops of glacial HOAc, dried under a stream of N₂, taken up in 50% EtOH, and passed through an On-Guard H® column. After drying, the samples were re-methylated with NaOH and CH₃I in DMSO and analyzed as described under Section 3.4.

3.6. NMR spectroscopy

Samples were deuterium-exchanged by lyophilization from D₂O and dissolved in 0.7 mL of D₂O. Proton, carbon, gradient-enhanced COSY (gCOSY), TOCSY, NOESY, and HSQC NMR spectra were acquired on Varian Inova-500 MHz and 600 MHz spectrometers at 70 °C using standard Varian pulse sequences. Chemical shifts were measured relative to internal acetone (δ_{H} 2.225 ppm, δ_{C} 31.07 ppm). Mixing times were 100 ms for TOCSY and 300 ms for NOESY.

3.7. Determination of D- or L-forms

A sample (1 g) was hydrolyzed in 2 M TFA at 125 °C for 90 min. The water and acid were evaporated by a stream of nitrogen, and the nonvolatile residue was heated with (R)-(–)-2-BuOH and 15 µL of TFA at 125 °C for 18 h. TMS derivatization and GC–MS were performed as described above for monosaccharide analysis. D-GlcA was derivatized with (R)-(–)-2-BuOH and (S)-(+)-2-BuOH to provide reference standards.

Acknowledgments

Studies of capsule polysaccharides in the Doering laboratory were supported by NIH Grant R01 GM71007 and in the Azadi laboratory by DOE Grant DE-FG09-93ER-20097. We thank Steven Levary for helpful suggestions on the manuscript and T.L.D. thanks Robert Cherniak for encouragement.

References

1. Chayakulkeeree, M.; Perfect, J. R. *Infect. Dis. Clin. North Am.* **2006**, *20*, 507–544.
2. Kwon-Chung, K. J.; Rhodes, J. C. *Infect. Immun.* **1986**, *51*, 218–223.
3. Cherniak, R.; Valafar, H.; Morris, L. C.; Valafar, F. *Clin. Diagn. Lab. Immunol.* **1998**, *5*, 146–159.
4. Vaishnav, V. V.; Bacon, B. E.; O'Neill, M.; Cherniak, R. *Carbohydr. Res.* **1998**, *306*, 315–330.
5. Klutts, J. S.; Doering, T. L. *J. Biol. Chem.* **2008**, *283*, 14234–14327.
6. Lahaye, M.; Ray, B. *Carbohydr. Res.* **1996**, *283*, 161–173.
7. Horsley, W. J.; Sternlicht, H. J. *Am. Chem. Soc.* **1968**, *90*, 3738–3748.
8. Batchelor, J. G.; Feeney, J.; Roberts, G. C. K. *J. Magn. Reson.* **1975**, *20*, 9–38.
9. Kennedy, J. F.; Robertson, S. M. *Carbohydr. Res.* **1978**, *67*, 1–15.
10. Petrzika, M.; Linow, F. *Eur. J. Mass Spectrom. Biochem., Med. Environ. Res.* **1982**, *2*, 53–61.
11. Seymour, F. R. *Methods Carbohydr. Chem.* **1993**, *9*, 59–85.
12. Cherniak, R.; Reiss, E.; Turner, S. H. *Carbohydr. Res.* **1982**, *103*, 239–250.
13. Corradini, C.; Canali, G.; Cavazza, A.; Delfino, D.; Teti, G. *J. Liq. Chromatogr. Related Technol.* **1998**, *21*, 941–951.
14. Anumula, K. R.; Taylor, P. B. *Anal. Biochem.* **1992**, *203*, 101–108.
15. York, W. S.; Darvill, A. G.; McNeil, M.; Stevenson, T. T.; Albersheim, P. *Methods Enzymol.* **1985**, *118*, 3–40.