

Masculinization and Tumour-Inducing Activity of Pregneninolone in the Female Gonads of the Viviparous Toothcarp *Xiphophorus helleri*.

In experimental sex inversion by the use of sex hormones in viviparous Toothcarps the duration of the experiments plays an important role. A high percentage of *Lebistes* males treated from birth with oestrogenic substances, appeared to develop ovotestes¹. The expected result of the treatment with androgenic hormones would be the development of ovotestes in the treated females. When young *Lebistes* females were treated with pregnenolone from birth for a period of two to three months, the ovaries showed a considerable regression without losing their specific female structure^{2,3}. After the cessation of the treatment, the ovaries returned to their normal state³. Treating young *Lebistes* from birth for six months with pregnenolone (Esthisterone, UCLAF, high dose of 0.03 mg, low dose of 0.015 mg, three times a week) out of 55 treated fishes 37 male gonads, 15 hermaphrodite gonads and 2 female gonads were observed⁴. Out of 16 control animals, 11 gonads were male and 5 female. As we personally could demonstrate masculinization and tumour-inducing activity of pregnenolone in the female gonads of *Lebistes*⁵, it was desirable for us to perform a corresponding investigation with another Toothcarp. As experimental material we chose the swordtail *Xiphophorus helleri*. 90 young fishes from the same litter were treated from birth during a 12 month period with pregnenolone

(0.03 mg, three times a week). The microscopical examination of the gonads revealed that 69 treated fishes were males, 18 had hermaphrodite gonads and 3 were females. Out of 21 control animals from the same litter 18 were of the male and 3 of the female sex. In 6 out of the 69 males a testicular tumour had developed. Type seminoma: masses of large, rounded, polyhedral cells with distinct cell boundaries, having large spherical vesicular nuclei and pale, poorly-staining cytoplasm, plentiful mitoses, stroma consisting of strands of connective tissue with blood vessels and lymphoid regions.

The masculinization of the fishes corresponded very well with that obtained in *Lebistes*^{4,5}.

Zusammenfassung. Es wird die maskulinisierende und Tumor erzeugende Aktivität von Pregneninolon beim lebendgebärenden Zahnkarpfen *Xiphophorus helleri* beschrieben.

A. STOLK

Histological Laboratory, Free University, Amsterdam (The Netherlands), July 31, 1961.

¹ P. BERKOWITZ, J. exp. Zool. 87, 233 (1941).

² W. EVERSOLE, Endocrinology 28, 603 (1941).


³ L. GALLIEN, C. R. Acad. Sci. (Paris) 223, 52 (1946).

⁴ T. MOHSEN, Proc. Egypt. Acad. Sci. 11, 67 (1955); C. R. Soc. Biol. (Paris) 149, 2232 (1955); Nature (London) 181, 1074 (1958).

⁵ A. STOLK, Naturwiss. 46, 274 (1959).

Purin-6-yltrimethylammonium Chloride: A New Purine Antimetabolite¹

Examination of the literature fails to disclose a single example of a purinyltrialkylammonium salt in the relatively wide spectrum of purine derivatives that have been synthesized and screened as potential antimetabolites.



It was found, in the course of a related study, that 6-chloropurine is readily transformed (78% yield) to purin-6-yltrimethylammonium chloride (I), m.p. 179–180°C², on treatment with trimethylamine in N,N-dimethyl-formamide. The toxicity and the effect of I on transplantable mouse tumor is herein described.

Acute toxicity in mice. The LD₅₀ was determined in 6 months old A₁/Sp mice. Two equal doses dissolved in sterile saline were administered intraperitoneally 6 h apart. The volume of the solution was 0.5 ml at each injection. The acute LD₅₀ was 500 mg/kg. All animals died within 12 h after the second injection. Liver, kidney, spleen, lungs, bone marrow and proximal jejunum were examined histologically in each animal. In all animals central portions of the liver lobules showed hyperemia and small areas of necrosis. No mitotic figures could be seen in the crypts of Lieberkühn. Other organs showed no abnormalities. Daily white blood counts in surviving animals showed a moderate drop in neutrophils (Figure 1).

Chronic toxicity. Six mice were treated with 150 mg/kg. The compound was administered intraperitoneally daily in 0.5 ml of sterile saline. All treated animals died between the 18th and 25th day of treatment. Before death all animals developed diarrhea. White blood counts, obtained every three days from a tail wound, showed disappearance of circulating neutrophils by the 15th day of treatment.

Histological examinations showed aplastic bone marrow in all animals, focal hemorrhagic enteritis in 5 animals and lung abscesses in 2 animals.

Effect on Ehrlich Ascites tumor. Two groups of eighteen 6-months-old Ehrlich Ascites tumor carrying male A₁/Sp mice were treated intraperitoneally daily for one week with 150 mg purin-6-yltrimethylammonium chloride. One group of animals was started on treatment four days after intraperitoneal injection of 0.2 ml of non-hemorrhagic ascites containing 4 million tumor cells. The treatment of the second group was started the day after the tumor inoculation. Equal numbers of tumor-inoculated mice of the same strain, age, and sex served as controls. The results are summarized in Figure 2. All animals in the control group had large amounts of ascites at death as did the animals dying between 5 and 9 days after tumor inoculation in the treated groups. Only two of the animals




Fig. 1

¹ Supported in part by grants CY-2903 and CY-4519 from the National Cancer Institute of the U.S.P.H.S.

² Anal. Calc. for C₈H₁₂N₆Cl: C, 44.97; H, 5.66; N, 32.78. Found: C, 44.96; H, 5.83; N, 32.98.




Fig. 2




Fig. 3

dying later in each of the treated groups had recognizable ascites at autopsy. In the second group there were 6 indefinite 'survivors' which were observed for 2 months after the experiment was terminated.

Effect on transplantable epidermoid carcinoma DC5. Twelve 6-months-old DC5 tumor bearing female MA/Sp mice were treated with 150 mg/kg purin-6-yltrimethylammonium chloride daily intraperitoneally for one week. A fragment of DC5 carcinoma was implanted under sterile conditions with a trocar into the left axilla 10 days before treatment was started. Equal numbers of animals

served as controls. At the start of treatment tumors ranged from 2 to 5 mm in diameter. The growth of the tumor was progressive in all control animals during the duration of the experiment. In all treated animals the tumor decreased in size by the time treatment was stopped and was not palpable in two animals. However, 5 days later it became apparent that the tumor had resumed its growth in all animals. The survival of the treated and untreated animals is summarized in Figure 3.

To obtain further data, the compound was submitted to the Cancer Chemotherapy National Service Center for routine screening. The compound had no effect against transplantable sarcoma 180 or against leukemia 1210. However, it was effective against carcinoma 755. The dose response curve is summarized in the Table.

Zusammenfassung. Purin-6-yltrimethylammoniumchlorid wurde synthetisiert. Es erwies sich als ein hämopoetisch toxischer Antimetabolit, wirksam gegen Ehrlich Ascites-Tumor, DC5- und Ca 755-Mäusekrebs.

J. P. HORWITZ and V. K. VAITKEVICIUS

Detroit Institute of Cancer Research and Henry Ford Hospital, Oncology Division, Detroit (Michigan), August 14, 1961.

| Daily dose (mg/kg) | Survivors | Response - Test/Control (T/C) Animal weight change (g) | Tumour weight (mg) | % (T/C) |
|--------------------|-----------|---|--------------------|---------|
| 165.00 | 06/10 | 01.7/04.4 | 10 /1478.0 | |
| 160.00 | 07/10 | 00.2/02.7 | 39.0/1716.0 | 02 |
| 80.00 | 08/10 | -00.6/02.7 | 56.0/1716.0 | 03 |
| 40.00 | 10/10 | 00.3/02.7 | 67.0/1716.0 | 03 |
| 20.00 | 09/10 | 00.2/02.7 | 94.0/1716.0 | 05 |
| 20.00 | 04/10 | -00.3/01.4 | 0.0/1499.0 | |
| 10.00 | 08/10 | -00.4/01.4 | 22.0/1499.0 | 01 |
| 5.00 | 08/10 | 00.3/01.4 | 338.0/1499.0 | 22 |

A Fungus Pigment of Novel Type, and the Nature of Fungus 'Lignin'

The presence or absence of lignin in fungi has been disputed, disagreement arising partly over the constitution of certain fungal materials and partly over the definition of a lignin. It is well-known that the fruits of many higher fungi contain hydrolysis-resistant structural materials which are of aromatic nature and may be polymeric; a major difference between these and true plant lignins is the virtual absence of methoxy groups from the typical fungal samples¹. In *Daldinia concentrica* and allied species of Ascomycetes the aromatic component of a hydrolysis-resistant structural substance in the fruits is formed by enzymic oxidation-polymerisation and is a partly quinonoid derivative of 1,8-dihydroxynaphthalene². In other species similar 'melanizing' enzymes promote the deposition of structural materials in differentiated sexual

structures, though the substrates have seldom been identified³. Materials of this kind resemble melanins rather than lignins.

The case is different with many Basidiomycetes which form superficially 'woody' sporophores. We have found that several pigmented species of this kind produce, even in vegetative mycelia, substances with the absorption spectra of oxygenated cinnamoyl derivatives. A simple example of such C_6C_3 metabolites is the methyl *p*-methoxycinnamate of *Lentinus lepideus*⁴. The more complex

¹ J. W. FOSTER, *Chemical Activities of Fungi* (New York 1949), p. 99. - K. FREUDENBERG, in K. PAECH and M. V. TRACEY, *Moderne Methoden der Pflanzenanalyse* (Berlin 1955), vol. 3, p. 499.

² J. D. BU'LOCK and D. C. ALLPORT, *J. chem. Soc.* 1958, 4090; 1960, 654.

³ P. E. BARBESGAARD and S. WAGNER, *Hereditas* 45, 564 (1959).

⁴ G. EBERHARDT, *J. Amer. chem. Soc.* 78, 2832 (1956).