

Initial Axon Collaterals from Nerve Cells in the Lateral Cervical Nucleus

The degeneration of terminal boutons in the lateral cervical nucleus (NCL) has been studied in the electron microscope after axonal transection¹. Although the lesions were made in the dorsal part of the lateral funiculus immediately below the nucleus and thus should cut off all known afferent fibres to it, there was a surprisingly small population of structurally altered boutons. A similar finding has been made in the cuneate nucleus in which only 18% of the boutons degenerated after lesions immediately below the nucleus².

To investigate if there are other afferent fibre systems to the NCL than that which was known previously³, studies have been undertaken on normal as well as on experimental material with the aid of the Golgi and the Nauta (1957) techniques, respectively. This is a preliminary note on the observation of initial axons collaterals from NCL-neurons in Golgi preparations.

Impregnation according to the rapid Golgi technique⁴ was carried out in 62 cats ranging in age from new-born to adult. Three of them, all 7 days old, were perfused with osmium tetroxide and potassium permanganate according to a method recently published⁵.

NCL-axons were found to give off collaterals at a distance of 70–130 μ from the cell body. The collaterals were very thin, leaving at a straight angle from the axons. So far only 1 collateral per axon has been observed. Most of the collaterals branched richly and terminated within the NCL with small terminal boutons or with what seemed to be free endings.

NCL has been studied with the Golgi technique before⁶ but initial axon collaterals have not been described. In this investigation, axon collaterals have been demonstrated in the perfused material only. Whether this means that perfusion is definitely superior to immersion fixation is not yet possible to decide from this material, especially when the hazardous outcome of the Golgi impregnation is considered.

The presence of initial axon collaterals fits well with what is known about the functional organization of the NCL. Neurophysiological studies have shown recurrent inhibition in the NCL after antidromic as well as orthodromic stimulation⁷. Whether the initial collaterals represent the whole number of unchanged boutons in the electron microscopical material¹ cannot be decided yet. The neurophysiological findings⁷ indicate the presence of internuncial neurones within the NCL. This would mean that at least a third group of boutons should be present in the nucleus beside those from the fibres ascending in the dorsal part of the lateral funiculus and the recurrent initial axon collaterals.

Zusammenfassung. Nach Perfusion mit Osmiumtetroxyd und Kaliumbichromat wurde das Rückenmark 7 Tage alter Katzen nach der raschen Methode von Golgi imprägniert. Es wurden initiale Axonkollateralen von Nervenzellen in Nucleus cervicalis lateralis beschrieben, was mit physiologischen Befunden übereinstimmt.

J. WESTMAN

Department of Human Anatomy, University of Uppsala (Sweden), 16th February 1967.

¹ J. WESTMAN and G. GRANT, *Acta Soc. Med. upsal.* 70, 259 (1965).

² F. WALBERG, *Expl Brain Res.* 2, 107 (1966).

³ A. BRODAL and B. REXED, *J. comp. Neurol.* 98, 179 (1953).

⁴ O. STRONG, *J. comp. Neurol.* 6, 101 (1896).

⁵ K. MOREST and R. MOREST, *Am. J. Anat.* 178, 811 (1966).

⁶ S. R. Y. CAJAL, *Histologie du système nerveux de l'homme et des vertébrés* (Consejo superior de investigaciones científicas, Madrid 1952), vol. I, p. 986; T. A. LEONTOVICH and G. P. ZHUKOVA, *J. comp. Neurol.* 121, 347 (1963); H. HA and C.-N. LIU, *Expl Neurol.* 8, 318 (1963).

⁷ G. GORDON and M. G. M. JUKES, *J. Physiol., Lond.* 169, 28 (1963); D. F. HORROBIN, *Q. Jl exp. Physiol.* 57, 351 (1966).

The Toxic Action of *Bacillus thuringiensis* 'Exotoxin' on *Drosophila* Reared in Yeast-Containing and Yeast-Free Media

A prepurified preparation of the so-called 'exotoxin' of *Bacillus thuringiensis* was prepared by differential precipitation with ethanol after the method of BENZ¹. Its toxic activity was bioassayed with larvae of *Drosophila* on two basically different rearing media.

Medium A was a corn-yeast-agar medium (water 750 ml, agar 9 g, sugar 50 g, corn semolina 100 g, dry yeast 20 g), while medium B was the synthetic medium C of SANG². Our third medium C was medium B with 2% of dry yeast added. After the media were cooked 0.66 ml of a 20% solution of nipagin in ethanol was added per 100 ml of medium and mixed thoroughly. Polystyrene beakers with a capacity of 200 ml were used for all tests. 0.5 ml of a 2.2% solution of streptomycin, and 2 ml of different dilutions of 'exotoxin' or water were pipetted into each beaker, and, in the case of the media B and C an additional 0.5 ml of a vitamin mixture. 30 ml of warm (not hot) medium were added per beaker and well mixed with the liquid before the medium solidified. Not all

brands of casein can be used for the preparation of synthetic *Drosophila* medium. We tested vitamin-free caseins manufactured by Nutritional Biochemical Co., Merck, and Fluka. Only the first preparation was good (maximal control mortality 6%), while the other two brands were toxic for *Drosophila* larvae (control mortalities of 45 and 100% respectively).


Drosophila eggs were collected from culture bottles, placed on fine meshed gauze and washed several times with water of 25°C in order to remove yeast and other impurities. The washed eggs were then kept on wet filter paper in a Petri dish. Fifty freshly hatched larvae were placed on the medium of each beaker. The beakers were closed with perforated polystyrene lids and kept at 24–25°C and 80% relative humidity. Flies that eclosed were recorded, and the results were corrected by ABBOT's formula.

The Figure presents results of such tests. They indicate a marked difference between the mode of action of the

¹ G. BENZ, *Experientia* 22, 81 (1966).

² J. H. SANG, *J. exp. Biol.* 33, 45 (1956).

'exotoxin' in the corn-yeast-agar medium (curve A) and the synthetic medium (curve B). The medium lethal concentration (LC_{50}) is much lower in the yeast-free synthetic



Probit regression lines of mortality of *Drosophila* reared on 3 media with different concentrations of 'exotoxin'. (A) corn-yeast-agar medium; (B) synthetic medium C of SANG²; (C) SANG's medium plus 2% of dry yeast. Each point represents 100 test larvae. The line of the 50% mortality level indicates the different LC_{50} values at the points where it crosses the probit regression lines. Ordinate: instead of probits, % mortality is given by a probability distribution scale. Abscissa: log 100 times concentration of 'exotoxin' preparation in medium.

medium, and the slope of curve B is steeper than that of curve A. These differences in slope and LC_{50} are mainly due to the presence or absence of dry yeast in the media. This is demonstrated by curve C which represents results of tests with medium C, i.e. synthetic medium to which 2% of dry yeast was added. This addition reduces the slope of curve B to a value which is practically identical with that of curve A, and the LC_{50} is also very similar to that with the corn-yeast-agar medium.

The results show that yeast reduces the toxic action of the 'exotoxin'. Bioassays on media that do not contain yeast are therefore more sensitive. As a consequence it must be postulated that bioassays for 'exotoxin' should either be made with yeast-free media, or with standardized media containing a defined proportion of yeast. A more detailed analysis of the yeast effect will be published elsewhere³.

Zusammenfassung. Die toxische Wirkung des sogenannten «Exotoxins» von *Bacillus thuringiensis* auf *Drosophila* wurde geprüft: (A) in Mais-Trockenhefe-Agar-Medium, (B) synthetischem Medium C nach SANG² und (C) synthetischem Medium plus 2% Trockenhefe. In hefefreiem Medium ist die LC_{50} bedeutend niedriger und die Probitkurve viel steiler als in hefehaltigen Medien. Hefe reduziert also die toxische Wirkung des «Exotoxins» beträchtlich.

G. BENZ and J.-M. PERRON⁴

Entomologisches Institut, Eidg. Technische Hochschule, Zürich (Switzerland), 7. Juni 1967.

³ J.-M. PERRON and G. BENZ, *J. Invertebrate Pathol.*, in print.

⁴ Supported by Fellowship No. 1560 of the National Research Council of Canada. - Present address: Département de Biologie, Université Laval, Québec, Canada.

Separation of 'Soluble' Immunoprecipitating Antigens Originating from Rabies Virus Infected Cells by Chromatography on Ecteola Cellulose and by Gel Filtration

Rabies virus has been successfully purified by chromatography on Ecteola cellulose^{1,2}. Preliminary experiments have revealed that most of the 'soluble' antigens can be eluted from Ecteola cellulose using solutions of lower ionic strength than are required to elute the whole virus³. The present communication will present in more detail the chromatographic technique of separating the 'soluble' antigens originating from the infected cells and will substantiate the feasibility of their purification by gel-filtration on Sephadex G-200.

The crude 'soluble' antigen preparation was obtained from infected tissue culture fluids by zinc acetate precipitation and subsequent removal of virus from the redissolved sediment through high speed centrifugation³. Prior to chromatography, this preparation was extensively dialyzed against 0.01 M Tris-(hydroxymethyl) aminoacetate buffer (TB) pH 7.0. Six ml of the antigen

preparation was applied to a column (dia. 1.6 · 27 cm) of Ecteola cellulose (Serva, Heidelberg). For elution, a continuous gradient of increasing NaCl concentration (220 ml TB-220 ml 0.45 M NaCl in TB) was used. Fractions (5 ml) were collected and analyzed for protein⁴ and precipitating antigen⁵ using 0.75 ml aliquots. Fluorescein isothiocyanate-labeled antirabies γ -globulin (Baltimore Biological Labs.) mixed with non-infected tissue culture fluid material was used for the fluorescent precipitin test.

¹ J. B. THOMAS, A. S. RICKER, G. M. BAER and R. K. SIKES, *Virology* 25, 271 (1965).

² A. R. NEURATH, T. J. WIKTOR, M. V. FERNANDES and H. KOPROWSKI, unpublished data. The preliminary experiments were performed at the Wistar Institute of Anatomy and Biology, Philadelphia, Pa. The presented results were obtained while A.R.N. worked at Wyeth Laboratories, Philadelphia, Pa. Rabies virus infected tissue culture material was obtained from the Wistar Institute.

³ A. R. NEURATH, T. J. WIKTOR and H. KOPROWSKI, *J. Bact.* 92, 102 (1966).

⁴ O. H. LOWRY, N. J. ROSEBROUGH, A. L. FARR and R. J. RANDALL, *J. biol. Chem.* 193, 265 (1951).

⁵ A. R. NEURATH, *Z. Naturf.* 20b, 974 (1965).