

Contraction of Frog Stomach Muscle and Frog Heart in Electrolyte-Free Solutions

Frog stomach muscle when frequently washed with half-isotonic (0.112 M) solution of sucrose, and frog heart perfused with similar solution lose nearly all sodium in 1 h^{1,2}, but exhibit spontaneous contractions³⁻⁶, with conducted action potentials for several hours⁷⁻¹⁶. They lose about 90% of their sodium in the first 15 min, and in 1 h the sodium content of both becomes less than 0.05 mM/kg of wet muscle². There is however a striking difference between the mechanical performance of the stomach muscle and that of the heart. The stomach muscle shows spontaneous contractions at all temperatures (experimental range 15–35°C) for about 24 h. The frog heart beats for about 24 h at 5°C, for about 2 h at 25°C¹⁶, and at high temperatures (30–35°C) it passes into contracture and does not beat at all.

There is a remarkable difference between the behaviour of adenosinetriphosphate (ATP) in sucrose-soaked frog stomach muscle and in frog heart. The ATP content of frog stomach muscle does not differ in any way whether the muscle is soaked in saline or in sucrose solution¹⁷. But in frog heart both ATP and creatine phosphate fall in sucrose solution, more at 25°C than at 5°C, and the fall correlates with the decline in the force of contraction¹⁶.

These experiments therefore suggest that sodium in muscle has an important metabolic function of preserving the ATP and failure of function in sodium-free solutions is most likely due to loss of ATP rather than to any direct action on the excitability mechanism.

Zusammenfassung. Der Verlust der Reizbarkeit bei Froschherz- und Magenmuskulatur auf verschiedenen

Temperaturstufen verläuft parallel mit einem Adenosin-triphosphat-Verlust, was andeutet, dass das Nichtfunktionieren in natriumfreier Lösung von einem Adenosin-triphosphat-Verlust abhängig ist.

I. SINGH and S. I. SINGH

Department of Physiology, Medical College, Agra (India), 15 May 1967.

- ¹ E. BOZLER, *Am. J. Physiol.* **199**, 299 (1960).
- ² I. SINGH and N. V. RAJU, *Experientia* **21**, 17 (1965).
- ³ S. I. SINGH and I. SINGH, *Proc. Indian Acad. Sci.* **78**, 56 (1943).
- ⁴ I. SINGH, *Curr. Sci.* **13**, 231 (1944).
- ⁵ I. SINGH, K. B. SEHRA and S. I. SINGH, *Proc. Indian Acad. Sci.* **27**, 259 (1945).
- ⁶ I. SINGH, K. B. SEHRA and S. I. SINGH, *Curr. Sci.* **14**, 152 (1945).
- ⁷ I. SINGH and J. BHATT, *Proc. Ind. Acad. Sci.* **46**, 211 (1957).
- ⁸ I. SINGH and A. K. ACHARYA, *Proc. Indian Acad. Sci.* **46**, 285 (1957).
- ⁹ R. L. KOLODNY and W. G. VAN DER KLOOT, *Nature* **190**, 786 (1961).
- ¹⁰ W. G. VAN DER KLOOT and N. S. RUBIN, *J. gen. Physiol.* **46**, 35 (1962).
- ¹¹ I. SINGH, *Am. J. Physiol.* **203**, 422 (1964).
- ¹² I. SINGH, *Experientia* **20**, 33 (1964).
- ¹³ I. SINGH, *Archs int. Physiol.* **71**, 361 (1963).
- ¹⁴ I. SINGH, *Archs int. Physiol.* **72**, 378 (1964).
- ¹⁵ I. SINGH and S. I. SINGH, *Experientia* **22**, 165 (1966).
- ¹⁶ W. G. VAN DER KLOOT and B. J. DANE, *J. gen. Physiol.* **48**, 199 (1966).
- ¹⁷ T. J. SARMA, personal communication.

3,4-Dihydroxycinnamic Acid, an Antithiamine Factor of Fern

According to our previous publications, fern (*Pteridium aquilinum*) contains antithiamine substances^{1,2}. These early experiments have shown that the active agent was a thermostable, water-soluble, small molecule, which moved in an electric field to the anode^{3,4}.

Later a good purification of the antithiamine agent was achieved, although the substance could not be isolated in pure form. This antithiamine factor, called Hydrolysate II, was a red-brown amorphous substance, free of N, P, S and halogenes⁵. The specific antithiamine activity of this factor was confirmed by the experiments with the single nerve fibre of the frog by VON MURALT and PETROPOULOS⁶ and PETROPOULOS⁷.

These experiments have recently been resumed. The antithiamine activity was determined by the thiochrome and microbiological methods. The purification process, which has been substantially altered, is summarized in Figure 1. The antithiamine activity of the dark-brown, oily ethyl acetate phase increased, while the activity of the water phase remained unchanged. The most active fraction of the column chromatography (I) was rechromatographed.

The purification effect of the column chromatography has been established by polyamid thinlayer chromato-

grams. Seventeen components could be separated from the ethyl acetate phase (see also Figure 2). One of the components (shaded spot) possessed the highest antithiamine activity ($R_f = 0.47$). This could be isolated as a uniform substance.

This compound is a microcrystalline substance with a melting point 190–192°C. Its antithiamine activity is very high, namely 1800 µg/mg, according to the thiochrome method (time effect curve) and 2300 µg/mg determined microbiologically. The activity is pH and temperature-dependent, being higher in slightly alkaline solutions (pH 7.8), than at pH 6.0.

The substance is thermostable; the antithiamine activity remained unchanged after boiling 2 h by reflux. It is soluble in water and ether and easily soluble in ethanol,

- ¹ J. C. SOMOGYI, *Int. Z. Vitaminforsch.* **21**, 341 (1949).
- ² J. C. SOMOGYI and A. V. MURALT, *Helv. Physiol. Acta*, **7**, C 56 (1949).
- ³ J. C. SOMOGYI, *Helv. Physiol. Acta*, **9**, C 79 (1951).
- ⁴ J. C. SOMOGYI, *Antianeurin-Faktoren* (Hans Huber Verlag 1952).
- ⁵ J. C. SOMOGYI and A. KOLLER, *Int. Z. Vitaminforsch.*, **29**, 234 (1959).
- ⁶ A. V. MURALT and S. V. PETROPOULOS, personal communication (1959).
- ⁷ S. V. PETROPOULOS, *J. Cell Comp. Physiol.*, **56**, 7 (1960).