

## Partial Complementation in *Aspergillus niger* Heterocaryon

Auxotrophic mutants require essential growth factors for normal growth on minimal medium. However, non-allelic mutants can usually complement each other in heterocaryons or diploids<sup>1-3</sup> under favorable conditions and show the normal phenotypic features of the wild type, or approach them, on the same medium.

Rarely 2 genetically dissimilar nuclei in 1 unit of cytoplasm do not complement each other but only do so if combined in a diploid nucleus<sup>2,3</sup>. In my case it has been found that 2 auxotrophic mutants will complement each other in a heterocaryon by showing growth on minimal medium which approaches wild type but it will not conidiate at all.

Yellow mutant  $Y_1$  of *A. niger* was produced by exposure to 2537 Å UV-light and 2 auxotrophic mutants requiring histidine (hist) and hypoxanthine (hypox) respectively were induced in the same way. The optimal supplementation to minimal medium for histidine and hypoxanthine was 0.3 mg/ml and 0.1 mg/ml respectively. Both mutants grew alone and responded with similar type of growth at the same age and temperature but the hist  $Y_1$  mutant did not conidiate as heavily as hypox  $Y_1$  mutant. The mean spore size of the 3 strains did not differ significantly. A forced heterocaryon was made between the 2 auxotrophic mutants by taking a heavily sporing suspension of each mutant and putting equal amounts in a flask of liquid minimal medium supplemented with their growth factors. The flasks were incubated for 5 days at 25°C. A mesh of submerged mycelium developed in the flask. It was isolated, spread on the surface of minimal agar and incubated for nearly a week at 30°C. At various places on the plate mycelium grew vigorously after incubation. Such mycelium was cut out, transferred to minimal medium and incubated for a further 3-5 days at 25°C. Despite effective hyphal growth on subsequent incubation for at least 20 days no conidia were seen.

Confirmation that the mycelium was heterocaryotic was provided by transfer to complete medium. After incubation heavy yellow conidiation developed; single spore isolations were tested for nutritional requirements and it was shown that both hist and hypox nuclei were present.

Forty milligrams of D-Camphor flower (natural) was sterilized in a petri dish for 5 lbs at 10 min and then sterilized minimal medium was poured on the top<sup>4</sup>. Plates so prepared had the heterocaryon inoculated on them and were left under a glass cover at room temperature for a

month. Only 2 plates showed conidia at a few points; they were transferred to minimal medium, germinated satisfactorily, grew and conidiated. These conidia were presumed to be diploid, were isolated and tested by single spore culture, the measurement of the conidial size and by the treatment of the conidia with *p*-fluorophenyl-alanine<sup>5</sup>. The result showed that these were true diploid. The volume of the individual conidium was not quite double than that of the parents conidia<sup>4</sup>.

The partial complementation in heterocaryon observed in these experiments could be caused by any of the following possibilities.

It might possibly be due to a disproportion of nuclear types in the heterocaryon which would lead to insufficient nutritional cooperation but, as soon as sufficient nuclei fuse together to make diploids, they would overcome the nutritional deficiency in a limited area and a diploid could resume full growth with conidiation. A second possibility might be a dilution effect on a secondary gene product arising by interaction in the cytoplasm. Here again in diploid nuclei due to the physically closer association of the genetic material, their products could interact before they had become diluted in the cytoplasm<sup>6</sup>.

**Zusammenfassung.** Erzwungene Heterokaryen-Vereinigung zwischen Hypoxanthin und Histidin abhängiger Mutanten von *Aspergillus niger*, die auf minimalem Nährboden ohne Sporenproduktion wachsen. Das Phänomen wird als teilweise Ergänzung bezeichnet und als Gleichgewichtsstörung der Kerntypen im Heterokaryon oder als Verdünnungseffekt eines Genproduktes zweiter Ordnung im Zytoplasma gedeutet.

I. PRASAD

Department of Microbiology, Medical College of Virginia, Richmond (Virginia 23219, USA), 25 November 1968.

<sup>1</sup> G. PONTECORVO, J. A. ROPER and E. FORBES, J. gen. Microbiol. 8, 198 (1953).

<sup>2</sup> D. APIRION, Genetics 53, 935 (1966).

<sup>3</sup> C. F. ROBERTS, Aspergillus Newsletter 3, 6 (1962).

<sup>4</sup> J. A. ROPER, Experientia 8, 14 (1952).

<sup>5</sup> P. LHOAS, Nature 190, 744 (1961).

<sup>6</sup> L. A. CASSELTON and D. LEWIS, Genet. Res. 9, 63 (1967).

## Sex Ratio Alteration in Litter of Parents Submitted to Experimental Neurosis

There are some data concerning the effect of stressful situations on sex distribution of children born in the 9-12 months following the stress. SNYDER<sup>1</sup> observed that military pilots of high performance have much more girls than boys after combat-fighting; the sex ratio diminished from 105.37 of controls to 59.32, which means that nearly  $\frac{2}{3}$  of the children were females. (Sex ratio = number of males born related to 100 females born in a given population.) According to these data, it seemed worth while to test the effect of anxiety on sex ratio in animals.

Wistar albino rats, both males and females, were submitted to experimental neurosis lasting 3 months, which was accompanied by overt signs of anxiety. As anxiety

always goes side-by-side with a large output of adreno-medullar hormones<sup>2</sup>, both epinephrine and norepinephrine were administered to rats for 3 months, in order to test whether the effect of experimental neurosis might be attributed to the output of these hormones. This supposition seemed to be possible, because norepinephrine has an effect on progeny: it significantly reduces the litter size<sup>3</sup>. As control animals partly conditioned but not


<sup>1</sup> R. G. SNYDER, Hum. Biol. 33, 1 (1961).

<sup>2</sup> L. LEVI, Psychosom. Med. 27, 80 (1965).

<sup>3</sup> A. SAI-HALÁSZ, Experientia 21, 155 (1965).

neurotized rats served, partly ones treated with Ringer solution. Both parents, i.e. father and mother were submitted to the same experimental procedure, which has been published in detail elsewhere<sup>4,5</sup>. In total 156 deliveries were observed, 102 cases in the experimental, and 54 in the control groups. Both litter size and sexual distribution was checked. The numerical results are presented in the Table.

The results show that while the litter size, i.e. the number of young, is reduced only in the norepinephrine group, the sex distribution is in the range of normal limits both in the injected and in the control groups, i.e.



Number of females born related to 100 males born.

Group	No. of observations	No. of young	Males	Females	Sex ratio
Neurotized	36	273	107	166	64
Epinephrine	32	237	125	112	112
Norepinephrine	34	194	99	95	104
Controls	54	397	193	204	95

Number of young born and their sex distribution after experimental neurosis, epinephrine and norepinephrine treatment lasting for 3 months.

the sex ratio is between 95 and 112. In the experimental neurotic group, however, it diminishes to 64, which means a high predominance of females over males in the next generation. This result is in accordance with the data observed in men by SNYDER<sup>1</sup>. The excess of females is even more striking when the number of females born is related to 100 males born (see Figure).

As for the mechanism of action of the above effect, it cannot be attributed to a larger output of catecholamines; as demonstrated, adrenomedullar hormones have no effect on sex ratio, as experimental neurosis has. So the mechanism of this effect is as yet unknown.

In mammalian populations, the propagation of species is related to the number of females. So the above results, i.e. the excess of females after the neurotic situation of parents, can be seen as one feature of the biological principle of SPENCER<sup>6</sup>. He established more than 100 years ago that those species which are rather much pursued and chased and therefore live in a stressful situation, have a more enhanced propagation, and it is this way they secure their survival.

**Zusammenfassung.** Sex ratio, d.h. die numerische Verteilung der Geschlechter untereinander, wurde bei Ratten nach einer dreimonatigen experimentellen Neurose beider Eltern registriert. Die Sex ratio sank bis 64, was eine starke Überzahl zugunsten der Weibchen unter den Abkömmlingen bedeutet: auf 100 Männchen fallen 156 Weibchen. Diese Reaktion kann als eine Form des SPENCERschen biologischen Gleichgewichtsprinzips bewertet werden: In Stress-Situationen wird die Erhaltung der Art gewährleistet durch die Tatsache, dass die Nachkommenschaft mehr weibliche Individuen aufweist.

A. SAI-HALÁSZ

*Centr. Neurosurgical Institute,  
Budapest XIV (Hungary), 17 October 1968.*

<sup>4</sup> A. SAI-HALÁSZ, Proc. 3rd IFA Congr. (Excerpta Med., Amsterdam 1961), p. 995.

<sup>5</sup> A. SAI-HALÁSZ, Kand. Ért. Acad. hung. 59 (1966).

<sup>6</sup> H. SPENCER, *Die Prinzipien der Biologie* (Schweizerbart, Stuttgart 1877), vol. 2.

### Action de la (+)-catéchine sur *Pseudomonas fluorescens*

*Pseudomonas fluorescens* est une bactérie mobile, gram négatif, vivant en aérobiose. Elle sécrète un pigment jaune à fluorescence verte qui s'accumule dans le milieu de culture. Ce pigment, de nature pyrrolique, présente un maximum d'absorption entre 390 et 430 nm; il a pour origine un précurseur endogène à fluorescence bleue. Selon LENHOFF et al.<sup>1</sup>, cette bactérie possède deux sortes de respiration terminale: a) cytochromique, cyanosensible; b) flavinique, cyanorésistante. Les travaux de GOUDA et GREPPIN<sup>2,3</sup> montrent que l'orientation qualitative et quantitative de cette alternative respiratoire est essentiellement réglée par le rapport de la pression partielle de l'oxygène à la concentration en substrat nutritif, mis à la disposition de la bactérie; d'autre part, la vitesse de variation de ce

rapport en fonction du nombre de germes, joue un rôle important dans l'induction respiratoire terminale.

L'alternative respiratoire a des conséquences importantes sur la disponibilité en énergie (la phosphorylation oxydative étant plus efficace dans le système terminal cytochromique) et de ce fait sur la physiologie et la mor-

<sup>1</sup> H. M. LENHOFF, D. J. D. NICHOLAS et N. O. KAPLAN, J. biol. chem. 220, 983 (1956).

<sup>2</sup> S. GOUDA et H. GREPPIN, C. r. Soc. Phys. Hist. Nat. 1, 3 (1966).

<sup>3</sup> S. GOUDA et H. GREPPIN, en préparation.