

### Alteration in the SAT-Chromosome of *Allium cepa* Affecting the Size of the Nucleolus

The nucleolus is formed in a generally well-defined locus of the chromosome<sup>1-3</sup> cytologically discernible as a secondary constriction called the nucleolar organizer<sup>2</sup>. An important question in the study of the relationship between SAT-chromosomes and nucleolus is the correlation between alterations taking place in the organizing area and the morphology and physiology of the nucleolus.

In the course of cytological studies of *Allium cepa* we observed a bulb which showed marked differences of size between the nucleoli of each nucleus. By means of a

silver impregnation technique<sup>4</sup>, it was observed that all the meristematic cells which had 2 nucleoli were found to have one of normal size, while the other had a diameter

<sup>1</sup> E. HEITZ, *Planta* 12, 775 (1931).

<sup>2</sup> B. McCLINTOCK, *Z. Zellforsch. mikrosk. Anat.* 21, 294 (1934).

<sup>3</sup> W. BEERMANN, *Chromosoma* 11, 263 (1960).

<sup>4</sup> M. E. FERNÁNDEZ-GÓMEZ, J. C. STOCKERT, J. F. LÓPEZ-SÁEZ and G. GIMÉNEZ-MARTÍN, *Stain Technol.* 44, 48 (1969).




Fig. A and B. Meristematic cells showing the different-sized nucleoli. Silver stain, phase contrast. C. Metaphase of the anomalous bulb. Only 1 of the SAT-chromosomes (arrows) presents the satellite. D. Nucleolar chromosomes of another metaphase showing the absence of satellite in one of them (arrow). E. Normal metaphase of *Allium cepa*.

between 40 and 50% smaller than the normal one (Figure A and B). The symmetrical position of these 2 different nucleoli in sister-nuclei seemed to point to the probability that each was organized by a different, specific chromosome.

With a view to analysis of the chromosomes, a number of specimens were prepared in accordance with the technique of TJIO and LEVAN<sup>5</sup>. All the metaphases showed the 16 chromosomes typical of the onion, but in the pair normally possessing satellites<sup>6-8</sup>, only one of the chromosomes possessed one (Figure C and D), a form of heterozygosis similar to that already observed by GIMÉNEZ-MARTÍN<sup>9</sup>, in the satellite pair of *Allium cepa*. Bulbs with nucleoli of the same size were found to have the normal SAT-chromosomes (Figure E). The presence of different-sized nucleoli is therefore correlated with the absence of a satellite or the corresponding secondary constriction in one of the chromosomes of *Allium cepa*.

This absence of the secondary constriction might be due simply to a physiological condition<sup>10</sup>, since there is evidence that the morphology of the secondary constrictions is controlled by genes within the chromosome itself<sup>11</sup>. But many studies have demonstrated the connection between the amount of chromosome material responsible for organizing the nucleolus and the volume of the nucleolus<sup>2,3,12,13</sup> or the number of ribosomal cistrons present<sup>14,15</sup>. The possibility of deletion of the secondary nucleolar constriction<sup>16</sup> in the anucleolate mutant in *Xenopus laevis*<sup>17</sup> is demonstrated by the fact that in this mutant the ribosomal cistrons are not merely inactivated but absent<sup>15</sup>. We might interpret the case we are presenting as a partial deletion of 1 nucleolar organizer, of which the cytological expression would be the difference in size between the 2 nucleoli, due to the heterozygosis.

**Resumen.** Se describe el hallazgo de un bulbo de *Allium cepa* en el cual los dos nucleolos de cada núcleo muestran una clara diferencia de tamaño. El nucleolo menor posee aproximadamente la mitad del diámetro del nucleolo normal. Esta alteración de la morfología nucleolar se correlaciona con la ausencia de satélite en uno de los dos cromosomas normalmente satelizados.

J. C. STOCKERT<sup>18</sup>, J. M. SOGO,  
J. L. Díez and G. GIMÉNEZ-MARTÍN

*Departamento de Citología,  
Instituto de Biología Celular, C.S.I.C.,  
Madrid 6 (Spain), 17 March 1969.*

- <sup>5</sup> J. TJIO and A. LEVAN, An. Estac. Exp. Aula Dei 2, 21 (1951).
- <sup>6</sup> A. LEVAN, Hereditas 15, 347 (1931).
- <sup>7</sup> A. LEVAN, Hereditas 16, 257 (1932).
- <sup>8</sup> M. R. AHUJA, J. Hered. 49, 247 (1958).
- <sup>9</sup> G. GIMÉNEZ-MARTÍN, Fiton 16, 129 (1961).
- <sup>10</sup> H. J. BARR, Natn. Cancer Inst. Monogr. 23, 411 (1966).
- <sup>11</sup> R. P. DONAHUE, W. B. BIAS, J. H. RENWICK and V. A. McKUSICK, Proc. natn. Acad. Sci. U.S.A. 61, 949 (1968).
- <sup>12</sup> M. LIN, Chromosoma 7, 340 (1955).
- <sup>13</sup> A. LONGWELL and G. SVIHLA, Expl. Cell Res. 20, 294 (1960).
- <sup>14</sup> F. M. RITOSSA and S. SPIEGELMAN, Proc. natn. Acad. Sci. U.S.A. 53, 737 (1965).
- <sup>15</sup> M. L. BIRNSTIEL, H. WALLACE, J. L. SIRLIN and M. FISCHBERG, Natn. Cancer Inst. Monogr. 23, 431 (1966).
- <sup>16</sup> J. KAHN, Q. Jl. microsc. Sci. 103, 407 (1962).
- <sup>17</sup> T. R. ELSDALE, M. FISCHBERG and S. SMITH, Expl. Cell Res. 14, 642 (1958).
- <sup>18</sup> The work reported in this paper was undertaken during the tenure of a Research Training Fellowship awarded by the International Agency for Research on Cancer.

### Chromosomes of Three Asian Mammals: *Meriones meridianus* (Rodentia: Gerbillinae), *Spermophilopsis leptodactylus* (Rodentia: Sciuridae), *Ochotona rufescens* (Lagomorpha: Ochotonidae)

Chromosomal analysis of many mammalian species during the last decade has demonstrated the potential usefulness of cytological characters in taxonomic investigation. Acquisition of 3 Asian mammals now provides additional information concerning chromosomal relationships within the rodent genus *Meriones*, between the 2 families of Lagomorpha, and between Asian and North American ground squirrels.

**Materials and methods.** The following specimens were studied: (1) *Meriones meridianus* Pallas, Iran: 21 km west of Sarakhs, one male; (2) *Ochotona rufescens* Gray, West Pakistan: 24 km S. Quetta (Chiltan Mtn.) 4 females and 3 males; (3) *Spermophilopsis leptodactylus* Lichtenstein, Iran: 24 km south of Sarakhs, 1 female and 2 males.

Chromosomes were analyzed from cell suspensions of femoral bone marrow<sup>1</sup>. Karyotypes were arranged in pairs according to size and centromere position (metacentric, submetacentric, acrocentric). Because morphological differentiation was difficult in *O. rufescens*, acrocentrics (telocentrics) and subtelocentrics were grouped together and similarly the biarmed chromosomes (metacentrics and submetacentrics) were grouped together. The fundamental number of chromosome arms (FN) is computed by counting biarmed autosomes as 2 arms and acrocentric autosomes as 1 arm; in *Meriones* the sex chromosome arms are included to facilitate comparison with prior publications.

**Results.** The diploid chromosome number (2n) of *Meriones meridianus* is 50 and the karyotype contains 12 metacentric, 14 submetacentric, and 22 acrocentric autosomes (Figure 1). A large submetacentric and a medium-sized submetacentric comprise the sex chromosomes and they are designated X and Y respectively because sex chromosomes of identical morphology have been found in 4 other species of *Meriones*<sup>1,2</sup>. The FN is 78.

*Ochotona rufescens* has a 2n of 60 and a karyotype containing 24 biarmed and 34 acrocentric and subtelocentric autosomes, a medium-sized submetacentric X and a minute, dot-like Y chromosome (Figure 2). The FN is 82.

*Spermophilopsis leptodactylus* has a 2n of 38. The autosomes are composed of 14 metacentrics, 20 submetacentrics, and 2 acrocentrics with prominent satellites (Figure 3). A medium-sized submetacentric X and a very small biarmed Y constitute the sex chromosomes (Figure 3). The FN is 70.

**Discussion.** 9 previously studied species of *Meriones* displayed 2n's of 40-72 and FN's of 72-78<sup>1-6</sup>. The chro-

- <sup>1</sup> C. F. NADLER and D. M. LAY, Z. Säugetierk. 32, 285 (1967).
- <sup>2</sup> D. M. LAY and C. F. NADLER, Cytogenetics 8, 35 (1969).
- <sup>3</sup> R. MATTHEY, Rev. Suisse Zool. 60, 225 (1953).
- <sup>4</sup> R. MATTHEY, Caryologia, Pisa 6, 1 (1954).
- <sup>5</sup> R. MATTHEY, Säugetierk. Mitt. 5, 145 (1957).
- <sup>6</sup> A. ZAHAVI and J. WAHRMAN, Mammalia 21, 341 (1957).