

enzymes seem to consist of isoenzymes, as was recently shown for pneumococcal neuraminidase by other methods<sup>8</sup>.

Attempts to characterize the neuraminidases of para-influenza viruses by isoelectric focusing were unsuccessful because a considerable loss in enzyme activity was encountered in the purification process. Similar difficulties have been previously reported<sup>9</sup>.

The results reported here explain why *V. cholerae* neuraminidase is less sensitive than the *Cl. perfringens* enzyme to inhibition by polyanions at pH 5, and why influenza virus neuraminidases are much more readily inhibited even at pH values higher than 5<sup>10</sup>, under conditions when they must have a net positive electric charge as shown here. The isoelectric point of influenza virus neuraminidase is relatively high, which is consistent with the low electrophoretic mobility (at pH 8.9) of the enzyme compared with that of other influenza virus envelope proteins<sup>4,11</sup>.

**Zusammenfassung.** Die isoelektrischen Punkte der Neuraminidasen von *Vibrio cholerae* (pH 4.80) und *Clostridium*

*perfringens* (pH 4.95) wurden mittels isoelektrischer Fokussierung bestimmt. Neuraminidasen zweier verschiedener Influenzaviren (A<sub>2</sub>/Aichi/68 und B/Mass/66) wurden entsprechend analysiert. Die Virus-Neuraminidasen waren heterogen und hatten wesentlich höhere isoelektrische Punkte als die bakteriellen Enzyme.

A. R. NEURATH, R. W. HARTZELL  
and B. A. RUBIN

Wyeth Laboratories, Inc., Department of Research and Development, P.O. Box 8299, Philadelphia (Pennsylvania 19101, USA), 25 May 1970.

<sup>8</sup> S. W. TANENBAUM, J. GULBINSKY, M. KATZ and S. C. SUN, Biochim. biophys. Acta 198, 242 (1970).

<sup>9</sup> R. DRZENIEK, J. T. SETO and R. ROTT, Biochim. Biophys. Acta 128, 547 (1966).

<sup>10</sup> R. DRZENIEK, Nature, Lond. 211, 1205 (1966).

<sup>11</sup> W. G. LAVER and R. C. VALENTINE, Virology 38, 105 (1969).

## The Prosthetic Group of the Green Chromoprotein of *Patella* Ova

The colour of the mature ova of limpets varies from dark brown to olive-green<sup>1</sup>. A green chromoprotein has been isolated from ripe olive-green ovaries of *Patella caerulea*<sup>2</sup>, but the nature of the prosthetic group has not been elucidated<sup>2,3</sup>. The present communication describes the identification of the prosthetic group as a close derivative of chlorophyll *a*.

*Patella vulgata* were collected at Roscoff in France and *Patella caerulea* were collected in Malta. The main green pigment in ova of *Patella vulgata* was isolated in the form of a methyl ester which proved to be methyl phaeophorbide *a*. The ripe ova were dried in vacuo and extracted with methanolic HCl (6–8%, w/w) for 72 h at 0°C. The dark green extract was transferred to chloroform and submitted to thin-layer chromatography on silica gel G. A fast moving green pigment (R<sub>f</sub> 70–80) was isolated with chloroform-*n*-butanol (190:10, v/v) and purified with carbon tetrachloride-methyl acetate (100:50, v/v) as solvents. The pigment showed absorption maxima at 274, 326, 418, 475, 512, 542, 565, 612 and 668 nm and a fluorescence maximum at 670 nm. It behaved as an authentic sample of methyl phaeophorbide *a* in thin-layer chromatography.

Green chromoprotein was isolated from the ova of *Patella caerulea* as described previously<sup>2</sup>. The chromoprotein showed absorption maxima at 395, 585 and 625 nm in 0.1 M ammonium bicarbonate (pH 8). A 0.5% solution of the chromoprotein in 0.1 M ammonium bicarbonate was digested with trypsin (0.1 mg/ml) at 15°C for 48 h. The solution became yellow with a greenish tinge and showed a large absorption maximum at 410 nm, two smaller maxima at 610 and 665 nm, and an inflexion at 540 nm. A pigment with similar absorption was obtained by refluxing lyophilized chromoprotein with boiling methanol for 8 h. This pigment was esterified with 14% boron trifluoride in methanol, transferred to chloroform and purified by thin-layer chromatography on silica gel G with carbon tetrachloride-methyl acetate (100:50, v/v) as solvent. A fast moving yellowish green pigment (R<sub>f</sub> 80) was obtained. The pigment showed absorption maxima at 276, 319, 355, 385, 416, 505, 535, 560, 605 and 660 nm in dioxan. The absorption spectrum showed the features of a chlorin<sup>4</sup> and the IR-spectrum showed a peak in the

C=C stretching region at about 1625 cm<sup>-1</sup> which is found in chlorins<sup>5</sup>. The pigment was identified as chlorin *a*<sub>6</sub> methyl ester in view of the preparation of methyl phaeophorbide *a* from the ova. It is probable that the isocyclic ring was broken during the extraction of the pigment from the chromoprotein with boiling methanol.

The present findings would appear to show that the prosthetic group of the green chromoprotein of *Patella* ova is a close derivative of chlorophyll *a*, namely, phaeophytin *a*<sub>5</sub> or phaeophorbide *a*<sub>6</sub>. This provides a simple explanation for the reversible yellow colour change of the chromoprotein at alkaline pH (see <sup>2</sup>) in terms of enolization of the hydrogen atom at carbon C-10 to the carbonyl group at carbon C-9 as in the ether-methanolic KOH phase test for chlorophylls<sup>6</sup>.

**Résumé.** Les ovaires des Patelles *Patella vulgata* et *Patella caerulea* contiennent un chromoprotéide vert qui a été précédemment étudié. Dans le présent travail, le groupe prosthétique de ce pigment est identifié. Il s'agit d'un tétrapyrrole dont la structure est proche de celle de la chlorophylle *a*. L'identification repose sur l'isolement des esters méthyliques du phéophorbide *a* d'une part (à partir des gonades entières) et de la chlorine *a*<sub>6</sub> d'autre part (à partir du chromoprotéide).

W. H. BANNISTER, M. CHOUSSEY,  
M. VUILLAUME and M. BARBIER

Department of Physiology, Royal University of Malta, Msida (Malta), and  
Institut de Chimie des Substances Naturelles, CNRS, F-91 Gif-sur-Yvette (France), 15 May 1970.

<sup>1</sup> J. H. ORTON, J. mar. Biol. Ass. U.K. 15, 851 (1928).

<sup>2</sup> W. H. BANNISTER, J. V. BANNISTER and H. MICALLEF, Comp. Biochem. Physiol. 37, 319 (1969).

<sup>3</sup> T. W. GOODWIN and M. M. TAHA, Biochem. J. 47, 244 (1950).

<sup>4</sup> J. E. FALK, *Porphyryns and Metalloporphyryns* (Elsevier Publishing Company, Amsterdam 1964).

<sup>5</sup> H. R. WETHERELL, M. J. HENDRICKSON and A. R. MCINTYRE, J. Am. chem. Soc. 81, 4517 (1959).

<sup>6</sup> A. STOLL and E. WIEDEMANN, Fortschr. chem. Forschung 2, 538 (1952).