

assumption that, on the average, for each cell which migrated out of our circumscribed area, another cell would migrate into it. The establishment and maintenance of the cell lines have been described<sup>3</sup>. For growth studies, well-dispersed cells in about 6 ml of growth medium were placed in small plastic flasks (Falcon plastics) at a concentration of 10,000 to 20,000 cells per flask and incubated in a humidified CO<sub>2</sub> incubator. A pattern of squares was scratched on the bottom of the flask, and the number of cells in each square was counted 4 to 6 h after subcultivation and daily thereafter at  $\times 100$  magnification under an inverted microscope with the illumination and diaphragm regulated to obtain maximum contrast. Each cell line's growth rate was determined in duplicate during at least 3 different passage levels.

Data from typical growth experiments are presented in Figures 1 and 2. Cells were randomly dispersed over the surface of the flask except at the edges where cells became confluent sooner. Although early experiments indicated that the location or nature of the pattern of squares did not make any difference, we always used a pattern of 3 to 6 squares, side-by-side, half way between the middle and edge of the flask and parallel to its long axis. Duplicate growth curves obtained with inocula of 10,000 to 20,000 cells were parallel; larger inocula sometimes led to areas of confluency on day 3 which caused difficulties in numeration.

Comparisons between results obtained from standard growth studies performed by trypsinization-resuspension methods and those obtained by counts of growing cells in plastic flasks showed no significant differences in doubling times. There were no variations at different passage levels during phase II<sup>5</sup>, nor when experiments were performed by different individuals. It was initially feared that debris from cells which did not attach to the plastic surface might exert a toxic effect and inhibit the multiplication of attached cells, but replacing the medium in the flasks

after 6 h incubation with fresh medium had no effect on the growth rate. This experiment also indicated that none of the increase in cell numbers after the first counts were performed was due to attachment of cells formerly in suspension to the plastic surface.

**Discussion.** This method makes possible the accurate determination of growth curves and doubling times of human fibroblasts in cell culture. The technique is simple, fast and economic. Only 20,000 to 40,000 cells and a few plastic flasks suffice for the whole experiment versus several 100,000 cells for trypsinization-resuspension methods which also entail more manipulation of cells with concomitant introduction of errors.

We have found this technique particularly useful for our comparative studies of abnormal human cell lines, but it could also be adapted to screening of various chemical and biological agents for their effects on cell multiplication, or to other routine growth studies.

**Résumé.** Description d'une technique simple permettant de mesurer le rythme de croissance in vitro des fibroblastes humains. Les suspensions cellulaires sont cultivées en flacons plastiques et les cellules présentes dans des surfaces délimitées sont dénombrées à des temps réguliers. Les applications de cette technique sont discutées.

SUSAN CURE and A. BOUÉ<sup>6</sup>

*Centre International de l'Enfance,  
Laboratoire de la S.E.S.E.P.,  
Château de Longchamp, F-75016 Paris, (France),  
1 December 1972.*

<sup>5</sup> L. HAYFLICK, *Expl Cell Res.* 37, 614 (1965).

<sup>6</sup> These investigations were supported by grants from the Delegation Générale à la Recherche Scientifique et Technique (D.G.R.S.T.).

## CONGRESSUS

### Israel

#### 1st International Congress for Bacteriology

*in Jerusalem, 2-7 September 1973.*

This will be the first international congress of the newly formed Bacteriology Section of the International Association of Microbiological Societies.

Further information about the congress may be obtained from the Congress Secretariat, P.O. Box 16271, Tel Aviv, Israel.

### Turkey

#### IAEA Symposium on Radioimmunoassay and Related Procedures in Clinical Medicine and Research

*in Istanbul, 10-14 September 1973.*

Further information by the scientific Secretaries: Dr. E. J. Garcia and Dr. E. H. Belcher, International Atomic Energy Agency, Kärlntner Ring 11-13, A-1010 Wien (Austria).

### Austria

#### First International Congress for Aerosols in Medicine

*in Vienna, 19-21 September 1973*

Aerosols in Medicine' (Advantages and Dangers). Main Topics: 19 September: Environmental Aerosols (Air Pollution), Hygienic Aspects of Aerosols.

Secretary of the Congress: Mrs. E. Weidenhaus, Wiener Medizinische Akademie, Stadiongasse 6-8, A-1010 Vienna, Austria.

### Switzerland

#### Symposium on Thirst organized by the Swiss Union of Societies of Experimental Biology

*in Lugano, 20 and 21 October 1973.*

The symposium will be held under the auspices of the International Commission 'Regulation of Food and Water Intake' of the International Union of the Physiological Sciences. Further Information by: Prof. Georges Peters, Institut de Pharmacologie de l'Université, 21, rue du Bugnon, CH-1011 Lausanne/Switzerland.