

Mutants of *Saccharomyces cerevisiae* with an Altered Regulation of Chorismate Mutase

J. Zeyer, P. Kradolfer and R. Hütter

Mikrobiologisches Institut, ETH-Zentrum, CH-8092 Zürich

Mutants of *Saccharomyces cerevisiae* with an increased sensitivity to the tryptophan analogue DL-5-methyl-tryptophan (5MT) were isolated (SCHÜRCH, MIOZZARI and HÜTTER, J. Bact. 117, 1131–1140, 1974). A number of these 5MT sensitive mutants were characterized and assigned to one of the following 3 classes: class I, strains with altered activity and/or feedback sensitivity of anthranilate synthase; class II, strains with elevated uptake of 5MT; class III, mutants with altered regulation of the tryptophan-biosynthetic enzymes which do not exhibit increases in activity in the presence of 5MT. The object of the following study was to characterize 3 5MT sensitive mutants which do not belong to one of the above mentioned groups. The 3 mutants were shown to possess the following characteristics: They belong to the same complementation group and are considered allelic, although they show slightly different phenotypes. – The 5MT inhibition no longer occurs in the presence of tryptophan, anthranilic acid or indole. – Enzymatic analysis revealed changes in chorismate mutase: a) greatly increased activity of chorismate mutase; b) diminished feedback sensitivity of this enzyme to tyrosine; c) reduction in its ability to be activated by tryptophan. – Genetic studies indicate that this mutation is dominant and is localized in the chorismate mutase gene. – Two possibilities to explain the phenotype of the mutation are being studied. We are either dealing with a promotor mutation in the chorismate mutase structural gene, or the mutation affects the equilibrium between an inactive (or less active) and an active form of the enzyme.

Some Applications of an Immobilized Esterase

J. Konecny

Ciba-Geigy AG, K-121.4.03, CH-4002 Basel

The immobilized esterase described earlier (KONECNY and SLANICKA, Biochim. biophys. Acta 403, 573, 1975; Path. Microbiol. 42, 245, 1975) was used for the quantitative alkalimetric titration of cephalosporins, for the preparation of larger quantities of deacetyl cephalosporins, and as an enzyme thermistor for their calorimetric assay. The highly active catalyst required for these applications was prepared by coupling the enzyme to zirconia-clad controlled-pore amino glass via glutaraldehyde, and had an activity of 200 units/g. The titrations and the preparative runs were carried out in 10–20 mM phosphate in a differential recirculation reactor (FORD, LAMBERT, COHEN and CHAMBERS, Enzyme Engin. 2, 267, 1972) at pH 7 and pH 8 and 25°C, the reaction time for complete conversion being about 60 and 30 min, respectively. In 80 successive runs at pH 8 the immobilized enzyme retained 70% of its initial activity, the rate of loss being highest at the beginning. In the analytical runs the titres agreed within 2% with the theory. The enzyme thermistor was the earlier version of the two devices described by MOSBACH and coworkers (MOSBACH and DANIELSSON, Biochim. biophys. Acta 364, 140, 1974; MOSBACH, DANIELSSON, BORGERUD and SCOTT, ibid. 403, 256, 1975). In current work the sluggish response of the system arising from its high heat capacity was reduced to a few minutes by substituting the isothermal Tronac calorimeter for the thermistor.

Product Inhibition During the Rifamycin S Fermentation

M. T. Küenzi, J. Gruner, A. Fiechter and J. Nüesch

Ciba-Geigy AG, Werk Klybeck, CH-4002 Basel; and Mikrobiologisches Institut, ETHZ, Weinbergstrasse 38, CH-8006 Zürich

The production of rifamycin S by a mutant strain of *Nocardia mediterranei* in a complex medium was investigated. The formation of the antibiotic was associated with the growth of the organism. Growth and production occurred in two phases. After a first phase of rapid growth and production metabolic activity slowed down and continued at a reduced rate until the energy source in the medium was exhausted. Substrate limitation and product inhibition were examined as possible reasons for this behaviour. Improved mixing conditions reduced pellet formation but did not change the course of the fermentation. Thus, substrate limitation could be excluded as an explanation for the drop in growth and production rates. It was found that rifamycin S inhibits its own synthesis as well as the growth of the producing organism. By using an in vitro system the specific rifamycin S synthesizing activity (RS-activity) of mycelium from different stages of a fermentation could be measured. This RS-activity paralleled the production rates calculated from a fermentation. The values increased in the first part of a fermentation and then dropped rapidly. Addition of rifamycin S to a growing culture inhibited this initial increase in RS-activity and prevented in this way the organism from producing higher quantities of the antibiotic. The implications of these results for the development of an improved production process are discussed.

Some Properties of the Endotoxin Unit of *Bacillus thuringiensis*

B. Trümpler and P. Lüthy

Mikrobiologisches Institut der ETH, Universitätsstrasse 2, CH-8092 Zürich

In a previous study it was shown that purified crystal suspensions of *Bacillus thuringiensis* var. *thuringiensis*, which were dissolved by non-proteolytic solvents such as NaOH, mercaptoethanol/urea or mercaptoethanol/guanidine hydrochloride, yielded peptides with a molecular weight $\geq 800,000$. This material retained its full toxic activity. Further degradation by purified gut juice proteases resulted in a toxic component with a molecular weight of 100,000 accompanied by a non-toxic low molecular weight fraction $\leq 10,000$. The molecule of MW 100,000 is referred to as the toxic unit. Using the same technique, endotoxin crystals belonging to the varieties *kurstaki*, *entomocidus*, *aizawai*, *morrisoni*, *tolworthi*, *darmstadtensis* and *toumanoffi* were analyzed. It was demonstrated by gel filtration and by gel electrophoresis that the predominant protein component of each of the above mentioned varieties appeared in the molecular weight range of 100,000. The low molecular weight fragments were also present. According to our results, it may be presumed that the toxic units of the different varieties were very similar. In further experiments the high molecular weight component (MW $\geq 800,000$) was heated to 100°C for 1 h. Enzymatic degradation of the heat treated protein yielded only a low molecular weight component whereas the toxic unit was missing. Thus, the intact structure of this protein is essential for the formation of the toxic unit by gut juice proteases. Finally, the SH-groups of the toxic protein unit were carboxymethylated

by iodoacetic acid. Carboxymethylation results in a complete loss of toxic activity. This shows that the toxic unit must be present in its secondary or even tertiary structure in order to display its activity.

In vitro Activity of the Endotoxin of *Bacillus thuringiensis*

P. Geiser, H.-R. Ebersold and P. Lüthy
Mikrobiologisches Institut
ETH-Zentrum, CH-8092 Zürich

The mode of action of the endotoxin formed by *Bacillus thuringiensis* is still unknown, mainly because no activity of the crystal protein could be demonstrated in any system, except within susceptible insects. Using primary cell cultures derived from gut epithelium and from hemocytes of *Pieris brassicae*, we were able to induce changes by the addition of endotoxin. Damaging effects were not limited to the culture of gut epithelial cells but occurred also within hemocytes.

The endotoxin was obtained by digesting purified crystal suspensions with gut juice proteases of *P. brassicae*. The resulting toxic unit with a molecular weight of 100,000 was isolated by gel filtration. The concentration of endotoxin added to the cell cultures amounted to 30 µg/ml. The first morphological changes of gut epithelial cells could be observed within 15 to 30 min following the addition of endotoxin. The cells doubled in size by forming bubble-like protrusions. High pH-values of the medium and the presence of proteases seemed to stimulate this reaction. The addition of endotoxin to hemocyte cultures caused identical morphological changes within 15 to 30 min. Vital staining 2 h after the addition of the endotoxin showed that 80 to 100% of the cells had lost their viability. Again, the presence of proteolytic enzymes enhanced the destruction of the hemocytes. Controls were conducted with heat inactivated endotoxin. In this case, no morphological alterations were observed and after 2 h, 95 to 100% of the insect cells had retained their viability.

Supported by SNSF, grant 3.4810.75.

PRAEMIA

Preis der Schweizerischen Gesellschaft für Mikrobiologie, 1976

Anlässlich der Jahresversammlung in Genf vom 17. bis 19. Juni 1976 hat die Schweizerische Gesellschaft für Mikrobiologie Herrn PD Dr. H. Bienz den Preis 1976 der Gesellschaft verliehen. Der Laureat, geb. 1940, arbeitet am Institut für Mikrobiologie und Hygiene der Universität Basel und hat sich durch seine Leistungen auf dem Gebiet der Virusforschung und der Mikrobiologieausbildung von Medizinern ausgezeichnet. Herr PD Dr. Bienz ist der erste Preisträger.

17. Juni 1976

Prof. Dr. R. Hütter, Präsident

Prize 'Biochemical Analysis' 1978

The prize of DM 10,000.— is donated from Boehringer in Mannheim, and is awarded every two years at the conference 'Biochemische Analytik' in Munich for outstanding work in the field of biochemical instrumentation and analysis. The donation will take place between 18 and 21 April during the 1978 conference. One or several papers concerning one theme each, either published or accepted for publication between 1 October 1975 and 30 September 1977, may be sent to: Prof. Dr I. Trautschold, Secretary of 'Biochemical Analysis', Medizinische Hochschule Hannover, Karl-Wiechert-Allee 9, D-3000 Hannover 61, Federal Republic of Germany.

CONGRESSUS

Italy The 4th International Symposium on Mass Spectrometry in Biochemistry

at Riva del Garda, Lake of Garda, 20–22 June 1977

The Symposium will be devoted to topics such as: Gas chromatography-mass spectrometry, mass fragmentography, stable isotope measurements, field ionization, field desorption, chemical ionization, high resolution studies and data of acquisition and processing. The areas of application will include biochemistry, medicine, toxicology, drug research, forensic science, clinical chemistry and pollution.

Those wishing to present a communication (approx. 20 min) are requested to submit the title and an abstract of not more than 200 words written in English before 1 March 1977. Further details by: Dr Alberto Frigerio, Istituto di Ricerche Farmacologiche 'Mario Negri', via Eritrea 62, I-20157 Milano, Italy.

Switzerland The 6th Annual Conference of the International Society for Experimental Hematology

in Basel, 28–31 August 1977

Scientific program: Regulation and differentiation of hemopoetic stem cells; hemopoetic inductive micro environment; exogenous effects of hemopoiesis; experimental models of hematologic disorders. – Immunity: lymphocytes, monocytes, macrophages; histocompatibility testing; cell interactions; immune suppression and stimulation; immune deficiency states; immunotherapy; cellular engineering; clinical and experimental neoplasias of hemopoetic tissues. – Bone marrow transplantation: experimental models; clinical marrow grafts for immune deficiency syndromes; aplastic anaemia, malignant disease and other indications; grafts across MHC and ABO barriers. Further information by: Congress Secretariat, ISEH, P.O. Box 129, CH-4004 Basel, Switzerland.