

bivalents (figure, A). Bivalents with 1 or 2 chiasmata are more in number. Subsequently most chiasmata get terminalized during late diplotene and diakinesis (figure, B, C). Finally the highly condensed bivalents settle at the equator (figure, D). Absence of the peculiar premeta-phase stretch of the bivalents, seems to be unique for this roach, while it is quite common in other genera like *Periplaneta*, *Blatta*, *Nauphoeta*, etc. The X-chromosome shows a somewhat precocious condensation. The condensation is completed by the beginning of diakinesis. Segregation of the X is reductional at Meiosis I.

Of the 11 bivalents 7–8 show one chiasma each at late diplotene and diakinesis, while 3–4 show 2 chiasmata each as a rule (figure, B, C). In the latter group, one bivalent occasionally shows 3 chiasmata. 20 spermatocytes were examined, from the preparations of each of the males collected, to work out the average chiasma frequency.

The average chiasma frequency per bivalent is 1.3, and per cell it is 14.6. The recombination index (haploid

number plus mean of chiasmata, according to Darlington) is therefore 26.6. From the foregoing account it is quite clear that in *Blattella suppellectilium* there is intimate pairing of the homologues with clear evidence of crossing over, not necessarily in the terminal regions of the bivalents. Further, the recombination index is also fairly high. One might expect the genus *Blattella* to have produced a good number of species. On the contrary, it has produced 3 species only. The reason for this lack of diversity may be the almost contiguous distribution of populations of this genus throughout the tropical and temperate regions, though probably a native of North Africa<sup>12</sup> and reached India and other tropical countries via Europe<sup>13</sup>, with a concomitant lack of isolating factors. This study on *Blattella suppellectilium* supplies yet another example of chiasmate type of meiosis in cockroaches.

12 D. R. Ragge, Frederick Warne & Co. Ltd., Lond. Publ., p. 64 (1965).

13 H. M. Lefroy, Thacker Spink & Co. Publ. Lond., p. 60 (1909).

### Concrement formation encountered in the rat pineal gland

F. Erdinc

*Department of Histology and Embryology, Istanbul Medical Faculty, University of Istanbul, Istanbul (Turkey), 30 September 1976*

**Summary.** Morphological evidence is given for the concretum formation encountered in the rat pineal gland, the occurrence of which has not been reported so far (for the rat).


The concretum formations (brain sand, *acervuli cerebri*) in the pineal gland are areas of calcification appearing in the form of concentrically arranged rings or as amorphous structures which are commonly found in the human pineal glands and in the glands of some animals as horse, donkey and cattle<sup>1</sup>. However, the occurrence of these formations in the rat pineal gland has not been observed<sup>2</sup> and reported so far.

In the course of histological examinations of 2.5–3 months old normal adult albino rat pineal glands *acervulus* formation was observed in a female rat pineal gland (figure) in the vicinity of which the homogenous material is also detectable. Although the origin and the pattern of *acervulus* formation have not been clearly revealed so far, it is generally considered that the initial stage is the formation of colloid in the pinealocyte cytoplasm and then the participation of parenchymal and the connective tissue

elements<sup>1,3,4</sup>. Its composition is found to be close to myelin sheaths of the nerve fibres<sup>4–6</sup> and, on the other hand, mineralogically similar to enamel<sup>7</sup>.

Reports in relation with the formation of pineal *acervuli* are contradictory; it has been attributed to degeneration or ageing processes<sup>3</sup>, while most of the studies do not consider it as a pathological sign or regression<sup>1,5,8,9</sup>. Reports on the occurrence of *acervuli* in the pineal glands of the newborn<sup>5</sup> and of children<sup>10</sup>, and whereas its absence in the aged<sup>8</sup> support the second view. Also we have encountered *acervuli*, although in small quantities, in the lamb pineal glands, which also support the view that *acervulus* formation shows individual variations depending on the individual conditions<sup>9,9</sup>. Our observation of pineal *acervulus* in the rat strengthens this opinion furthermore, indicating that *acervulus* formation can commonly occur in the rat as well, and cannot be considered as specific to some of the mammalian species, or dependent on some external factors, since we encountered it in one animal although all of the animals were kept under the same conditions.

Since functional significance of pineal *acervuli* still remains obscure, it is not possible to explain the *acervulus* formation in the rat pineal gland or to answer the question: why not in all rats, and what is this special factor? We hope that further studies on the pineal gland will give us the answers.



Concrement in the pineal gland of a rat. Fixation: Bouin's fluid. Stain: Hematoxylin-eosin.  $\times 1200$ .

- 1 K. Scharenberg and L. Liss, *Progr. Brain Res.* 10, 216 (1965).
- 2 B. S. Şayli, Thesis, University of Ankara, Medical Faculty, Ankara (1962).
- 3 L. Thiéblot and H. Le Bars, in: *La glande pinéale ou épiphyse*. Librairie Maloine S. A., Paris 1955.
- 4 A. M. Khelinskii, *Probl. Endokr.*, Moskva 15, 50 (1969).
- 5 G. Heidel, *Anat. Anz.* 116, 139 (1965).
- 6 A. Bayer and G. Bayerova, *Endokrinologie* 52, 1 (1967).
- 7 C. P. Mabie and B. M. Wallace, *Calc. Tiss. Res.* 16, 59 (1974).
- 8 M. Milcu and Gh. Vrejoiu, *Rum. med. Rev.* 3, 13 (1959).
- 9 W. B. Quay, *Progr. Brain Res.* 10, 69 (1965).
- 10 E. Wildi and E. Frauchiger, *Progr. Brain Res.* 10, 222 (1965).