

- 4 Ardry, R.M., *Anals pharm. fr.* 14 (1956) 97.
- 5 Chauvin, R., and Herbert, S., *C.r. Soc. Biol., Paris* 150 (1956) 332.
- 6 Kalinin, A.P., and Radbil, O.S., *Endocr. Hormonother.* 8 (1962) 119.
- 7 Vittek, J., and Janči, J., *Royal Jelly*, p.272. SVPL, Bratislava 1968.
- 8 Dixit, P.K., and Patel, N.G., *Nature* 202 (1964) 189.
- 9 Kramer, K.J., Tager, H.S., Childs, C.N., and Speirs, R.D., *J. Insect Physiol.* 23 (1977) 203.
- 10 Takahashi, K., Kiuchi, K., Endo, M., Furuno, Y., Shinozuka, T., and Hasegawa, S., *C.r. Soc. Biol.* 157 (1963) 428.
- 11 Heyl, H.L., *Science* 89 (1939) 540.
- 12 Donault, M.P., *C.r. Acad. Sci.* 257 (1963) 258.
- 13 Jakab, M., Orban, G., Richter, P., and Fekete, G., *Acta pharm. hung.* 30 (1960) 170.
- 14 Šterba, R., Kvičalová, E., Šaško, E., and Danielisová, E., *Parfum. Kozmet.* 43 (1964) 307.
- 15 Brown, W.H., Felauer, E.E., and Freure, R.J., *Can. J. Chem.* 39 (1961) 1086.
- 16 Pain, J., Barbier, M., Bohdanovsky, D., and Lederer, E., *Comp. Biochem. Physiol.* 6 (1962) 233.
- 17 Carter, H.E., and Hirschberg, C.B., *Biochemistry* 7 (1968) 2296.
- 18 Gilbert, L.I., Bollenbacher, W.E., and Granger, N.A., *A. Rev. Physiol.* 42 (1980) 493.
- 19 Clayton, R.B., *J. Lipid Res.* 5 (1964) 3.
- 20 Schildknecht, H., Birringer, H., and Maschwitz, U., *Angew. Chem.* 6 (1967) 558.
- 21 Schildknecht, H., *Angew. Chem.* 4 (1966) 421.
- 22 Barker, S.A., Forster, A.B., and Lamb, D.C., *Nature, Lond.* 184 (1959) 634.
- 23 Callow, R.K., Johnston, J.F., and Simpson, J., *Experientia* 15 (1959) 421.
- 24 Riad-Fahmy, D., Read, G.F., and Walker, R.F., *Postgr. Med. J.* 56, Suppl. 1 (1980) 75.

0014-4754/84/010104-03\$1.50 + 0.20/0

© Birkhäuser Verlag Basel, 1984

Calcium transients in a crustacean motoneuron soma: Detection with arsenazo III

G. Czernasty, M. Thieffry and I. Parker

Laboratoire de Neurobiologie, U.E.R. Sciences Exactes et Naturelles, 33, rue St Leu, F-80039 Amiens (France), Laboratoire de Neurobiologie Cellulaire, C.N.R.S., F-91190 Gif-sur-Yvette (France), and Department of Biophysics, University College London, Gower Street, London WC1E6BT (England), 3 June 1982

Summary. Long lasting action potentials can be triggered in crayfish giant motor neurons by a short depolarizing pulse under different conditions. The concomitant increase in absorbance of the Ca indicator arsenazo III preloaded into the soma, confirms previous observations suggesting that these potential changes are related to a Ca inward current through the soma membrane.

The giant motoneurone (MoG) is a large cell, found bilaterally in the crayfish abdominal ganglia. Early work¹ suggested that the cell soma does not show any regenerative action potential in response to a depolarizing pulse, but it is now known that regenerative responses can be recorded under certain conditions. For instance, when the K⁺ conductance of the membrane is reduced (either by TEA⁸ or following inactivation by a conditioning depolarisation), a slow action potential (SAP) may be triggered by a short depolarization. This response is most probably due to an influx of Ca²⁺ ions, since it depends upon the presence of Ca²⁺, but not Na⁺ in the bathing solution, and is blocked by Co²⁺ and Mn²⁺, but not by TTX². In the preparations pretreated with TEA, the SAPs can also be recorded at more hyperpolarized levels than the normal resting potential. TEA is effective either injected into the MoG soma or added to the saline³. In this paper most of the results were obtained in this way when TEA was added to the solution. We report below that the internal free Ca concentration increases during the SAP, as demonstrated by the change in absorbance of the Ca²⁺ indicator dye arsenazo III preloaded into the MoG soma. This observation strengthens previous observations suggesting that SAP is associated with an influx of Ca²⁺ ions.

Methods. Experiments were carried out on the giant motor neurons of the crayfish *Procambarus clarkii*. The ventral chord was dissected and perfused with Van Harreveld's solution containing (in mM): NaCl 195; KCl 5.4; CaCl₂ 13; MgCl₂ 2.6; Tris maleate 10, adjusted to pH 7.2 by NaOH. The 2nd, 3rd and 4th ganglia were desheathed and used for the experiments. The membrane potential was monitored by conventional electrophysiological techniques using a 3 M KCl electrode. A 2nd electrode filled with an aqueous

solution of 1 mM arsenazo III (Sigma grade 1) was also inserted in the MoG soma. The dye was injected by negative current pulses (500 msec, 1 Hz) which hyperpolarized the cell by 30–40 mV. The injection was stopped when the soma appeared just detectably stained as judged by eye. The Arsenazo electrode was then used for passing both the conditioning depolarizing current pulse and the brief (2–5 msec) pulse triggering the SAP. The intracellular calcium transients were monitored by recording the differential absorbance of the dye at 650 and 700 nm as previously described^{4,5}.

Results. Typical responses obtained in TEA-treated preparations are shown in figure 1. The SAP was almost unchanged in Na-free solutions (fig. 1A). On the contrary, when Ca²⁺ was omitted from the bathing solution, the SAP was abolished (fig. 1B). These observations, together with the fact that Co²⁺ and Mn²⁺, but not TTX, block the SAP make it very likely that the response is due to an influx of Ca²⁺ ions^{2,3}. It is therefore interesting to look for possible changes in intracellular ([Ca²⁺]_i) during the SAP. Simultaneous records of membrane potential and arsenazo absorbance from a MoG cell soma are shown in figure 2. Following a conditioning depolarization from the resting potential (V_r = 66.9 mV ± 5.3; N = 59), a long lasting action potential is elicited by a brief depolarizing pulse in preparations not treated with TEA (fig. 2A). During the SAP a rise in arsenazo absorbance is seen, which returns slowly to the baseline after the cell has repolarized. This indicates that the intracellular free Ca²⁺ level increases steadily throughout the duration of the SAP. Control records obtained at a wavelength of 570 nm (the isosbestic point of the dye)⁵ demonstrated that the arsenazo Ca²⁺ signal is not appreciably contaminated by any other optical changes




Figure 1. Effects of Na^+ and Ca^{2+} removal upon the SAPs triggered at the resting potential in MoG pretreated for at least 30 min in solutions containing 75 mM TEA. *A* Before (left) and after (right) substitution of Na^+ by an equimolar concentration of choline. *B* Before (left) and after (right) total substitution of Ca^{2+} by an equimolar concentration of Mg^{2+} . Top trace: intracellular potential; bottom trace: triggering current.




Figure 2. Arsenazo light responses (upper trace) and slow action potentials (lower trace) recorded in different conditions. The optical record is a differential recording of light transmission at the wavelength pair 650–700 nm. An upward deflection corresponds to an increase in dye absorbance at 650 nm, and signals an increase in free (Ca^{2+}) . The starting point of the arsenazo trace was reset at the beginning of each sweep. *A* In normal saline, the soma potential was shifted from the resting potential to -45 mV about 5 sec before applying the triggering pulse. *B*, *C*, *D* Responses of another cell in the presence of 75 mM TEA. Single depolarizing pulses applied at holding potentials of -65 mV (*B*) and -50 mV (*C*) triggered action potentials of different duration. *D* Two depolarizing pulses were applied 2 sec apart at a holding potential of -58 mV. In response to the 2nd pulse, the duration of both signals was decreased. Calibrations: Membrane potential = 20 mV, arsenazo signal expressed as $\Delta I = 1.7 \times 10^{-3}$ (A) and 3.6×10^{-3} (B, C, D).

during the SAP. In a saline containing 75 mM TEA, a similar transient increase in absorbance occurred during the SAP (fig. 2B).

The relationship between the duration of the SAP and that of the rising phase of the arsenazo signal could be directly verified by two different methods using the same cell: 1. The SAP duration strongly depends on the holding membrane potential. When the latter was shifted from -65 mV (fig. 2B) to -50 mV (fig. 2C), both the duration of the SAP and the rising phase of the arsenazo signal increased by about 50%. However, the rate of rise of the arsenazo signal then decreased, the amplitude being similar to that observed at -65 mV. 2. The SAP duration (and to a much lesser extent its amplitude) is very sensitive to repetitive activation². Two short depolarizing pulses applied 2 sec apart to the same cell triggered SAPs of decreasing dura-

tion (fig. 2D) though the 2nd SAP had been triggered at a slightly more depolarized level. Again the duration of rise of arsenazo signal fitted that of the SAP.

Unlike the other motoneurons, MoG does not arborize in the neuropile. Its neurite is very thin at the beginning and leaves the soma obliquely in a dorso-caudal direction^{6,7}. The area analyzed by the optical system could thus be limited by an image of a circular diaphragm focussed onto the larger part of the soma. This minimized a possible contribution of other parts of the cell to the recorded arsenazo signal.

This gives additional evidence that the SAP results from an inward Ca current through the soma membrane. Quantitative measurements of $[\text{Ca}^{2+}]_i$ are now under investigation. The records in figure 2 show a steady increase in $[\text{Ca}^{2+}]_i$, which suggests that most of the Ca^{2+} signal is due to a

steady inward flux. If most of the Ca^{2+} signal were due to a sudden surge at the beginning of the SAP, the arsenazo signal should not show this steady rise.

Furthermore, the time constant of the membrane ($\cong 4$ msec at rest) is much too small to account for the SAP duration. Since the SAP is still observed in Na-free solutions and requires external Ca^{2+} (fig. 1), the increase in the arsenazo signal occurring during the long lasting depolarization is most probably due to the Ca^{2+} entering the cell rather than

to Ca^{2+} released from cytoplasmic pools secondary to depolarization. Thus a more precise characterization of the Ca channels and of the mechanisms underlying their inactivation may be expected from the use of the arsenazo dye. The kinetics of Ca^{2+} entry and Ca^{2+} removal obtained from the arsenazo signal will be especially useful for studying whether internal Ca^{2+} concentration affects Ca conductance on this preparation as it does in other preparations^{9,11}.

- 1 Takeda, K., and Kennedy, D., J. Cell comp. Physiol. 48 (1964) 435.
- 2 Czternasty, G., and Bruner, J., Neurosci. Lett. suppl. 7 (1981) 379.
- 3 Czternasty, G., Kado, R.T., and Bruner, J., Neurosci. Lett. suppl. 10 (1982) 127.
- 4 Parker, I., in: Detection and measurements of free Ca^{2+} in cells, p. 269. Eds C.C. Ashley and A.K. Campbell. Elsevier, Amsterdam 1979.
- 5 Miledi, R., Parker, I., and Zhu, P.H., J. Physiol., Lond. 333 (1982) 655.
- 6 Selverston, A.I., and Kennedy, D., Endeavour 28 (1969) 107.
- 7 Mittenthal, J.E., and Wine, J.J., J. comp. Neurol. 177 (1978) 311.
- 8 Narahashi, T., Physiol. Rev. 54 (1974) 813.
- 9 Tillotson, D., Proc. natl Acad. Sci. USA 76 (1979) 1497.
- 10 Standen, N.B., Nature 293 (1981) 158.
- 11 Ashcroft, F.M., and Stanfield, P.R., Science 213 (1981) 224.

0014-4754/84/010106-03\$1.50 + 0.20/0
© Birkhäuser Verlag Basel, 1984

Nervous and behavioral responses to light in colonies of the free-living bryozoan, *Selenaria maculata* (Busk)¹

M.S. Berry and P.J. Hayward

Department of Zoology, University College of Swansea, Singleton Park, Swansea SA2 8PP (Wales, Great Britain),
22 Novembre 1982

Summary. Bursts of electrical impulses which spread throughout the entire colony were recorded from individual zooids of *Selenaria*. Strong illumination caused a rapid increase in the frequency of bursts and also caused the colonies to start moving, usually towards the light. The recorded electrical activity may act to trigger locomotor behavior.

Bryozoa are generally small, colonial, sessile animals commonly found encrusting as flat sheets on rocks or seaweed². Adjacent zooids in the colony are interconnected, and in some species at least there appears to be a colonial nervous system³⁻⁵. Thorpe et al.^{6,7} recorded nerve impulses traveling across the colony of *Membranipora membranacea* (L.) in response to electrical or mechanical stimulation of the frontal membrane of a single zooid. The impulses were recorded from the surface of the colony and were found to influence retraction of the lophophores (ciliated food-collecting tentacles).

In contrast to these simple, sessile colonies, the family Selenariidae (Cheilostomata, Anasca) includes free-living species which show some remarkable behavior patterns^{8,9}. The conical colony of *S. maculata* (Busk.) is illustrated in figure 1. The first reported observations of living *Selenaria* were made by Cook and Chimonides^{8,9} who found that the peripheral avicularian setae (which in other Bryozoa act to clean the colony surface of detritus) can raise and support the colony, and propel it over the substratum at rates of up to 1 m/h. Colonies move laterally in a random manner except when strongly illuminated from one side when they move towards the light^{8,9}. We have used electrophysiological techniques to investigate the neural basis of locomotion in *Selenaria*, and the response to light.

Methods. Colonies of *S. maculata* were collected from near Townsville (Queensland, Australia) and flown to Swansea where they were maintained in synthetic seawater (Instant Ocean) at 26°C. Glass-tipped suction electrodes of about 100 µm diameter were used to record from the frontal membrane of individual zooids, and the electrical activity was amplified, displayed on a storage oscilloscope (Tektronix 5111) and photographed with a Polaroid camera.

Strong illumination was provided by a 3 kW tungsten lamp, the light first passing through water to absorb the heat. The level of illumination was varied between 1.3 and 42 W/m² for wavelengths of 400–700 nm (behavioral responses⁹ are produced by wavelengths of 400–560 nm).

Results and discussion. Under normal laboratory lighting (about 1.3 W/m²) colonies usually showed no locomotion and little movement of the setae. Recordings from individual zooids showed bursts of impulses (fig. 2) occurring at irregular intervals of at least 1 min and often more than 10 min. In confirmation of the studies by Cook and Chimonides^{8,9} increased irradiance (up to 42 W/m²) initiated or increased setal movements within 2 min and colonies were observed to move (usually towards the light) within 10 min. Increased light intensity produced a rapid increase in frequency of bursts which persisted for the duration of the illumination and then returned to previous levels within




Figure 1. Lateral view of a living colony of *Selenaria maculata* (Busk) supported above the substratum (a glass plate) by the peripheral and subperipheral avicularian setae. Diameter of colony, 15 mm. (Photograph courtesy of Mr P.J. Chimonides).