

## CORRIGENDUM

In Table 2 of this article, the second and third columns were switched for the lysine residues. The correct table is given here. The authors thank Anja Böckmann (PBIL-IBPC, Lyon Gerland, France) for bringing this error to their attention.

**Table 2.** Activation energies ( $E_a$ ) and the pre-exponential factors ( $A$ ) [see Eq. (6)] of the  $H^{N\epsilon}$  protons of the four arginine and the  $HN\alpha$  protons of six lysine residues in human ubiquitin at pH 7.45.

Residue	$\ln A$	$E_a$ [kJ mol <sup>-1</sup> ]
R42	32.5 ± 3.5	64.3 ± 8.6
R54	33.0 ± 0.8	68.1 ± 1.9
R72	37.2 ± 0.6	75.6 ± 1.5
R74	36.5 ± 0.2	73.5 ± 0.4
K6	27.1 ± 0.9	41.8 ± 2.1
K11	36.2 ± 1.0	67.1 ± 2.4
K29	40.2 ± 17.6	79.0 ± 44.3
K33	34.5 ± 1.1	63.3 ± 2.8
K48	35.4 ± 0.6	64.8 ± 1.4
K63	36.9 ± 0.7	69.2 ± 1.8

*T. Segawa, F. Kateb, L. Duma,  
G. Bodenhausen, P. Pelupessy\**

**Exchange Rate Constants of Invisible  
Protons in Proteins Determined by  
NMR Spectroscopy**

*ChemBioChem* **2008**, *9*, 537–542

DOI 10.1002/cbic.200700600