




Solar Cells


A. Ghicov, S. P. Albu, R. Hahn, D. Kim, T. Stergiopoulos, J. Kunze, C.-A. Schiller, P. Falaras, P. Schmuki*

TiO₂ Nanotubes in Dye-Sensitized Solar Cells: Critical Factors for the Conversion Efficiency

Particle vs tube: The present paper systematically investigates a range of fundamental geometrical and structural features of TiO₂ nanotube layers and their effect on the dye-sensitized solar cell conversion efficiency, to deduce the most promising strategies for improvement. It is found that the performance of the cells strongly depends on the morphology and crystalline structure of the nanotubes.



Chem. Asian J.
DOI: 10.1002/asia.200800441




Protein Aggregation


A. Taniguchi, Y. Sohma, Y. Hirayama, H. Mukai, T. Kimura, Y. Hayashi, K. Matsuzaki, Y. Kiso*

"Click Peptide": pH-Triggered in Situ Production and Aggregation of Monomer A β 1–42

Into neutral: We demonstrate the unique features of a pH click peptide based on an O-acyl isopeptide method. Under acidic conditions, the click peptide remains in a monomeric form. Upon increase of the pH to 7.4, the click peptide is quickly able to convert into A β 1–42 through an O-to-N intramolecular acyl migration. Further study using this pH click peptide would elucidate the pathological role of A β 1–42 in Alzheimer's disease.



ChemBioChem
DOI: 10.1002/cbic.200800765




Carbon Nanotubes

P. Gründler,* O. Frank, L. Kavan, L. Dunsch

Carbon Nanotube Electrodes for Hot-Wire Electrochemistry

Hot-wired electrodes: Thin metallic wires ($d=25\ \mu\text{m}$) are covered with a $3\ \mu\text{m}$ layer of single-walled carbon nanotubes (SWCNTs; see image) by electrophoresis from a suspension containing excess ionic surfactant. A pure SWCNT surface is achieved by heating the electrode in air. Strong differences between covered and bare metallic electrodes occur with in situ heating during electrochemical experiments.



ChemPhysChem
DOI: 10.1002/cphc.200800659




Halogenated Fullerenes

E. Kemnitz,* S. I. Troyanov*


Connectivity Patterns of Two C₉₀ Isomers Provided by the Structure Elucidation of C₉₀Cl₃₂

Two for the price of one: The first halogenated derivative of C₉₀, C₉₀Cl₃₂ (see structure; gray C, green Cl), is obtained by chlorination of a higher fullerene mixture with SbCl₅. Its molecular structure, elucidated by single crystal X-ray diffraction, reveals the presence of two isomeric C₉₀ cages that correspond to C_{2v} isomer 46 and C_s isomer 34. The addition of 32 chlorine atoms is the maximum degree of chlorination achieved for fullerenes.



Angew. Chem. Int. Ed.
DOI: 10.1002/anie.200806332

Pd-Catalyzed Carbonylation



Eur. J. Inorg. Chem.
DOI: 10.1002/ejic.200801047


S. Yamashita, N. Kurono, H. Senboku, M. Tokuda, K. Orito*

Synthesis of Phenanthro[9,10-*b*]indolizidin-9-ones, Phenanthro[9,10-*b*]quinolizidin-9-one, and Related Benzolactams by Pd(OAc)₂-Catalyzed Direct Aromatic Carbonylation

Phenanthro[9,10-*b*]indolizidin-9-ones, phenanthro[9,10-*b*]quinolizidin-9-one, and related benzolactams were obtained by phosphane-free Pd(OAc)₂-catalyzed direct aromatic carbonylation from the corresponding amines. This constitutes a formal synthesis of the representative phenanthroindolizidine and -quinolizidine alkaloids (±)-tylophorine, (±)-antofine, and (±)-cryptopleurine.



Subphthalocyanine Chemistry




Eur. J. Org. Chem.
DOI: 10.1002/ejoc.200801250


D. González-Rodríguez, T. Torres*

Peripheral Functionalization of Subphthalocyanines

Some useful procedures for the incorporation of diverse functional groups in the periphery of the subphthalocyanine macrocycle are described.



Asymmetric Synthesis



Chem. Eur. J.
DOI: 10.1002/chem.200802383


P. Selig, E. Herdtweck, T. Bach*

Total Synthesis of Meloscine by a [2 + 2]-Photocycloaddition/Ring-Expansion Route

The stereogenic centers at C3 and C12 of meloscine (**3**) can be established in the photochemical key step **1** → **2**. 1,2-Retro-benzilic acid rearrangement to a five-membered ring, reductive amination, Claisen rearrangement, and ring-closing metathesis are further key steps in the transformation of cyclobutane **2** into the target molecule **3** (14 steps, 9% overall yield). Enantioselective access to (+)-meloscine was possible when the [2 + 2]-photocycloaddition was conducted in the presence of a chiral template.



Fuel Cell Membranes



ChemSusChem
DOI: 10.1002/cssc.200800237

J. L. Horan, A. Genupur, H. Ren, B. J. Sikora, M.-C. Kuo, F. Meng, S. F. Dec, G. M. Haugen, M. A. Yandrasits, S. J. Hamrock, M. H. Frey, A. M. Herring*

Copolymerization of Divinylsilyl-11-silicotungstic Acid with Butyl Acrylate and Hexanediol Diacrylate: Synthesis of a Highly Proton-Conductive Membrane for Fuel-Cell Applications

Highly conductive to high conductivity: Polyoxometalates were incorporated in the backbone of a hydrocarbon polymer to produce proton-conducting films. These first-generation materials contain large, dispersed clusters of polyoxometalates. Although the morphology of these films is not yet optimal, they already demonstrate practical proton conductivities and proton diffusion within the clusters appears to be very high.

