

[Chem. Pharm. Bull.]
31(2) 730—732 (1983)

Studies on Tetrahydroisoquinolines. XX.¹⁾ A Novel Synthesis of 8-Aryl-1,2,3,4-tetrahydroisoquinolines²⁾

HIROSHI HARA, OSAMU HOSHINO, and BUNSUKE UMEZAWA*

Faculty of Pharmaceutical Sciences, Science University of Tokyo,
12, Ichigaya Funagawara-machi, Shinjuku-ku, Tokyo 162, Japan

(Received August 11, 1982)

Treatment of a mixture of the *p*-quinol acetate (**4**) and aryl ethers with trifluoroacetic acid gave 8-aryl-1,2,3,4-tetrahydroisoquinolines (**1a—f**) in good yields. Similar reaction of **4** and corypalline (**3**) afforded a corypalline dimer (**5**).

Keywords—oxidation; lead tetraacetate; *p*-quinol acetate; trifluoroacetic acid; deuterium-exchange; NMR; arylation; nucleophilic substitution; tetrahydroisoquinoline

8-Aryl-1,2,3,4-tetrahydroisoquinolines (**1**) are regarded as the lower homologs of aporphines (**2**); that is, the former compounds lack the C-7 methylene group in the C-ring of the latter compounds. In spite of the potential biological activity³⁾ of the former compounds, an efficient synthesis has not yet been developed. A significant drawback in the known methods⁴⁾ is the difficulty of preparing **1**, which possesses the same oxygenation pattern as the natural aporphines, in addition to the large number of reactoin steps. Here we wish to report a convenient synthetic method starting from 7-hydroxy-6-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline (corypalline) (**3**).

As an extension of the intermolecular nucleophilic reaction⁵⁾ at the C-8 position of 1,2,3,4-tetrahydroisoquinolines *via* *p*-quinol acetate (**4**), phenylation with phenyl ethers or a phenol was tried. The key compound **4**, which was prepared from corypalline (**3**) as usual,⁶⁾ smoothly reacted with veratrole in the presence of trifluoroacetic acid in CH₂Cl₂ to give 8-(3,4-dimethoxyphenyl)corypalline (**1a**) in 70% yield. Structure assignment was based on nuclear magnetic resonance (NMR) spectroscopic evidence. Namely, three methoxyl signals at δ 3.86, 3.89, and 3.93 showed that veratrole was substituted at the C-5 or C-8 position. Moreover, the up-field shift of the C-1 methylene protons (δ 3.18) of **1a** as compared to those (δ 3.44) of




Chart 1

TABLE I. Melting Points and Yield of the Products

Compound	mp (°C)	Recryst. solvent	Yield (%)
1a	183—185	iso-PrOH	70
1b	129—130	Ether-acetone	42
1c	203—204	iso-PrOH	52
1d	163—166	<i>n</i> -Hexane	43
1e	152—155 ^{a)}	MeOH	29
1f	207—210	iso-PrOH-ether	36
5	227—229	iso-PrOH	22

a) Methiodide of the acetate.

TABLE II. IR and NMR Spectral Data for the Products (**1a—f**)

Compound	IR (cm ⁻¹) OH	NMR (δ)			
		NMe	1-H ₂	OMe	Others
1a	3535	2.34	3.18	3.86, 3.89, 3.93	
1b	3540	2.41	3.26	3.66, 3.90, 3.91, 3.93	6.65 (5-H), 6.74, ^{a)} 6.84 ^{a)}
1c	3525	2.37	3.21	3.90	6.01 (OCH ₂ O)
1d	3200—3500	2.38	3.37	3.75 (2×OMe), 3.80	
1e	3530	2.32	3.17	3.67, 3.73, 3.83	
1f	3250, ^{b)} 3560 ^{b)}	2.31	3.09	3.68 (2×OMe), 3.82 (2×OMe)	6.19 (3'-and 5'-H), 6.55 (5-H)

a) Doublet, *J*=8.6 Hz.

b) Broad absorption.

TABLE III. Microanalytical or High Resolution Mass Spectral Data for New Compounds

Compound	Formula	Molecular weight	Analysis (%)					
			Calcd			Found		
			C	H	N	C	H	N
1a	C ₁₉ H ₂₃ NO ₄	329.38	69.28	7.04	4.25	69.24	7.19	4.07
1b	C ₂₀ H ₂₅ NO ₅	359.41	66.83	7.01	3.90	66.66	7.09	3.76
1c	C ₁₈ H ₁₉ NO ₄			313.1314			313.1306 ^{b)}	
1d	C ₁₉ H ₂₃ NO ₄	331.40	69.28	7.04	4.25	69.08	7.10	4.09
1e^{a)}	C ₂₂ H ₂₈ NO ₅ ⁺ ·I ⁻			386.1967			386.1956 ^{b)}	
1f	C ₂₀ H ₂₅ NO ₅ ·0.25H ₂ O	363.91	66.00	7.06	3.85	65.91	7.03	3.77

a) Methiodide of the acetate.

b) High resolution mass spectral data measured with a Hitachi model RMU-6E mass spectrometer.

corypalline is clearly due to the anisotropic effect induced by the newly introduced C-8 phenyl group.

The position of substitution in the present reaction was also clarified by a deuterium-exchange experiment. When corypalline (**3**) was heated in 1*N* NaOD-D₂O at 100°C for 8 h, the signal intensity at δ 6.29 was reduced markedly. Thus, 43 % of the original hydrogen was replaced by deuterium. On the other hand, 8-(2,3,4-trimethoxyphenyl)corypalline (**1b**), which was prepared analogously from **4** and 1,2,3-trimethoxybenzene, was found to have no replaceable hydrogen by the same treatment as above. Accordingly, the absence of hydrogen *ortho* to the C-7 phenolic group in **1b** was confirmed unequivocally.

p-Quinol acetate (**4**) reacted with corypalline (**3**) to give a dimer (**5**), which was identical with an authentic sample⁷⁾ obtained by the phenol coupling of **3**. This reaction further verified the position of substitution, confirming the above assignment.

The intermediate in the present reaction was supposed to be an *o*-quinonoid cation (6),⁵⁾ which was attacked immediately by the activated benzene nucleophile.

A variety of phenyl ethers could be used as the nucleophile with satisfactory results. Yields and spectral data of the products are shown in Tables I and II.

The similar reaction of 4 with other phenolic 1,2,3,4-tetrahydroisoquinolines is currently being investigated in order to obtain mixed dimers.

Experimental

All melting points were measured on a Büchi melting point apparatus and are uncorrected. NMR spectra were taken with a JEOL model JNR-4H-100 (100 MHz) or a Hitachi model R-24B instrument in CDCl₃ solution with Me₄Si as an internal standard. Infrared (IR) spectra were run on a Hitachi model 215 spectrophotometer in CHCl₃ solution. Preparative thin-layer chromatography (TLC) was performed on Silica gel HF₂₅₄ (Merck). Microanalytical data for all new compounds are listed in Table III.

General Procedure for the Preparation of 8-Aryl-1,2,3,4-tetrahydroisoquinolines (1)—Trifluoroacetic acid (0.5 ml) was added to a CH₂Cl₂ (10 ml) solution of an aryl ether (1.5 eq) and the *p*-quinol acetate (4) obtained from corypalline (3) (100 mg) as described previously,⁶⁾ and the whole was stirred at room temperature for 1 h. The reaction mixture was washed with sat. NaHCO₃ aq. and brine, and the CH₂Cl₂ solution was dried over anhydrous K₂CO₃. Evaporation of the solvent under reduced pressure gave a crude product which was purified by preparative TLC (developing solvent; CHCl₃:MeOH=10:1).

Preparation of Corypalline Dimer (5)—Trifluoroacetic acid (1 ml) was added to a stirred mixture of corypalline (3) (100 mg) and *p*-quinol acetate (4) obtained from 3 (100 mg) in CH₂Cl₂ (20 ml) and stirring was continued at room temperature for 1 h. Usual work-up gave an amorphous mass (188 mg), which was separated by preparative TLC (developing solvent; CHCl₃:MeOH=20:3) to yield recovered corypalline (3) (55 mg), mp 164–166°C, from the upper zone, and the dimer (5) (43 mg, 22%), mp 227–229°C (iso-PrOH), from the lower zone. The latter product was shown to be identical with an authentic sample⁷⁾ by mixed melting point determination and spectral comparison.⁸⁾

Acknowledgement The authors are indebted to Dr. T. Moroe of Takasago Perfumery Co., Ltd. for his kind supply of the starting material. Thanks are also due to Miss Y. Katoh for her technical assistance, to Sankyo Co., Ltd. for elemental analyses, and to Miss N. Sawabe and Mrs. F. Hasegawa of this Faculty for NMR and mass spectral measurements.

References and Notes

- 1) Part XIX: H. Hara, O. Hoshino, T. Ishige, and B. Umezawa, *Chem. Pharm. Bull.*, **29**, 1083 (1981).
- 2) A preliminary communication has appeared: H. Hara, O. Hoshino, and B. Umezawa, *Heterocycles*, **15**, 911 (1981).
- 3) J. Sam, R.M. Shafik, and K. Aparajithan, *J. Pharm. Sci.*, **59**, 59 (1970); W. L. Mendelson, C. B. Spainhour, Jr., S.S. Jones, B.L. Lam, and K.L. Wert, *Tetrahedron Lett.*, **21**, 1393 (1980).
- 4) Y. Ahmad and D.H. Hey, *J. Chem. Soc.*, **1961**, 3882; C.R. Ellefson, *J. Org. Chem.*, **44**, 1533 (1979); L.K. Dyall and C.J. Pullin, *Aust. J. Chem.*, **32**, 345 (1979); L.N. Pridgen, *J. Heterocyclic Chem.*, **17**, 1289 (1980); O.R. Andresen and E.B. Pedersen, *Heterocycles*, **19**, 1467 (1982).
- 5) H. Hara, O. Hoshino, and B. Umezawa, *Chem. Pharm. Bull.*, **29**, 51 (1981).
- 6) B. Umezawa, O. Hoshino, Y. Terayama, K. Ohyama, Y. Yamanashi, T. Inoue, and T. Toshioka, *Chem. Pharm. Bull.*, **19**, 2138 (1971).
- 7) B. Umezawa, O. Hoshino, H. Hara, and J. Sakakibara, *Chem. Pharm. Bull.*, **16**, 381 (1968).
- 8) NMR in 1N NaOD-D₂O.