

## Communications to the Editor

[Chem. Pharm. Bull.]  
31(3)1112-1114(1983)

ISOLATION AND STRUCTURE OF MAGNOSALIN AND MAGNOSHININ,  
NEW NEOLIGNANS FROM MAGNOLIA SALICIFORIA MAXIM

Tohru Kikuchi,<sup>\*,a</sup> Shigetoshi Kadota,<sup>a</sup> Kazuo Yanada,<sup>a</sup> Ken Tanaka,<sup>a</sup>  
Kazuo Watanabe,<sup>a</sup> Masao Yoshizaki,<sup>b</sup> Toshio Yokoi,<sup>c</sup> and Tetsuro Shingu<sup>c</sup>

Research Institute for Wakan-Yaku (Oriental Medicines), Toyama Medical  
and Pharmaceutical University,<sup>a</sup> 2630 Sugitani, Toyama 930-01, Japan,  
Faculty of Pharmaceutical Sciences, Toyama Medical and Pharmaceutical  
University,<sup>b</sup> 2630 Sugitani, Toyama 930-01, Japan and  
Faculty of Pharmaceutical Sciences, Kobe-Gakuin University,<sup>c</sup>  
Ikawadani-cho, Tarumi-ku, Kobe 673, Japan

The structures of magnosalin and magnoshinin, new neolignans isolated  
from buds of *Magnolia salicifolia* MAXIM., were determined to be **1** and  
**4**, respectively, on the basis of chemical and spectroscopic evidence.

KEYWORDS — magnosalin; magnoshinin; neolignan; Shin-i; *Magnolia  
salicifolia*

Dried buds of several *Magnolia* species are well known as a crude drug "Shin-i"  
and used in some Chinese medical prescriptions, especially for nasal empyema.  
Chemical investigation on their constituents has been done by several authors, but  
it mainly concerned so far with the essential oils.<sup>1)</sup> A few lignans<sup>2)</sup> and an  
unidentified alkaloid<sup>3)</sup> were also reported.

In Japan, mostly the buds of *Magnolia salicifolia* MAXIM. (Japanese name: tamu-  
shiba) are used as "Shin-i".<sup>4)</sup> In the course of our search for biologically active  
substances from this plant, we isolated four phenolic alkaloids<sup>5)</sup> and two new neo-  
lignans, named magnosalin and magnoshinin. The present communication describes the  
isolation and structure elucidation of the latter compounds.

Air-dried plant material (1 Kg), collected at Toga-mura District, Toyama Prefec-  
ture, was roughly pulverized and extracted with ether at room temperature and then  
with boiling MeOH. Each extract was separated into basic, acidic, and neutral frac-  
tions in the usual manner. The neutral fraction of the MeOH extract, after removal  
of the essential oil by steam distillation, was subjected to alkaline hydrolysis and  
the neutral fraction of the hydrolyzate was roughly separated by silica gel column  
chromatography using hexane, hexane-CH<sub>2</sub>Cl<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, and MeOH-CH<sub>2</sub>Cl<sub>2</sub>. The CH<sub>2</sub>Cl<sub>2</sub>  
eluate was further separated by preparative TLC on Merck Kieselgel 60 PF<sub>254</sub> with  
benzene to give magnosalin (**1**) (140 mg) and magnoshinin (**4**) (105 mg).

Magnosalin (**1**), mp 98-99°C (from ether-hexane),  $[\alpha]_D^{20}$  0° (CHCl<sub>3</sub>),<sup>6)</sup> has the mole-  
cular formula C<sub>24</sub>H<sub>32</sub>O<sub>6</sub> and showed IR absorptions at 1612, 1600, and 1515 cm<sup>-1</sup> and UV  
bands at 208, 231, and 298 nm (log ε: 4.61, 4.23, and 4.03). Its mass spectrum showed  
characteristic peaks at m/z 208 (base peak, C<sub>12</sub>H<sub>16</sub>O<sub>3</sub><sup>+</sup>) and 360 (M<sup>+</sup> - CH<sub>3</sub>CH=CHCH<sub>3</sub>) along  
with the molecular ion peak at m/z 416. The NMR spectrum exhibited signals at δ 1.17

(6H, diffused d,<sup>7)</sup>  $J=6$  Hz,  $\text{sec-CH}_3 \times 2$ ), 1.75 (2H, m,  $\text{CH}_3\text{-CH-CH} \times 2$ ), and 3.27 (2H, diffused d,<sup>7)</sup>  $J=9$  Hz,  $\text{Ar-CH-CH} \times 2$ ) as shown in Fig. 1, suggesting a symmetric feature of its structure. Decoupling and NOE experiments between the signals at  $\delta$  1.75 and  $\delta$  3.27 clearly indicated that the hydrogens concerned are in vicinal, *cis*-arrangement. Furthermore, NOE's were observed between methoxys, aromatic hydrogens, and benzylic hydrogens (Fig. 1).

From the above observations, two possible structures, **1** (racemate) and **2**, can be deduced. Here, we propose the structure **1** for magnosalin, because a serious steric strain may be expected in the alternative structure **2**.

It should be noted that recently Yamamura et al.<sup>8)</sup> isolated a novel neolignan, heterotropan, from *Heterotropa takaoi* MAEKAWA and established its structure to be **3**. Magnosalin, a stereoisomer of **3**, is the second example of this class of neolignan having a cyclobutane system.





Fig.1. NMR (Normal, INDOR, and NOE) Spectra of Magnosalin ( $\text{CDCl}_3$ , 90 MHz)



Magnoshinin (**4**), mp 113.5–115°C (from ether),  $[\alpha]_D^{20}$  0° ( $\text{CHCl}_3$ ),<sup>6)</sup> has the molecular formula  $\text{C}_{24}\text{H}_{30}\text{O}_6$  and showed UV absorptions at 232, 260<sub>sh</sub>, 269, 278, 292<sub>sh</sub>,

310, and 320 nm (log  $\epsilon$ : 4.32, 4.11, 4.21, 4.22, 3.96, 3.95, and 3.92), suggesting the presence of a styrene chromophore. Its mass spectrum showed significant peaks at  $m/z$  414 (base peak,  $M^+$ ), 399, 383 ( $M^+ - OCH_3$ ), 368, 246 ( $a$ ,  $M^+ - C_9H_{12}O_3$ ), and 231.<sup>9)</sup> The NMR spectrum ( $CDCl_3$ ) exhibited signals at  $\delta$  1.10 (3H, d,  $J=6$  Hz,  $sec-CH_3$ ), 1.76 (3H, d,  $J=1.5$  Hz,  $CH=C-CH_3$ ), 2.27 (1H, dq,  $J=0.8, 6$  Hz,  $CH-CH-CH_3$ ), 3.44, 3.61, 3.97 (each 3H, s,  $OCH_3 \times 3$ ), 3.90 (9H, s,  $OCH_3 \times 3$ ), 4.70 (1H, d,  $J=0.8$  Hz,  $Ar-CH-CH$ ), 6.33, 6.49, 6.59 (each 1H, s, aromatic H), and 6.57 (1H, q,  $J=1.5$  Hz,  $Ar-CH=C-CH_3$ ) and their splitting patterns were easily analyzed by the decoupling method.

The above data, coupled with the biogenetic consideration, led us to suppose that the structure of magnoshinin might be **4**, in which the stereochemistry of 1,2-substituents may be *trans*-diaxial from the coupling constant (0.8 Hz) between 1- and 2-hydrogens.<sup>10)</sup>

In order to confirm the substitution pattern of methoxyl groups on the dihydronaphthalene nucleus, we next examined the degradation of **4**. Osmium tetroxide oxidation of **4**, followed by  $HIO_4$  oxidation, afforded a keto-aldehyde (**5**),  $C_{24}H_{30}O_8$ , mp 131–132°C, IR (KBr)  $\nu_{max}$ : 1700 and 1670  $cm^{-1}$ , NMR ( $CDCl_3$ ):  $\delta$  2.08 (3H, s,  $COCH_3$ ), 6.40, 6.44, 7.18 (each 1H, s, aromatic H), and 10.62 (1H, s,  $CHO$ ).

Selective decarbonylation of **5** with tris(triphenylphosphine)rhodium chloride<sup>11)</sup> in benzene gave a decarbonyl compound (**6**) in 12% yield, oil,  $C_{23}H_{30}O_7$  ( $M^+$ : 418.2035. Calcd: 418.1992), NMR ( $CDCl_3$ ):  $\delta$  1.01 (3H, d,  $J=7$  Hz,  $sec-CH_3$ ), 2.04 (3H, s,  $COCH_3$ ), 3.39 (1H, dq,  $J=11.5, 7$  Hz,  $CH-CH_3$ ), 3.62, 3.76 (each 3H, s,  $OCH_3 \times 2$ ), 3.78, 3.85 (each 6H, s,  $OCH_3 \times 4$ ), 4.99 (1H, d,  $J=11.5$  Hz,  $Ar-CH-CH$ ), 6.34, 6.58 (each 1H, d,  $J=3$  Hz, aromatic H), 6.50, and 6.78 (each 1H, s, aromatic H).

Thus the structure of magnoshinin was established to be **4** (racemate).<sup>12)</sup>

Biological activities of **1** and **4** are currently under investigation.

#### REFERENCES AND NOTES

- 1) M. Nagasawa, T. Murakami, K. Ikeda, and Y. Hisada, *Yakugaku Zasshi*, **89**, 454 (1969) and references cited therein.
- 2) H. Kakisawa, T. Kusumi, H. Y. Hsu, and Y. P. Chen, *Bull. Chem. Soc. Japan*, **43**, 3631 (1970).
- 3) M. Kimura, M. Yoshizaki, I. Muro, and D. Shiho, *Yakugaku Zasshi*, **85**, 570 (1965).
- 4) K. Kimura, K. Hata, and M. Yoshizaki, *Shoyaku-gaku Zasshi*, **15**, 50 (1961).
- 5) H. Watanabe, M. Ikeda, K. Watanabe, and T. Kikuchi, *Planta Medica*, **42**, 213 (1981); I. Kimura, M. Kimura, M. Yoshizaki, K. Yanada, S. Kadota, and T. Kikuchi, *ibid.*, in press.
- 6) The CD curves of **1** and **4** showed no Cotton effect. We are grateful to Dr. H. Ishii and Dr. T. Ishikawa, Chiba University, for CD measurements.
- 7) Complex pattern may be due to virtual coupling.
- 8) S. Yamamura, M. Niwa, M. Nonoyama, and Y. Terada, *Tetrahedron Lett.*, 4891 (1978). We are grateful to Dr. S. Yamamura, Keio University, for the copies of NMR and IR spectra of heterotropan for comparison.
- 9) We are indebted to Dr. A. Kato, Niigata College of Pharmacy, for high-resolution MS measurement of magnoshinin.
- 10) F. D. Hostettler and M. K. Seikel, *Tetrahedron*, **25**, 2325 (1969); A. F. A. Wallis, *Tetrahedron Lett.*, 5287 (1968); *idem*, *Aust. J. Chem.*, **26**, 585 (1973) and references cited therein.
- 11) J. Tsuji and K. Ohno, *Tetrahedron Lett.*, 3969 (1965).
- 12) The 5,6,7-trisubstituted structure is excluded from the biogenetic point of view.

(Received January 24, 1983)