

[Chem. Pharm. Bull.]
31(4)1222-1227(1983)

The Reaction of 6-Phenylthiouridine with Sulfur Nucleophiles: A Simple and Regiospecific Preparation of 6-Alkylthiouridines and 6-Alkylthiouridylic Acids

HIROMICHI TANAKA,^a SHUJI IIJIMA,^a AKIRA MATSUDA^a HIROYUKI HAYAKAWA,^a
TADASHI MIYASAKA,^{*,a} and TOHRU UEDA^b

School of Pharmaceutical Sciences, Showa University,^a Hatanodai 1-5-8, Shinagawa-ku, Tokyo 142, Japan and Faculty of Pharmaceutical Sciences, Hokkaido University,^b Kita-12, Nishi-6, Kita-ku, Sapporo 060, Japan

(Received October 1, 1982)

6-Alkylthiouridines and 6-alkylthiouridylic acids were synthesized from the corresponding 6-phenylthiouridine derivatives *via* a regiospecific nucleophilic reaction.

Keywords—uridine; 6-alkylthiouridine; 6-alkylthiouridylic acid; regiospecific reaction; ¹H-NMR

For the preparation of pyrimidine nucleosides, the ribosidation method has been extensively studied.¹⁾ This method is known to be applicable to C-5 substituted pyrimidines but, in general, not to C-6 substituted ones. In the case of the latter, the preferential formation of the N-3 ribosylated product usually takes place, presumably due to the steric hindrance of the substituent in the C-6 position. For example, Winkley and Robins noted that treatment of the bistrimethylsilyl derivative of 6-methylthiouracil with 2,3,5-tri-*O*-benzoyl- β -D-ribofuranosyl bromide in acetonitrile gave 6-methylthio-3-(2,3,5-tri-*O*-benzoyl- β -D-ribofuranosyl)uracil as the sole product in 72% yield.²⁾ To eliminate such undesired reactions, there is a need for a procedure to transform naturally occurring pyrimidine nucleosides to the C-6 substituted analogues.

We recently reported³⁾ that the lithiation of 2',3'-*O*-isopropylidene-5'-*O*-methoxymethyluridine with lithium diisopropylamide occurs at its C-6 position in a regiospecific manner and that the reaction of the resulting dianion with various types of electrophiles represented a new and general route to 6-substituted uridines.

Among the electrophiles employed in this reaction, diphenyl disulfide was quite effective, producing 2',3'-*O*-isopropylidene-5'-*O*-methoxymethyl-6-phenylthiouridine (**1**) in high yield.³⁾ On the other hand, the reaction with an alkyl counterpart such as dimethyl disulfide gave a complex mixture of products, from which the 6-methylthio derivative (**2**) was isolated in only 9.3% yield.⁴⁾ This prompted us to devise an alternative route to the 6-alkylthiouridine derivatives. In this paper, we describe an efficient method for synthesizing 6-alkylthiouridines and 6-alkylthiouridylic acids which have, so far, been difficult to obtain.

A fairly recent study by one of the authors showed that the 5-bromo uridine derivative **3** is susceptible to nucleophilic reaction with benzyl mercaptan to yield the 6-benzylthio (**6**):




Fig. 1




Fig. 2





Chart 1

34%) and 5-benzylthio (7: 40%) derivatives.⁵⁾ The formation of 7 could be attributed to nucleophilic displacement ($4 \rightarrow 5$) in the intermediate 4 which competes with elimination of hydrogen bromide from 4 (Chart 1). Based on the above mechanistic interpretation, we reasoned that a uridine derivative bearing a suitable leaving group at the C-6 position should furnish the desired 6-sulfur substituted product exclusively because displacement at C-5 by the sulfur nucleophile could be avoided in the 5,6-dihydro intermediate, and it appeared that the 6-phenylthiouridine derivative 1 would fulfil this requirement.

We first examined the reaction of 1 with methylmercaptide anion. When 1 in *N,N*-dimethylformamide (DMF) was treated with 5 eq of aqueous sodium methylmercaptide (sodium methanethiolate) at room temperature for 1.5 h, the 6-methylthio derivative (2) was obtained in 91.5% yield after extraction with ethyl acetate followed by chromatographic purification on a silica gel column. The ^1H -nuclear magnetic resonance (NMR) spectrum of 2 (CDCl_3 δ : H-1' 6.07 doublet; H-5 5.49 singlet; SMe 2.48 singlet) was consistent with the expected structure and indistinguishable from that of a previously prepared sample.⁴⁾ As expected, the C-5 substituted product was hardly detectable by careful thin-layer chromatographic (TLC) analysis (benzene: AcOEt=1:1) and ^1H -NMR study (100 MHz, in CDCl_3) of



the crude **2**. The structure of **2** was further confirmed by treating it with 50% aqueous trifluoroacetic acid to give the corresponding free nucleoside **8** (80.8%) whose physical data have been reported.⁵⁾ Under similar conditions, a crystalline 6-mercaptouridine derivative **9** (mp 190–191°C) was also prepared from **1** in quantitative yield by the use of sodium hydrosulfide. This provides a more convenient method for the 6-mercaptouridine derivative than the previously reported sulphydrolysis of 6-benzylthiouridine⁵⁾ and 6-methylaminouridine⁶⁾ derivatives.

The free 6-phenylthiouridine (**10**)³⁾ also followed this reaction course, providing another route to the free 6-alkylthiouridines. The reactions of **10** with alkyl mercaptans were carried out in pyridine in the presence of triethylamine at room temperature. Under these conditions, the 6-ethylthio (**11**), 6-benzylthio (**12**), and 6-butylthio (**13**) derivatives were obtained in 90.5, 77.7, and 93.8%, respectively. We believe that this constitutes an effective route to the free 6-alkylthiouridines, since only mild reaction conditions and simple isolation procedures (evaporation of the solvent and washing of the residue with benzene) are required. Functionalized mercaptans such as β -mercaptoethanol and ethyl thioglycolate also appear to be effective, providing **14** and **15** in good yields. In contrast, the reaction of **10** with *tert*-butyl and isobutyl mercaptans failed. Steric repulsion between the 6-phenylthio group and these rather bulky reagents when adding to the 5,6-double bond is presumably responsible for the above failures.

We next investigated the displacement of the 6-phenylthio group at the nucleotide level, since the reaction proceeded under very mild conditions as mentioned above. Selective phosphorylation of the 5'-hydroxyl group in **10** was achieved according to the published procedure⁷⁾ to give 6-phenylthio-5'-UMP (**16**) in 62% yield. When **16** was treated with sodium hydrosulfide in DMF at room temperature, the desired 6-mercapto-5'-UMP (**17**) was obtained in 92% yield. Further derivatization of **17** was carried out by alkylation with methyl iodide to provide 6-methylthio-5'-UMP (**18**).




Fig. 3

TABLE I. ¹H-NMR Data for 6-Alkylthiouridines in DMSO-*d*₆ (after addition of D₂O)

C-6 substituent	H-5	H-1' (<i>J</i> _{1',2'})	H-2'	H-3'	H-4'	CH ₂ -5'	Other protons
SCH ₃ (8)	5.45	5.67 (3.4 Hz)	4.57	4.07	3.68	~3.39	CH ₃ 2.50
SPh (10)	4.67	5.79 (3.9 Hz)	4.65	4.12	3.73	~3.40	Ph 7.63
SCH ₂ CH ₃ (11)	5.53	5.69 (3.9 Hz)	4.57	4.01	3.71	~3.36	CH ₂ 3.04, CH ₃ 1.32
SCH ₂ Ph (12)	5.61	5.69 (3.9 Hz)	4.57	4.06	3.67	~3.40	Ph 7.48–7.31, CH ₂ 4.33
S(CH ₂) ₃ CH ₃ (13)	5.53	5.72 (3.9 Hz)	4.57	4.08	3.68	~3.40	SCH ₂ 3.02, (CH ₂) ₂ 1.72–1.31, CH ₃ 0.92
SCH ₂ CH ₂ OH (14)	5.58	5.73 (3.4 Hz)	4.57	4.07	— ^{a)}	~3.39	SCH ₂ 3.12, CH ₂ OH — ^{a)}
SCH ₂ CO ₂ C ₂ H ₅ (15)	5.52	5.74 (3.9 Hz)	4.59	— ^{a)}	3.70	~3.36	SCH ₂ 4.08, CO ₂ C ₂ H ₅ 4.17 and 1.22
CH ₃	5.58	5.47 (3.9 Hz)	4.57	4.09	3.69	~3.42	CH ₃ 2.28
H	5.68	5.78 (4.9 Hz)	4.11	~3.90	— ^{a)}	~3.50	H-6 7.88

^{a)} Not resolved.

The ¹H-NMR data for the free 6-alkylthiouridines (**8** and **11–15**) involved in the present study are listed in Table I, together with those for 6-phenylthiouridine (**10**), 6-methyluridine, and uridine. It is well known from crystallographic⁸⁾ and NMR^{9,10)} studies that 6-substituted pyrimidine nucleosides prefer the *syn*-conformation: the C-2 carbonyl is situated on the same side as the ribose ring. All the 6-alkylthiouridines, including 6-phenylthiouridine (**10**), listed in Table I display ¹H-NMR features similar to those of 6-methyluridine. That is, their H-2' and H-3' signals are located downfield compared to those of uridine. This is a well-known charac-

teristic of nucleosides in the *syn*-conformation.^{5,10} As can be seen from Table I, change of the alkyl group in the C-6 substituent of these nucleosides has little effect on the chemical shifts of H-5 with the exception of **10**, in which H-5 appears at higher field by *ca.* 0.9 ppm. Examination of a molecular model strongly suggested that the H-5 in **10** can take a position directly above the phenyl ring, and thus, should be shielded by the ring current effect. Our previous finding that the *endo* methyl signal of the isopropylidene group in 2',3'-*O*-isopropylidene-6-phenylthiouridine is not affected by an anisotropic effect of the C-6 substituent¹¹ provides further support for the above conformation of **10**.

Experimental

Melting points were determined with a Yanagimoto micro melting point apparatus and are uncorrected. ¹H-NMR spectra were measured with a JEOL JNM-FX 100 NMR spectrometer with an appropriate internal standard: tetramethylsilane (TMS) or sodium 2,2-dimethyl-2-silapentane-5-sulfonate (DSS). The abbreviations used are as follows: s, singlet; d, doublet; dd, double doublet; t, triplet; q, quartet; m, multiplet; br, broad. Mass spectra (MS) were taken on a JEOL JMS-D 300 spectrometer. Ultraviolet (UV) spectra were recorded on Hitachi 340 or Shimadzu UV-240 spectrometer. Column chromatography was carried out on Merck Silica Gel 60 or Florisil (Floridin, Inc.). Column chromatographic purification of 5'-phosphates was performed on DEAE-Cellulofine AL (Chisso, Inc.). TLC was performed on silica gel (Precoated silica gel plate F₂₅₄, Merck) or Avicel SF (Funakoshi Pharmaceutical, Inc.).

2',3'-*O*-Isopropylidene-5'-*O*-methoxymethyl-6-methylthiouridine (2)—To a suspension of **1** (500 mg, 1.15 mmol) in 15% aqueous NaSMe (2.69 ml, 5.75 mmol), 0.3 ml of DMF was added. The resulting solution was stirred for 1.5 h at room temperature. After being quenched with AcOH (0.33 ml), the reaction mixture was poured into saline. Extraction with AcOEt followed by chromatographic purification on a silica gel (10 g) column (1% EtOH in CHCl₃) gave 394 mg (91.5%) of **2** as a syrup. MS *m/z*: 374/(M⁺), 359 (M-15), 158 (B+1). NMR (CDCl₃) δ: 1.35 (3H, s, isop.Me), 1.56 (3H, s, isop.Me), 2.48 (3H, s, SMe), 3.36 (3H, s, CH₂OCH₃), 3.72–3.79 (2H, m, CH₂-5'), 4.20–4.37 (1H, m, H-4'), 4.65 (2H, s, CH₂OCH₃), 4.88 (1H, dd, H-3'), 5.22 (1H, dd, H-2'), 5.49 (1H, s, H-5), 6.07 (1H, d, *J*=1.5 Hz, H-1'), 8.02 (1H, br, NH).

6-Methylthiouridine (8)—Treatment of **2** (276 mg) with 50% aqueous CF₃COOH (5 ml) for 2 d at room temperature followed by chromatographic purification (10% EtOH in CHCl₃) gave **8** (173 mg, 80.8%), which was crystallized from MeOH: mp 171.5–172.5°C (resolidified at 173.5°C, dec. at 223°C); lit.⁵¹ mp 163–165.5°C (resolidified at 166°C, dec. at 230°C). NMR: see Table I.

2',3'-*O*-Isopropylidene-5'-*O*-methoxymethyl-6-mercaptopuridine (9)—A DMF (5 ml) solution of **2** (500 mg, 1.15 mmol) and 70% NaSH (200 mg, 2.50 mmol) was stirred for 2.5 h at room temperature. After being neutralized with Amberlite IR-120-B (H⁺ form), the mixture was evaporated to dryness. Chromatographic purification on Florisil (50% MeOH in CH₂Cl₂) gave 413 mg (100%) of **9**. Crystallization from isopropanol gave an analytical sample: mp 190–191°C. Anal. Calcd for C₁₄H₂₀N₂O₇S: C, 46.66; H, 5.59; N, 7.77. Found: C, 46.42; H, 5.60; N, 7.53. MS *m/z*: 361 (M+1), 345 (M-15), 144 (B+1). UV λ_{max}^{methanol} nm (ε): 316 (23300) and 238 (8400), λ_{min}^{methanol} nm (ε): 263 (600) and 222 (5500). NMR (DMSO-*d*₆) δ: 1.26 (3H, s, isop. Me), 1.46 (3H, s, isop. Me), 3.24 (3H, s, CH₂OCH₃), 3.40–3.74 (2H, m, CH₂-5'), 3.85–4.04 (1H, m, H-4'), 4.53 (2H, s, CH₂OCH₃), 4.69 (1H, dd, H-3'), 4.97 (1H, d, H-2'), 5.30 (1H, d, H-5), 7.66 (1H, s, H-1'), 10.13 (1H, br, NH).

6-Phenylthiouridine (10)—An aqueous 50% CF₃CO₂H solution (30 ml) containing **1** (3.0 g, 6.87 mmol) was stirred for 2 d at room temperature. The mixture was then evaporated to dryness and the resulting residue was chromatographed on a silica gel column (5% EtOH in CHCl₃) to afford **10** (2.07 g, 85.5%). Crystallization from EtOAc gave an analytical sample: mp 156–159°C. Anal. Calcd for C₁₅H₁₆N₂O₆S: C, 51.13; H, 4.58; N, 7.95. Found: C, 51.08; H, 4.64; N, 7.77. UV λ_{max}^{methanol} nm (ε): 282 (11000), λ_{min}^{methanol} nm (ε): 245 (5400). NMR: see Table I.

6-Ethylthiouridine (11)—Et₃N (1.6 ml) and EtSH (0.85 ml) were added to a pyridine (3 ml) solution of **10** (403 mg, 1.14 mmol). After being stirred overnight at room temperature, the reaction mixture was evaporated to dryness. The resulting residue was washed with benzene to leave 314 mg (90.5%) of **11**. Crystallization from H₂O gave an analytical sample: mp 190.5–191.5°C. Anal. Calcd for C₁₁H₁₆N₂O₆S: C, 43.41; H, 5.30; N, 9.21. Found: C, 43.20; H, 5.27; N, 9.31. MS *m/z*: 286 (M-H₂O), 172 (B+1). UV λ_{max}^{H₂O} nm (ε): 283 (14700), λ_{min}^{H₂O} nm (ε): 249 (3300). NMR: see Table I.

6-Benzylthiouridine (12)—Et₃N (1.24 ml) and benzyl mercaptan (1.0 ml) were added to a pyridine (3 ml) solution of **10** (302 mg, 0.86 mmol). After being stirred overnight at room temperature, the reaction mixture was evaporated to dryness. The resulting residue was washed with benzene to leave 253 mg (77.7%) of **12**. Crystallization from EtOH gave pure **12**: mp 174.5–176°C; lit.⁵¹ mp 169–170°C, resolidified at 172°C, dec. at 225°C. NMR: see Table I.

6-Butylthiouridine (13)—Et₃N (1.5 ml) and butyl mercaptan (1.2 ml) were added to a pyridine (3 ml)

solution of **10** (385 mg, 1.09 mmol). After being stirred for 2 d at room temperature, the reaction mixture was evaporated to dryness. The resulting residue was washed with benzene to leave 340 mg (93.8%) of **13**. Crystallization from EtOH gave an analytical sample: mp 148.5–150°C. *Anal.* Calcd for $C_{13}H_{20}N_2O_6S$: C, 46.97; H, 6.07; N, 8.43. Found: C, 46.72; H, 6.05; N, 8.59. UV $\lambda_{\max}^{\text{methanol}}$ nm (ϵ): 279 (13000), $\lambda_{\min}^{\text{methanol}}$ nm (ϵ): 247.7 (3400). NMR: see Table I.

6-(β -Hydroxy)ethylthiouridine (14)— Et_3N (1.4 ml) and 2-mercaptoethanol (0.7 ml) were added to a pyridine (3 ml) solution of **10** (366 mg, 1.00 mmol). After being stirred for 2 h at room temperature, the reaction mixture was evaporated to dryness. The resulting residue was chromatographed on a silica gel column (15% EtOH in CHCl_3) to afford 243 mg (75.9%) of **14**. Crystallization from EtOH gave an analytical sample: mp 166–167.5°C. *Anal.* Calcd for $C_{11}H_{16}N_2O_7S$: C, 41.24; H, 5.04; N, 8.75. Found: C, 41.48; H, 5.16; N, 8.63. MS m/z : 302 ($\text{M}-\text{H}_2\text{O}$), 217 ($\text{B}+\text{CHOH}$), 188 ($\text{B}+1$). UV $\lambda_{\max}^{\text{methanol}}$ nm (ϵ): 279.5 (13300), $\lambda_{\min}^{\text{methanol}}$ nm (ϵ): 237.5 (3800). NMR: see Table I.

6-Ethoxycarbonylmethylthiouridine (15)— Et_3N (1.52 ml) and ethyl thioglycolate (1.2 ml) were added to a pyridine (3 ml) solution of **10** (384 mg, 1.09 mmol). After being stirred for 3 h at room temperature, the reaction mixture was evaporated to dryness. The resulting residue was chromatographed on a silica gel column (5% EtOH in CHCl_3). This afforded 301 mg (76.3%) of **15**. Crystallization from EtOH gave hygroscopic crystals: mp 112–114°C. UV $\lambda_{\max}^{\text{methanol}}$ nm: 277.5, $\lambda_{\min}^{\text{methanol}}$ nm: 245. MS m/z : 230 ($\text{B}+1$). NMR: see Table I.

6-Phenylthiouridine 5'-Phosphate (16)—An ice-cooled $(\text{MeO})_3\text{PO}$ (2.5 ml) solution of **10** (324 mg, 0.92 mmol) was treated with 0.17 ml (2 equivalents) of POCl_3 . After being stirred for 6 h at 0°C, the reaction mixture was quenched with saturated NaHCO_3 (10 ml). The aqueous solution was washed with two 10 ml portions of ether, acidified with 1 N HCl to pH 5, diluted with H_2O to a volume of ca. 400 ml, and applied to a column of DEAE-Cellulofine (bicarbonate form, 2.4×26 cm). Elution of **16** was performed with a linear gradient of 0 to 0.4 M triethylammonium bicarbonate (800 ml each). Fractions of 18 ml were collected. Fractions No. 32–41 were combined and concentrated to give the triethylammonium salt of **16** (6330 ODU at 283 nm, 62%, calculated from ϵ of **16**=11000). UV $\lambda_{\max}^{\text{H}_2\text{O}}$ nm: 283, $\lambda_{\min}^{\text{H}_2\text{O}}$ nm: 246. NMR (D_2O) δ : 3.90–4.10 (3H, m, CH_2 -5' and H-4'), 4.46 (1H, t, H-3'), 4.83 (1H, dd, H-2'), 5.23 (1H, s, H-5), 6.08 (1H, d, $J=2.9$ Hz, H-1'), 7.57–7.67 (5H, m, phenyl). *Rf* values: see Table II.

6-Mercaptouridine 5'-Phosphate (17)— NaSH (100 mg) was added to a DMF (10 ml) solution of **16** (triethylammonium salt, 2100 ODU at 283 nm). After being stirred overnight, the reaction mixture was diluted with H_2O (300 ml), acidified with 1 N HCl to pH 4, and bubbled through with N_2 gas to remove H_2S . The resulting cloudy solution was applied to a column of DEAE-Cellulofine (bicarbonate form, 2.4×9 cm). The column was washed with H_2O (500 ml). Elution of **17** was performed with a linear gradient of 0 to 0.4 M triethylammonium bicarbonate, and 15 ml fractions were collected. Fractions No. 28–49 were combined and concentrated to give the triethylammonium salt of **17** (4725 ODU at 317 nm, 92%, calculated from ϵ of **17**=23300). UV $\lambda_{\max}^{\text{H}_2\text{O}}$ nm: 317 and 233, $\lambda_{\min}^{\text{H}_2\text{O}}$ nm: 262 and 220. *Rf* values: see Table II.

6-Methylthiouridine 5'-Phosphate (18)—A 1.2 ml aliquot of 1 N NaOH and 1.0 ml of MeI were added with stirring to a MeOH (20 ml) solution of **17** (triethylammonium salt, 5100 ODU at 317 nm). After 1 h at room temperature, the reaction mixture was diluted with H_2O (300 ml), acidified with 1 N HCl to pH 6, and applied to a column of DEAE-Cellulofine (bicarbonate form, 2.4×7 cm). Elution of **18** was performed with a linear gradient of 0 to 0.3 M triethylammonium bicarbonate (400 ml each). Fractions of 15 ml were collected. Fractions No. 21–39 were combined and concentrated to give the triethylammonium salt of **18** (3170 ODU at 282 nm, 94%, calculated from ϵ of **18**=15400). UV $\lambda_{\max}^{\text{H}_2\text{O}}$ nm: 282, $\lambda_{\min}^{\text{H}_2\text{O}}$ nm: 248. NMR (D_2O) δ : 2.56 (3H, s, SMe), 3.92–4.16 (3H, m, CH_2 -5' and H-4'), 4.45 (1H, t, H-3'), 4.84 (1H, dd, H-2'), 5.65 (1H, s, H-5), 5.98 (1H, d, $J=2.9$ Hz, H-1'). *Rf* values: see Table II.

TABLE II. *Rf* Values of Compounds **16**–**18**

Compound	Solvent system ^{a)}		
	A	B	C
5'-UMP	0.40	0.15	0.06
16	0.63	0.42	0.12
17	0.30	0.08	0.06
18	0.46	0.20	0.08

a) A, EtOH-1M NH_4OAc (5:4); B, EtOH-1M NH_4OAc (5:2);
C, isopropanol- NH_4OH - H_2O (7:1:2).

References and Notes

- 1) For example, H.F. Vorbrüggen, U. Niedballa, K. Krolikiewicz, B. Bennua, and G. Höfle, "Chemistry and Biology of Nucleosides and Nucleotides," ed. by R.E. Harmon, R.K. Robins, and L.B. Townsend, Academic Press, New York, San Francisco, and London, 1978, pp. 251—265, and references cited therein.
- 2) M.W. Winkley and R.K. Robins, *J. Chem. Soc. (C)*, **1969**, 791.
- 3) H. Tanaka, H. Hayakawa, and T. Miyasaka, *Chem. Pharm. Bull.*, **29**, 3565 (1981); *idem*, *Nucleic Acids Symposium Series*, **10**, 1 (1981); *idem*, *Tetrahedron*, **38**, 2635 (1982).
- 4) H. Tanaka and H. Hayakawa, unpublished result.
- 5) H. Inoue, S. Tomita, and T. Ueda, *Chem. Pharm. Bull.*, **23**, 2614 (1975).
- 6) H. Inoue, M. Takada, M. Takahashi, and T. Ueda, *Heterocycles*, **8**, 427 (1977).
- 7) M. Yoshikawa, T. Kato, and T. Takenishi, *Bull. Chem. Soc. Jpn.*, **42**, 3505 (1969).
- 8) D. Suck and W. Saenger, *J. Am. Chem. Soc.*, **94**, 6520 (1972).
- 9) As an example of a ^{13}C -NMR study, see: M.P. Schweizer and G.P. Kreishman, *J. Magn. Reson.*, **9**, 334 (1973).
- 10) As an example of a ^1H -NMR study, see: M.P. Schweizer, E.B. Banta, J.T. Witkowski, and R.K. Robins, *J. Am. Chem. Soc.*, **95**, 3770 (1973).
- 11) H. Tanaka, H. Hayakawa, and T. Miyasaka, *Chem. Pharm. Bull.*, **30**, 1117 (1982).