

Communications to the Editor

[Chem. Pharm. Bull.
31(4)1437-1439(1983)]

REGIOSELECTIVE INTRODUCTION OF p-COUMAROYL GROUP TO α -L-ARABINO-
PYRANOSIDES. TOTAL SYNTHESSES OF INUNDOSIDE-G AND INUNDOSIDE-D₁¹⁾

Yoshisuke Tsuda* and Md. Ekramul Haque

Faculty of Pharmaceutical Sciences, Kanazawa University,
13-1 Takara-machi, Kanazawa 920, Japan

Stannylation of cholesteryl α -L-arabinopyranoside with Bu₂SnO followed by p-coumaroylation gave the 3'-O-p-coumarate, which on heating in pyridine rearranged to the 4'-O-p-coumarate. Similarly inundoside-A was converted to its 3'-O-p-coumarate and then 4'-O-p-coumarate, which were identical with inundoside-G and inundoside-D₁, respectively, thus providing their total syntheses.

KEYWORDS — dibutyltin oxide; regioselective acylation; p-coumaroylation; acyl migration; inundoside-G; inundoside-D₁; triterpenoid glycoside; α -L-arabinopyranoside; Lycopodium inundatum; cholesteryl α -L-arabinopyranoside

Synthesis of naturally occurring acylated poly-hydroxy compounds such as acylated glycosides requires introduction of an acyl group regioselectively to a desired position in non-acylated substrates; that is, the differentiation of one hydroxyl group from the others.

This communication treats the above problem for α -L-arabinopyranosides, choosing cholesteryl α -L-arabinopyranoside (1a)²⁾ and serratenediol 3- α -L-arabinopyranoside (1b: inundoside-A)³⁾ as substrates. p-Coumaroyl group, one of the common acyl groups of shikimate origin, was selected as an acyl moiety. Introduction of p-coumaroyl group to 4'-OH of 1b would lead to the synthesis of inundoside-D₁, an acylated glycoside isolated from Lycopodium inundatum.³⁾ 2'-O-Acylation may be achieved via an O,O-isopropylidene derivative and 3'-O-acylation can be done regioselectively by using either Bu₂SnO or (Bu₃Sn)₂O method.⁴⁾ However, 4'-O-acylation can not be done regioselectively, since 4'-OH of arabinopyranosides is least reactive. We thought this can be done by acyl migration of the 3'-O-acyl group.

Stannylation of cholesteryl α -L-arabinopyranoside (1a) with Bu₂SnO (1.5 eq) in boiling dioxane for 7.5 h followed by acylation with freshly prepared p-acetoxycinnamoyl chloride (1.5 eq) at room temperature for 3 h yielded 3'-O-p-acetoxycinnamate (2a) (50.5%), mp 279-281°C, and 4'-O-p-acetoxycinnamate (3a) (13.5%), mp 157-160°C. The phenolic acetyl group of the former was easily removed solvolytically on treatment with boiling methanol to yield 3'-O-p-coumarate (4a), mp 289-290°C. The 3'-O-acyl group in 2a, when heated in pyridine at 130°C for 3 h, migrated to 4'-O-position, giving rise to an equilibrium mixture of 4'-O- (3a) and 3'-O-ester (2a) in a ratio of 3 : 2. The structures of these products and the

ratio were estimated by ^{13}C -NMR spectra. Chromatographic separation of the product effected the isolation of 4'-O-ester (3a), mp 275-278°C, in a pure form. Treatment of 3a with NaBH_4 in THF smoothly gave the deacetylated compound (5a), mp 289-290°C. 3'-O-p-Coumarate (4a) also migrated, on heating in pyridine at 130°C for 3 h, giving rise to a 2 : 3 mixture of 4a and 5a. Crystallization of the reaction mixture from CHCl_3 -MeOH gave 5a in a pure form.




Table I. ^{13}C -NMR δ_{C} of the Arabinose Moiety for the non-Acylated and Acylated α -L-Arabinopyranosides, 1a, 2a, 3a, 4a, 5a, 1b, 4b, and 5b, and Their Acylation Shift Values (in Parentheses) in Pyridine- d_5

Carbon No.	1a	2a	3a	4a	5a	1b	4b	5b
1'	103.1	103.0 (-0.1)	103.3 (+0.2)	103.0 (-0.1)	103.3 (+0.2)	106.8	106.9 (+0.1)	106.6 (-0.2)
2'	72.5	69.7 (-2.8)	72.7 (+0.2)	69.7 (-2.8)	72.7 (+0.2)	72.8	69.9 (-2.9)	72.8 (0)
3'	74.6	77.2 (+2.6)	72.7 (-1.9)	76.9 (+2.3)	72.3 (-2.3)	74.4	76.7 (+2.3)	73.3 (-1.1)
4'	69.5	67.1 (-2.4)	72.9 (+3.4)	67.2 (-2.3)	72.9 (+3.4)	69.0	66.9 (-2.1)	72.3 (+3.3)
5'	66.8	66.8 (0)	64.6 (-2.2)	66.9 (+0.1)	64.7 (-1.9)	66.1	66.2 (+0.1)	64.5 (-1.6)

Similarly stannylation of inundoside-A (1b) with Bu_2SnO (3.0 eq) in boiling dioxane followed by acylation with p-acetoxy-cinnamoyl chloride and crystallization of the product from hot CHCl_3 -MeOH gave, with concomitant loss of acetyl group, 3'-O-p-coumarate (4b), mp 297-299°C. Acetylation of this gave the tetraacetate, mp >300°C. This tetraacetate was found to be identical with inundoside-G tetraacetate (lit. mp >300°C) (^1H -NMR, IR, and TLC comparisons), for which the position of the p-coumaroyl group had not been established.³⁾

Heating of 3'-O-p-coumarate (4b) in pyridine at 110°C for 3 h gave ca. 1 : 1 equilibrium mixture of 3'-O- and 4'-O-p-coumarate (4b and 5b) with slight excess of the latter (estimated from ^{13}C -NMR). Preparative TLC and several crystallizations of the mixture from CHCl_3 -MeOH afforded 4'-O-p-coumarate (5b), mp 285-287°C, in a pure form, which was identical with inundoside- D_1 (lit. mp 286-290°C)³⁾ (NMR, IR, and TLC comparisons). Acetylation of this gave the tetraacetate, mp 275-276°C, identical with inundoside- D_1 tetraacetate (lit. mp 272-274°C)³⁾ (NMR, IR, and TLC comparisons).

The above syntheses provide not only the rigid proofs of the structures of inundoside- D_1 and inundoside-G but also the total syntheses of these acylated triterpenoid glycosides in a formal sense, since total synthesis of inundoside-A has already been accomplished.⁵⁾

ACKNOWLEDGEMENT One of the authors (Md. E. H.) is grateful to the Ministry of Education, Science and Culture for financial support during the course of this research work.

REFERENCES AND NOTES

- 1) Utilization of Sugars in Organic Synthesis. IX. Part VIII: Y. Tsuda, Md. E. Haque, and K. Yoshimoto, Chem. Pharm. Bull., 31, in press.
- 2) This was prepared by modified König-Knorr reaction of cholesterol and tri-O-acetyl- β -L-arabinopyranosyl bromide followed by methanolysis with NaOMe.
- 3) Y. Tsuda, M. Kaneda, N. Yasufuku, and Y. Shimizu, Chem. Pharm. Bull., 29, 2123 (1981).
- 4) see Part VIII.
- 5) Y. Tsuda and Md. E. Haque, Chem. Pharm. Bull., 30, 1500 (1982).

(Received March 5, 1983)