

[Chem. Pharm. Bull.]
36(9)3638—3641(1988)]

**Studies on Tetrahydroisoquinolines. XXXII.¹⁾ Conversion
of 7-Hydroxy-6-methoxytetrahydroisoquinolines to
6-Hydroxy-7-methoxytetrahydroisoquinolines**

HIROSHI HARA, TOSHIFUMI AKIBA, TAKANORI MIYASHITA,
OSAMU HOSHINO, and BUNSUKE UMEZAWA*

*Faculty of Pharmaceutical Sciences, Science University of Tokyo, 12,
Ichigaya Funagawara-machi, Shinjuku-ku, Tokyo 162, Japan*

(Received February 24, 1988)

The 7-hydroxy-6-methoxytetrahydroisoquinolines (**2** and **8**) were transformed to the corresponding 6-hydroxy-7-methoxytetrahydroisoquinolines (**4** and **14**) via the *p*-quinol acetates (**1** and **9**), the *p*-quinol ethers (**3a** or **3b**, and **10a**), and the 6-alkoxy-7-ols (**5** or **7**, and **13**), respectively.

Keywords—tetrahydroisoquinolinol; *p*-quinol acetate; *p*-quinol ether; reductive aromatization; transposition

We have already reported that the *p*-quinol acetate (**1**), derived from corypalline (**2**) by lead tetraacetate oxidation, reacts with various alcohols in the presence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ or concentrated H_2SO_4 to give rise to the corresponding *p*-quinol ether (**3**), as a result of elimination of the methoxyl group at the C-6 position.²⁾

If the *p*-quinol ether (**3**) has a readily removable alkyl group at the C-6 position, it may be a key precursor leading to the 6-hydroxy-7-methoxytetrahydroisoquinoline (**4**). Here we wish to report a procedure for conversion of 7-hydroxy-6-methoxytetrahydroisoquinolines to 6-hydroxy-7-methoxytetrahydroisoquinolines via the following steps; oxidation, ether-formation, reductive aromatization, methylation of the 7-hydroxyl group, and hydrolysis.

First, a conversion of corypalline (**2**) into the isomer (**4**) via the 4a,6-diallyl *p*-quinol ether (**3a**) was undertaken. Treatment of the *p*-quinol ether (**3a**), which was obtained by the reaction of **1** and allyl alcohol,²⁾ with NaBH_4 followed by heating on a boiling water bath, gave the allyloxy phenol (**5**) in a yield of 88%. The structure of the product was supported by its infrared (IR) and proton nuclear magnetic resonance ($^1\text{H-NMR}$) spectral data.

Methylation of **5** with diazomethane afforded the ether (**6**), then the allyl ether was cleaved by treatment with 10% Pd-C and *p*- $\text{TsOH} \cdot \text{H}_2\text{O}$ ³⁾ to give the expected 6-hydroxy-7-methoxytetrahydroisoquinoline (**4**) in an overall yield of 3.5% from **2**. The product (**4**) was identical with an authentic specimen.⁴⁾ The low overall yield was mainly due to the final step (22% yield). Hence isopropanol was employed for the formation of the *p*-quinol ether (**3**) instead of allyl alcohol. Thus, the *p*-quinol ether (**3b**)²⁾ was reduced with NaBH_4 followed by heating at 100°C (bath temperature) to give the isopropoxy phenol (**7**), which was subjected to methylation and subsequent hydrolysis with 48% HBr in AcOH ⁵⁾ to yield **2a** in an overall yield of 8.2%.

The procedure was applied to the 1-benzyl corypalline (**8**). Namely, the *p*-quinol acetate (**9**),⁶⁾ readily obtained from **8** as a mixture of two diastereoisomers, was treated with isopropanol in the presence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ at room temperature to give a single *p*-quinol ether (**10a** or **10b**) (14%) and a diastereoisomeric mixture of the 4,6-diisopropoxytetrahydroisoquinolines (**11**) (18%) on the basis of $^1\text{H-NMR}$ spectral analysis. This can presumably be explained in terms of the stereochemical 1,3-relationship of the two substituents at C-1 and C-




Chart 1

4a. Namely, two *p*-quinol ethers (**10a** and **10b**) were formed initially. One of these, the 1,4a-*trans* ether (**10a**), remained unchanged, while the other, the 1,4a-*cis* ether (**10b**), was further transformed into the 4,7-diisopropoxy derivatives (**11**) via the *p*-quinone methide (**12**) owing to its instability caused by repulsion between the two bulky groups in the 1,3-relationship.

Reductive aromatization of the *p*-quinol ether (**10a**) gave the 6-isopropoxy derivative (**13**) in 46% yield. The mixture of the diethers (**11**) was also converted to **13** by hydrogenolysis (H₂/10% Pd-C, 6N HCl) in 38% yield. Compound **13** was methylated and subsequently hydrolyzed with 48% HBr in AcOH, giving the 1-benzyl-6-hydroxy-7-methoxytetrahydro-

isoquinoline (**14**) in 55% yield (overall yield of 7.4% from **8**). Compound **14** was identical with an authentic sample⁴⁾ based on a comparison of their spectral data and mixed melting point determination.

Thus, an efficient procedure for transposition of the hydroxyl and methoxyl groups of 7-hydroxy-6-methoxytetrahydroisoquinolines has been developed.

Experimental

All melting points were measured on a Büchi melting point apparatus and are uncorrected. ¹H-NMR spectra were taken with a JEOL JNX-FX-100 (100 MHz) instrument in CDCl₃ solution with Me₄Si as an internal standard unless otherwise noted. IR spectra were run on a Hitachi 260 spectrometer in CHCl₃ solution. Mass spectra (MS) were run on a Hitachi RMU-7M mass spectrometer. Preparative thin layer chromatography (TLC) was performed on Silica gel 60 F₂₅₄ plates (Merck), 2.0 mm thick.

Reaction of the *p*-Quinol Acetate (9**) with Isopropanol in the Presence of Borontrifluoride Etherate**—The *p*-quinol acetate (**9**) was prepared quantitatively by Pb(OAc)₄ oxidation of 1-benzylcorypalline (**8**)⁶⁾ (300 mg), as described previously.⁶⁾ Without purification, **9** was dissolved in a mixture of iso-PrOH (36 ml) and BF₃·Et₂O (3.6 ml). The whole was stirred at room temperature for 2.5 h, and the reaction mixture was poured into 10% aqueous Na₂CO₃ to be alkalized. The product was extracted with CHCl₃, and the extract was washed with brine and dried over K₂CO₃. The solvent was evaporated off under reduced pressure to give an oil (298.3 mg), which was separated by preparative TLC [developing solvent, CHCl₃-AcOEt-MeOH (25:1:1)] to afford the *p*-quinol ether (**10a**) (55.8 mg, 14%) and the diisopropoxyphenol (**11**) (70.8 mg, 18%) (polarity: **10a** < **11**). **10a**: An oil. High-resolution MS, Calcd for C₂₃H₃₀NO₃ (M⁺ - 1): 368.2224. Found: 368.2232. ¹H-NMR δ: 0.93, 1.06, 1.35, 1.37 (each 3H, d, J = 5.7 Hz, OCHMe₂), 2.17 (3H, s, NMe), 3.27, 4.36 (each 1H, septet, J = 5.7 Hz, OCHMe₂), 5.66 (1H, s, 5-H), 6.36 (1H, s, 8-H), 7.30 (5H, s, PhH). IR ν_{max} cm⁻¹: 1665, 1640, 1620 (dienone). **11**: An oil. High-resolution MS, Calcd for C₂₄H₃₁NO₃: 369.2302. Found: 369.2333. ¹H-NMR δ: 1.22, 1.25, 1.37, 1.38 (each 3H, d, J = 5.7 Hz, OCHMe₂), 2.54, 2.58 (3H, each s, 3:5,⁷⁾ NMe), 3.69—3.96 (ca. 1.5H, m, C₄-H and OCHMe₂), 4.22 (ca. 0.5H, t, C₄-H), 4.58 (1H, septet, J = 5.7 Hz, OCHMe₂), 6.24, 6.44 (1H, each s, 3:5,⁷⁾ ArH), 6.88, 6.95 (1H, each s, 5:3,⁷⁾ ArH), 7.00—7.40 (5H, m, ArH). IR ν_{max} cm⁻¹: 3535 (OH).

6-Allyloxy-7-hydroxy-2-methyl-1,2,3,4-tetrahydroisoquinoline (5**)**—Sodium borohydride (7.5 mg) was added to a solution of the *p*-quinol ether (**3a**)²⁾ (36.2 mg) in MeOH (4 ml) and the whole was stirred at room temperature for 2 h. The solvent was evaporated off under reduced pressure and the residue was made alkaline with 10% aqueous Na₂CO₃. The product was extracted with a mixture of CHCl₃ and iso-PrOH (4:1), and the organic layer was washed with brine and dried over K₂CO₃. The solvent was evaporated off under reduced pressure to give a yellow oil (35 mg), which was heated on a boiling water bath under reduced pressure for 2 h to afford an oil (32.5 mg). The oil was purified by preparative TLC [developing solvent CHCl₃-AcOEt-MeOH (6:1:1)] to give the allyloxyphenol (**5**) (25.4 mg, 88%), mp 130—132°C (hexane). High-resolution MS, Calcd for C₁₃H₁₇NO₂: 219.1257. Found: 219.1243. ¹H-NMR δ: 2.42 (3H, s, NMe), 3.48 (2H, s, C₁-H), 4.57 (2H, dt, J = 5.7, 1.4 Hz, OCH₂CH=CH₂), 5.29 [1H, dq, J = 10, 1.4 Hz, OCH₂CH=CH₂ (*trans* H)], 5.39 [1H, dq, J = 17, 1.4 Hz, OCH₂CH=CH₂ (*cis* H)], 6.08 (1H, ddt, J = 17, 10, 5.7 Hz, OCH₂CH=CH₂), 6.56, 6.59 (each 1H, s, ArH). IR ν_{max} cm⁻¹: 3545 (OH).

7-Hydroxy-6-isopropoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline (7**)**—Sodium borohydride (18 mg, 1.5 eq) was added to a solution of the *p*-quinol ether (**3b**)²⁾ (89 mg) in MeOH (6 ml) and the whole was stirred at room temperature for 2 h. Usual work-up gave a yellow oil (90 mg), which was heated on a boiling water bath under reduced pressure for 2 h to afford an oil. The oil was purified by preparative TLC [developing solvent, CHCl₃-AcOEt-MeOH (6:1:1)] to give the isopropoxyphenol (**7**) (38 mg, 54%), mp 114—114.5°C (AcOEt). Anal. Calcd for C₁₃H₁₉NO₂: C, 70.55; H, 8.65; N, 6.33. Found: C, 70.69; H, 8.57; N, 6.25. ¹H-NMR δ: 1.35 (6H, d, J = 5.7 Hz, OCHMe₂), 2.46 (3H, s, NMe), 3.51 (2H, s, 1-H), 4.53 (1H, septet, J = 5.7 Hz, OCHMe₂), 6.58, 6.61 (each 1H, s, ArH), IR ν_{max} cm⁻¹: 3540 (OH).

1-Benzyl-7-hydroxy-6-isopropoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline (13**)**—(a) Sodium borohydride (5 mg) was added to the solution of the *p*-quinol ether (**10a**) (27 mg) in MeOH (4 ml) and the whole was stirred at room temperature for 3.5 h. Usual work-up gave a yellow oil (29 mg), which was heated on a boiling water bath under reduced pressure for 2 h to afford an oil (30 mg). The oil was purified by preparative TLC [developing solvent, CHCl₃-AcOEt-MeOH (15:1:1)] to give **13** (11 mg, 46%), as an oil. High-resolution MS, Calcd for C₂₀H₂₅NO₂: 311.1884. Found: 311.1894. ¹H-NMR δ: 1.36, 1.38 (each 3H, d, J = 5.7 Hz, OCHMe₂), 2.50 (3H, s, NMe), 3.78 (1H, t, J = 5.7 Hz, C₁-H), 4.55 (1H, septet, J = 5.7 Hz, OCHMe₂), 6.39, 6.56 (each 1H, s, ArH), 7.00—7.46 (5H, m, PhH). IR ν_{max} cm⁻¹: 3540 (OH).

(b) The isopropoxyphenol (**11**) (36 mg) was dissolved in 6N HCl and the mixture was shaken with 10% Pd-C (20 mg) in an atmosphere of hydrogen until uptake of the gas ceased. After the catalyst was removed by filtration, the filtrate was alkalized by adding 15N NH₄OH. The product was extracted with CHCl₃, and the extract

was washed with brine and dried over K_2CO_3 . The solvent was evaporated off under reduced pressure to give an oil (43 mg), which was purified by preparative TLC [developing solvent, $CHCl_3$ -AcOEt-MeOH (15:1:1)] to afford the 6-isopropoxy-7-ol (**13**) (12 mg, 38%), as an oil. The product was identical with **13** prepared by procedure (a).

6-Hydroxy-7-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline (4)—(a) A solution of **5** (72 mg) and diazomethane (excess) in ether (30 ml) and MeOH (8 ml) was allowed to stand at room temperature for 3 d. Evaporation of the solvent gave an oil, which was purified by preparative TLC [developing solvent, $CHCl_3$ -AcOEt-MeOH (10:1:1)] to afford the methyl ether (**6**) (36.4 mg, 47%), as an oil. High-resolution MS, Calcd for $C_{14}H_{19}NO_2$: 233.1414. Found: 233.1399. 1H -NMR δ : 2.48 (3H, s, NMe), 3.55 (2H, s, C_1 -H), 3.85 (3H, s, OMe), 4.58 (2H, dt, $J=5.7, 1.4$ Hz, $OCH_2CH=CH_2$), 6.10 (1H, ddt, $J=17, 10, 5.7$ Hz, $OCH_2CH=CH_2$), 6.53, 6.62 (each 1H, s, ArH). The methyl ether (**6**) (22 mg) was dissolved in a mixture of MeOH (2 ml) and water (0.4 ml), and 10% Pd-C (20 mg) and *p*-TsOH (20 mg) was added. The whole was refluxed with stirring for 24 h. After removal of the catalyst by filtration, the filtrate was alkalized with 10% aqueous Na_2CO_3 . The product was extracted with a mixture of $CHCl_3$ and iso-PrOH (4:1) and the extract was dried over K_2CO_3 . Evaporation of the solvent gave a brown oil (14 mg), which was purified by preparative TLC [developing solvent, $CHCl_3$ -AcOEt-MeOH (6:1:1)] to afford **4** (4 mg, 22%), mp 152–154°C (MeOH). 1H -NMR [$CDCl_3$ - CD_3OD (1:1)] δ : 2.59 (3H, s, NMe), 3.73 (2H, s, C_1 -H), 3.87 (3H, s, OMe), 6.56, 6.64 (each 1H, s, ArH). The product was identical with an authentic specimen as judged by mixed melting point determination and comparison of their spectral data.

(b) Methylation of the 6-isopropoxy-7-ol (**7**) (18 mg) with diazomethane (excess) in ether-MeOH for 3 d gave a crude product (18 mg), which was purified by preparative TLC [developing solvent $CHCl_3$ -AcOEt-MeOH (10:1:1)] to afford the methyl ether (**15**) (12.3 mg, 64%), as an oil. 1H -NMR δ : 1.35 (6H, d, $J=5.7$ Hz, $OCHMe_2$), 2.48 (3H, s, NMe), 3.54 (2H, s, C_1 -H), 3.84 (3H, s, OMe), 4.48 (1H, septet, $J=5.7$ Hz, $OCHMe_2$), 6.52, 6.64 (each 1H, s, ArH). The methyl ether (**15**) (11.2 mg) was dissolved in a mixture of AcOH (2 ml) and concentrated HBr (0.4 ml), and the whole was heated at 80°C with stirring for 3 h. The reaction mixture was made alkaline with 10% aqueous Na_2CO_3 , and the product was extracted with a mixed solvent [$CHCl_3$ -iso-PrOH (4:1)]. After drying over K_2CO_3 , the solvent was evaporated off under reduced pressure to give an oil (15 mg). Purification by preparative TLC [developing solvent $CHCl_3$ -AcOEt-MeOH (6:1:1)] afforded **4** (4.6 mg, 50%), which was identical with an authentic sample prepared by procedure (a).

1-Benzyl-6-hydroxy-7-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline (14)—The isopropoxy phenol (**13**) (60 mg) was dissolved in MeOH (8 ml), and diazomethane-ether solution (excess) was added to the solution. The whole was allowed to stand at room temperature for 1 d and the solvent was evaporated off under reduced pressure to give an oil (61 mg), which was purified by preparative TLC [developing solvent $CHCl_3$ -AcOEt-MeOH (15:1:1)] to afford 1-benzyl-6-isopropoxy-7-methoxy-2-methyl-1,2,3,4-tetrahydroisoquinoline (**16**) (51.7 mg, 83%), as an oil. 1H -NMR δ : 1.34, 1.36 (each 3H, d, $J=5.7$ Hz, $OCHMe_2$), 2.56 (3H, s, NMe), 3.50 (3H, s, OMe), 4.47 (1H, septet, $J=5.7$ Hz, $OCHMe_2$), 5.96, 6.59 (each 1H, s, ArH), 7.00–7.40 (5H, m, PhH). Concentrated HBr (2 ml) was added to a solution of the methyl ether (**16**) (32.4 mg) in AcOH (5 ml), and the whole was heated at 80°C with stirring for 2.5 h. The reaction mixture was basified with 10% aqueous Na_2CO_3 , and the product was extracted with $CHCl_3$. The organic layer was washed with brine and dried over Na_2SO_4 . Evaporation of the solvent under reduced pressure gave a yellow oil (28 mg), which was purified by preparative TLC [developing solvent, $CHCl_3$ -AcOEt-MeOH (10:1:1)] to afford **14** (15.5 mg, 55%), mp 160–162°C (lit.⁴⁾ 161–163°C). The product was identical with an authentic sample.⁴⁾ 1H -NMR δ : 2.60 (3H, s, NMe), 3.51 (3H, s, OMe), 5.83, 6.64 (each 1H, s, ArH), 7.02–7.40 (5H, m, PhH). IR ν_{max} cm^{-1} : 3545 (OH).

Acknowledgements We thank Dr. T. Moroe of Takasago Perfumery Co., Ltd. for providing the starting vanillin, Mr. M. Tanaka for his technical assistance, Sankyo Co., Ltd. for elemental analyses, and Miss N. Sawabe and Mrs. N. Yamatani of this Faculty for NMR and MS measurements.

References and Notes

- 1) Part XXXI: H. Hara, M. Murakata, O. Hoshino, B. Umezawa, and T. Inoue, *Chem. Pharm. Bull.*, **36**, 1869 (1988).
- 2) B. Umezawa, O. Hoshino, and Y. Yamanashi, *Chem. Pharm. Bull.*, **19**, 2147 (1971).
- 3) R. Boss and R. Scheffold, *Angew. Chem. Int. Ed. Engl.*, **15**, 558 (1976).
- 4) O. Hoshino, K. Ohyama, M. Taga, and B. Umezawa, *Chem. Pharm. Bull.*, **22**, 2587 (1974).
- 5) H. G. Theuns, H. B. M. Lenting, C. A. Selemink, H. Tanaka, M. Shibata, K. Ito, and R. J. J. Ch. Lousberg, *Heterocycles*, **22**, 1995 (1984).
- 6) B. Umezawa, O. Hoshino, Y. Terayama, K. Oyama, Y. Yamanashi, T. Inoue, and T. Toshioka, *Chem. Pharm. Bull.*, **19**, 2138 (1971).
- 7) These ratios indicate the existence of two diastereoisomers.