

ARAGUSPONGINES B, C, D, E, F, G, H, AND J, NEW VASODILATIVE BIS-1-OXAQUINOLIZIDINE ALKALOIDS FROM AN OKINAWAN MARINE SPONGE, XESTOSPONGIA SP.

Motomasa KOBAYASHI, Kazuyoshi KAWAZOE, and Isao KITAGAWA*

Faculty of Pharmaceutical Sciences, Osaka University, 1-6, Yamada-oka, Suita, Osaka 565, Japan

Nine new vasodilative alkaloids, araguspongines A, B (1), C (2), D (3), E (4), F (5), G (6), H (7), and J (8), were isolated from an Okinawan marine sponge, Xestospongia sp. On the basis of chemical and physicochemical evidence, the absolute stereostructures of araguspongines B, D, E, F, G, H, and J were determined respectively as 1, 3, 4, 5, 6, 7, 8, and the relative stereostructure of araguspongine C was determined as 2 having two 1-oxaquinolizidine moieties. Araguspongines B, D, and E each comprised a pair of the enantiomers, 1a and 1b, 3a and 3b, and 4a and 4b, respectively.

KEYWORDS marine sponge; Xestospongia sp.; alkaloid; araguspongine; 1-oxaquinolizidine; vasodilative activity; HPLC chiral analysis


In a continuing search for new bioactive substances from marine organisms,¹⁾ we have isolated nine new macrocyclic alkaloids, araguspongines A, B (1), C (2), D (3), E (4), F (5), G (6), H (7), and J (8) from an Okinawan marine sponge Xestospongia sp. This paper describes their structures.

An acetone extract of the titled fresh sponge (4 kg collected in July at Aragusuku-jima, Okinawa Prefecture) was partitioned into a water-AcOEt mixture and the water phase was further partitioned with 1-butanol. The 1-butanol solute (75 g) was dissolved in aq. (COOH)₂ solution (pH 3) and extracted with AcOEt. The aqueous phase was then treated with aq. NH₄OH (to pH 10) and extracted again with AcOEt to furnish an alkaloid fraction (28 g). The alkaloid fraction (4 g) was subjected to silica gel column chromatography (benzene-acetone-NH₄OH) and HPLC (μ -PORASIL, n-hexane-benzene-Et₂NH; COSMOSIL 5C₁₈, CHCl₃-MeOH-CH₃CN-H₂O-NH₄OH) to afford araguspongines A (30 mg)²⁾, B (1)(85 mg), C (2)(131 mg), D (3)(1 g), E (4)(1.3 g), F (5)(18 mg), G (6)(5 mg), H (7)(16 mg), and J (8)(22 mg).

Araguspongine D (3)³⁾, C₂₈H₅₀N₂O₂,⁴⁾ [α]_D -5.1° (CHCl₃), showed 14 carbon signals in the ¹³C NMR spectrum, suggesting that 3 had a C₂ symmetry. The IR spectrum of 3 showed Bohlmann absorptions⁵⁾ (2810, 2760 cm⁻¹) due to the trans-quinolizidine ring. Based on ¹H (500 MHz) and ¹³C (125 MHz) NMR analysis, 3 was concluded to be identical with xestospongine A⁶⁾ which was previously isolated from the Australian marine sponge Xestospongia exigua and its relative configuration reported. However, the [α]_D value of 3 significantly differed from that ([α]_D +10°) of xestospongine A. So that araguspongine D (3) was subjected to HPLC separation using a chiral column [CHIRALCEL OF (DAICEL), n-hexane-2-ProH-Et₂NH] to provide (+)-araguspongine D (3b, [α]_D +10°) and (-)-araguspongine D (3a, [α]_D -9°) in 3:7 ratio. The 3b thus obtained was identical with xestospongine A⁶⁾ in all respects.

Araguspongine J (8),⁷⁾ C₃₀H₅₄N₂O₂, [α]_D -23.4° (CHCl₃) showed Bohlmann absorptions (2800, 2750 cm⁻¹) in the IR spectrum and 15 carbon signals in the ¹³C NMR spectrum due to the C₂ symmetrical trans-1-oxaquinolizidine structure. The ¹H NMR spectrum (CDCl₃) of 8 showed a doublet methyl signal (δ 1.10, J=6.7 Hz) together with signals due to 10 α -H_{ax} (δ 3.03, d, J=8.9 Hz), 6 α -H_{ax} (δ 1.90, ddd, J=12.2, 12.2, 2.8 Hz), and 6 β -H_{eq} (δ 2.66, brd, J=ca 12.2 Hz) which were very similar to those observed in the ¹H NMR spectrum of 3. The location of the secondary methyl was determined at 3 β axial from the coupling pattern of the signals due to 2 α -H_{ax} (δ 3.46, ddd, J=10.4, 2.3, 2.3 Hz), 4 α -H_{ax} (δ 2.30, dd, J=11.0, 3.4 Hz), and 4 β -H_{eq} (δ 2.73, dd, J=11.0, 1.8 Hz). Consequently, the relative stereostructure of araguspongine J was revealed to be 8, which corresponded to a 3 β ,3' β -dimethyl analogue of araguspongine D (3).

Araguspongine G (6),⁸⁾ C₂₉H₅₂N₂O₂, [α]_D -16° (CHCl₃) showed ¹H and ¹³C signals ascribable to one half moiety of both araguspongines D (3) and J (8) in its ¹H and ¹³C NMR spectra. So the relative stereostructure of araguspongine G was clarified as 6 which had a hybrid structure of 3 and 8.



The ^1H NMR spectrum of araguspongine F (5), $\text{C}_{29}\text{H}_{52}\text{N}_2\text{O}_2$, $[\alpha]_D +8.6^\circ$ (CHCl_3) showed ^1H signals ascribable to the 3 α -methyl-trans-1-oxaquinolizidine structure (δ 0.76, 3H, d, $J=6.4$ Hz; 2.95, 1H, brt, $J=\text{ca } 9.5$ Hz, 2 α -H_{ax}; 1.80, 1H, dd, $J=11.0, 11.0$ Hz, 4 α -H_{ax}; 2.85, 1H, dd, $J=11.0, 4.0$ Hz, 4 β -H_{eq}; 3.02, 1H, d, $J=8.5$ Hz, 10 α -H_{ax}) together with those observed in the ^1H NMR spectrum of araguspongine D (3). Araguspongine H (7), $\text{C}_{30}\text{H}_{54}\text{N}_2\text{O}_2$, $[\alpha]_D -9^\circ$ (CHCl_3) showed ^1H and ^{13}C signals ascribable to one each of the 3 β -methyl-trans-1-oxaquinolizidine and 3 α -methyl-trans-1-oxaquinolizidine moieties in its ^1H and ^{13}C NMR spectra. Therefore, the relative stereostructures of araguspongines F and H were 5 and 7, respectively.

Araguspongines F (5), G (6), H (7), and J (8) were shown to be optically pure by HPLC analysis using a chiral column. Based on the ^1H and ^{13}C NMR analysis, it was concluded that 5, 6, 7, and 8 have the same skeletal conformation as araguspongine D (3) regardless of the presence of 3 α - and/or 3 β -methyl group(s) which may contribute independently to the optical rotatory property. We then applied the Hudson rule¹¹⁾ to 5, 6, 7, 8 and araguspongine D (3a or 3b) and the results were as shown in the Chart. Thus, the comparison of $[M]_D$ values among 5, 6, 7, 8, and (-)-araguspongine D (3a) disclosed that 5, 6, 7, 8, and 3a have the same absolute stereostructures. To determine the absolute stereostructures of these alkaloids, we next applied Horeau's method¹²⁾ to a diol 9, which was obtained by NaBH_3CN reduction of araguspongine J (8) in THF-MeOH (1:1) under reflux. The recovered α -phenylbutyric acid showed $[\alpha]_D +1.6^\circ$ (benzene), so that the 2R configuration in 8 and consequently the absolute stereostructures of (-)- and (+)-araguspongines D (3a and 3b), (+)-araguspongine F (5), (-)-araguspongine G (6), (+)-araguspongine H (7), and (-)-araguspongine J (8) were determined as shown. In addition, the absolute stereostructure of xestospongine A⁶⁾, which was identified with (+)-araguspongine D (3b), was clarified.

Araguspongine B (1), $\text{C}_{28}\text{H}_{50}\text{N}_2\text{O}_2$, $[\alpha]_D -0.8^\circ$ (CHCl_3) showed 14 carbon signals in the ^{13}C NMR spectrum but lacked Bohlmann absorption in its IR spectrum. Treatment with Al_2O_3 60 F₂₅₄ (Type E, Merck) (80°C, 5h) selectively isomerized 1 to furnish 3, so that it was concluded that 1 had a bis-cis-1-oxaquinolizidine structure while possessing the same relative configurations at 2,2' and 9,9' as those in 3. Upon irradiation of 4 α -H_{ax} in the ^1H NMR of 1, 7% NOE was observed on 10 α -H whereas irradiation of 10 α -H resulted in 8% NOE on 2 α -H_{ax}. So the relative stereostructure of araguspongine B was shown to be 1 having a bis-cis-1-oxaquinolizidine structure.

Araguspongine E (4),¹⁴ $C_{28}H_{50}N_2O_2$, $[\alpha]_D -1.1^\circ$ ($CHCl_3$) showed 1H and ^{13}C signals ascribable to one half moiety of both araguspongines B (1) and D (3) in the 1H and ^{13}C NMR spectra and was selectively converted to 3 on Al_2O_3 treatment. So that the relative stereostructure of araguspongine E was shown to be 4 which had a hybrid structure of 1 and 3. HPLC analysis using a chiral column showed that both 1 and 4 were mixtures of the respective enantiomers 1a and 1b (1:1) and 4a and 4b (3:2), respectively: $[\alpha]_D$ ($CHCl_3$) $+17^\circ$ (1a), -15° (1b), $+1^\circ$ (4a), and -2° (4b). Al_2O_3 treatment of 1a and 4a furnished 3a, whereas the similar treatment of 1b and 4b furnished 3b. Thus, the absolute stereostructures of 1a, 1b, 4a, and 4b were elucidated as shown.

Finally, araguspongine C (2),¹⁵ $C_{28}H_{50}N_2O_4$, $[\alpha]_D +11.1^\circ$ ($CHCl_3$) showed 14 carbon signals in the ^{13}C NMR spectrum, suggesting 2 to have a C_2 symmetry. The 1H NMR analysis in detail¹⁶ of 2 revealed the presence of a 9-hydroxy-cis-1-oxaquinolizidine structure in 2. Furthermore, the configuration of the 9-hydroxyl moiety in 2 was elucidated by a pyridine-induced shift study¹⁷ of the 1H NMR of 2. Thus, the $7\alpha-H_{ax}$ and $10\alpha-H$ signals were significantly shifted lower [$\Delta\delta = \delta_{C_5D_5N} - \delta_{CDCl_3} = +0.35$ and $+0.24$, respectively] due to the presence of the $9\alpha-OH$ group. Consequently, the relative stereostructure of araguspongine C was clarified as 2.

Araguspongines C (2), D (3), E (4), and J (8) showed stronger vasodilative activities than papaverine in a perfusion model experiment using an isolated mesenteric artery of the SD-rat.

ACKNOWLEDGMENT The authors are grateful to Prof. P. R. Bergquist, the University of Auckland, New Zealand for identification of the marine sponge and to Dr. M. Endo, Suntory Institute for Biomedical Research for identification of araguspongine D and undertaking the bioassay.

REFERENCES AND NOTES

- 1) The recent paper: a) I. Kitagawa, M. Kobayashi, B. W. Son, S. Suzuki, and Y. Kyogoku, *Chem. Pharm. Bull.*, **37**, 1230 (1989); b) I. Kitagawa, *Yakugaku Zasshi*, **108**, 398 (1988).
- 2) Based on 1H and ^{13}C NMR analyses, araguspongine A is presumed to have a hybrid structure of araguspongine C (2) and D (3).
- 3) 3: $\delta(CDCl_3)$: 3.35 (brd, $J=ca$ 10.7 Hz, 2-H), 3.06 (d, $J=8.2$ Hz, 10-H), 2.94 (ddd, $J=11.8, 4.0, 1.8$ Hz, 4- H_{eq}), 2.76 (brd, $J=ca$ 11.6 Hz, 6- H_{eq}), 2.18 (ddd, $J=12.0, 11.8, 3.1$ Hz, 4- H_{ax}), 1.96 (ddd, $J=11.6, 11.6, 2.7$ Hz, 6- H_{ax}); $\delta_c(CDCl_3)$: 95.9 (C-10), 75.3 (C-2), 54.3, 54.1 (C-6,4), 40.6 (C-9).
- 4) The molecular composition of compounds with the chemical formulae were determined by high resolution mass spectrometry. 5) F. Bohlmann, *Angew. Chem.*, **69**, 641 (1957).
- 6) M. Nakagawa, M. Endo, N. Tanaka, and G. Lee, *Tetrahedron Lett.*, **25**, 3227 (1984).
- 7) 8: $\delta_c(CDCl_3)$: 96.9 (C-10), 78.3 (C-2), 62.3 (C-4), 54.4 (C-6), 40.1 (C-9), 33.4 (C-3), 13.3 (3-Me).
- 8) 6: $\delta(CDCl_3)$: 3.46 (ddd, $J=11.0, 2.5, 2.1$ Hz, 2-H), 3.35 (brt, $J=ca$ 10.7 Hz, 2'-H), 3.08 (d, $J=8.5$ Hz, 10'-H), 3.01 (d, $J=8.2$ Hz, 10-H), 2.94 (ddd, $J=11.8, 4.3, 2.1$ Hz, 4'- H_{eq}), 2.73 (dd, $J=11.0, 1.8$ Hz, 4- H_{eq}), 2.30 (dd, $J=11.0, 3.4$ Hz, 4- H_{ax}), 2.19 (ddd, $J=12.2, 11.8, 3.4$ Hz, 4'- H_{ax}), 1.10 (d, $J=7.0$ Hz, 3-Me); $\delta_c(CDCl_3)$: 97.3 (C-10), 95.9 (C-10'), 78.3 (C-2), 75.7 (C-2'), 62.3 (C-4), 54.4, 54.2 (C-6,6',4'), 40.8, 40.2 (C-9,9'), 33.4 (C-3), 13.3 (3-Me).
- 9) 5: $\delta(CDCl_3)$: 3.35 (brt, $J=ca$ 10.7 Hz, 2'-H), 3.06 (d, $J=8.6$ Hz, 10'-H), 2.94 (brd, $J=ca$ 11.7 Hz, 4'- H_{eq}), 2.18 (ddd, $J=11.7, 11.7, 3.4$ Hz, 4'- H_{ax}); $\delta_c(CDCl_3)$: 95.9 (C-10,10'), 81.1 (C-2), 75.1 (C-2'), 62.2 (C-4), 54.3, 54.1 (C-6,6',4'), 40.6, 40.5 (C-9,9'), 35.4 (C-3), 14.9 (3-Me).
- 10) 7: $\delta(CDCl_3)$: 3.40 (brd, $J=ca$ 10.4 Hz, 2-H), 2.96 (d, $J=8.8$ Hz, 10-H), 2.95 (d, $J=8.5$ Hz, 10'-H), 2.90 (ddd, $J=10.1, 10.1, 2.4$ Hz, 2'-H), 2.83 (dd, $J=11.0, 4.0$ Hz, 4'- H_{eq}), 2.71 (dd, $J=11.3, 1.8$ Hz, 4- H_{eq}), 2.30 (dd, $J=11.3, 3.4$ Hz, 4- H_{ax}), 1.80 (dd, $J=11.0, 11.0$ Hz, 4'- H_{ax}), 1.09 (d, $J=7.0$ Hz, 3-Me), 0.76 (d, $J=6.7$ Hz, 3'-Me); $\delta_c(CDCl_3)$: 97.9 (C-10), 96.9 (C-10'), 82.3 (C-2'), 79.1 (C-2), 62.1 (C-4,4'), 54.3, 53.7 (C-6,6'), 40.9, 40.4 (C-9,9'), 35.2 (C-3'), 33.1 (C-3), 15.0 (3'-Me), 13.3 (3-Me).
- 11) C. S. Hudson, *J. Am. Chem. Soc.*, **31**, 66 (1909). 12) A. Horeau, *Tetrahedron Lett.*, **15**, 506 (1961).
- 13) 1: $\delta(CDCl_3)$: 4.30 (d, $J=2.8$ Hz, 10-H), 3.53 (brt, $J=ca$ 11.0 Hz, 2-H), 3.18 (ddd, $J=13.7, 13.4, 3.1$ Hz, 4- H_{ax}), 2.95 (ddd, $J=13.7, 3.1, 1.5$ Hz, 4- H_{eq}); $\delta_c(CDCl_3)$: 87.5 (C-10), 76.0 (C-2), 52.8 (C-4), 45.3 (C-6), 40.5 (C-9).
- 14) 4: $\delta(CDCl_3)$: 4.24 (brs, 10-H), 3.48 (brt, $J=ca$ 10.7 Hz, 2-H), 3.30 (brt, $J=ca$ 10.7 Hz, 2'-H), 3.13 (brt, $J=ca$ 14.1 Hz, 4- H_{ax}), ca 3.00 (10'-H, 6- H_{ax}), 2.91 (brdd, $J=ca$ 14.1, 3.4 Hz, 4- H_{eq}), 2.88 (brd, $J=ca$ 12.2 Hz, 4'- H_{eq}), 2.12 (ddd, $J=12.2, 12.2, 3.4$ Hz, 4'- H_{ax}); $\delta_c(CDCl_3)$: 95.7 (C-10'), 87.4 (C-10), 75.7, 75.2 (C-2,2'), 54.2, 53.9 (C-6',4'), 52.6 (C-4), 45.3 (C-6), 40.4, 40.1 (C-9,9').
- 15) 2: $\delta(CDCl_3)$: 4.06 (s, 10-H), 3.56 (brt, $J=ca$ 10.7 Hz, 2-H), 3.11 (brt, $J=ca$ 13.7 Hz, 4- H_{ax}), 3.03 (ddd, $J=10.0, 10.0, 3.1$ Hz, 6- H_{ax}), 2.97 (brdd, $J=ca$ 13.7, 3.4 Hz, 4- H_{eq}), 2.34 (brd, $J=ca$ 10.0 Hz, 6- H_{eq}), 1.72 (m, 7- H_{ax}); (C_5D_5N): 2.07 (dddd, $J=13.0, 13.0, 13.0, 4.1, 4.1$ Hz, 7- H_{ax}); $\delta_c(CDCl_3)$: 90.4 (C-10), 76.5 (C-2), 70.8 (C-9), 52.6 (C-4), 44.3 (C-6).
- 16) For examples: irradiation of $10\alpha-H$ resulted in 9% NOE on $2\alpha-H_{ax}$ and that of $4\alpha-H_{ax}$ resulted in 5% NOE on $10\alpha-H$. Furthermore, NOE was also observed between $4\beta-H_{eq}$ and $6\alpha-H_{eq}$.
- 17) a) P. V. Demarco, E. Farkas, D. Doddrell, B. L. Mylari, and E. Wenkert, *J. Am. Chem. Soc.*, **90**, 5480 (1968); b) I. Kitagawa, M. Kobayashi, M. Hori, and Y. Kyogoku, *Chem. Pharm. Bull.*, **37**, 61 (1989).

(Received May 9, 1989)