

Preparation of Chiral, Highly Functionalized Cyclopentanes and Its Application to the Synthesis of Prostaglandin E₁


Kouji OKANO, Hiroshi SUEMUNE and Kiyoshi SAKAI*

Faculty of Pharmaceutical Sciences, Kyushu University, 3-1-1, Maidashi, Higashi-ku, Fukuoka 812, Japan. Received December 16, 1988


This paper describes the preparation of the polyfunctionalized cyclopentane ((±)-5) by silica gel-catalyzed air-oxidation, and its kinetic resolution by means of microbial reduction with *Rhodotorula rubra* to afford optically pure (–)-5 (>99% ee). Starting with (–)-5, a new route to prostaglandin E₁ was established.

Keywords prostaglandin E₁; air-oxidation; microbial reduction; chemoselective oxidation; stereoselective reduction

Previously, we reported¹⁾ that the highly functionalized cyclopentenone ((±)-2) was obtained by silica gel-catalyzed air-oxidation of the keto-diester ((±)-1) and that the kinetic resolution of (±)-2 by microbial reduction afforded optically pure (+)-2. Compound (+)-2 could be converted into (–)-α- and (+)-β-cuparenone *via* nickel-catalyzed stereoselective 1,4-addition of (*p*-tolyl)₂Zn as a key step (Chart 1). Silica gel-catalyzed air oxidation coupled with microbial reduction seems to provide optically pure, synthetically useful cyclopentanones. As a part of our synthetic studies on biologically active compounds based on a five-membered ring, we wish to report a further application of the above methodology to the synthesis of prostaglandin (PG) E₁.



In analogy with (±)-2, the cyclopentenone ((±)-4) obtained by base-catalyzed cyclization of the 1,4-diketone ((±)-3) underwent facile air-oxidation during silica-gel column chromatography with the AcOEt–hexane solvent system to afford two oxygenated products, (±)-5 (34%) and (±)-6 (15%) (Chart 2).



The structures of (±)-5 and (±)-6 were determined by means of spectral analysis. In the mass spectra (MS), 5 and 6 showed the molecular ion peak at *m/z* 356, and this finding suggests the addition of one oxygen atom to 4 (molecular weight, 340). The infrared (IR) spectra showed the absorption band of OH at 3450 cm^{–1}, in addition to the signals of hydroxy protons at δ4.05–4.09 in the proton

nuclear magnetic resonance (¹H-NMR) spectra. The position of the hydroxy function was deduced on the basis of the ¹H-NMR spectra. While the signals of the C₅-geminal protons in 5 were observed separately at δ2.70 and 2.89 (1H each), the C₅-geminal protons in 6 appeared at lower field (δ3.30 and 3.35) as signals with homoallylic coupling (*J* = 1.5 Hz) between C₅-H₂ and C₃-CH₂. From these results, it was concluded that the hydroxy function in 5 and 6 should be at C₄ and at C₂, respectively.

For the preparation of the chiral 5, enantioselective reduction of (±)-5 with microorganisms was tested. On screening of microbial reduction using 40 species of yeast, *Rhodotorula rubra* CCY 20-7-1 and *Schizosaccharomyces octosporus* were found to be effective for the kinetic resolution, and products of high optical purity were obtained. On reduction using 500 mg of the substrate (±)-5, *R. rubra* afforded (–)-5 (46%, >99% ee) as a recovered substrate, and (+)-7 (48%, >99% ee) as a reduced product after treatment with CH₂N₂²⁾ (Table I, entry 1), and *S. octosporus* gave (–)-5 (80%, 8% ee) and (+)-7 (14%, >99% ee) (Table I, entry 2). The optical purity of each product was determined from the 270 MHz ¹H-NMR spectral analysis by using a chiral shift reagent Tris[3-(heptafluoropropyl)-hydroxymethylene]-*d*-camphorato]europium(III) (Eu(hfc)₃) after conversion into (+)- or (–)-5.

According to the quadrant rule for *R. rubra* proposed by Nakazaki,³⁾ the recovered (–)-5 was concluded to be the *R*-isomer. The sequence for the synthesis of PGE₁ starts from (–)-5 with stereoselective catalytic hydrogenation (5% Pd/C in MeOH at room temperature) to give the saturated ketone ((–)-8, 75%), [*α*]_D²⁵ –71.1° (*c* = 1.89, CHCl₃). NaBH₄ reduction of (–)-8 at –20°C afforded the 1,4-*trans*-diol ((–)-9, 98%) as a single product, [*α*]_D²⁴ –10.8°

TABLE I.

Entry	Yeast	Yield (%)	[<i>α</i>] _D ^{a)}	% ee ^{b)}
1	<i>Rhodotorula rubra</i> CCY 20-7-1	(+)-7 48 (–)-5 46	+65.4° –119.6°	>99 >99
2	<i>Schizosaccharomyces octosporus</i>	(+)-7 14 (–)-5 80	+63.1° –9.3°	>99 8

a) In CHCl₃. b) Determined by 270 MHz ¹H-NMR analysis by using Eu(hfc)₃ (0.13 eq).




Chart 3




Chart 4




Chart 5

($c=2.54$, CHCl_3). These reductions proceeded stereoselectively, similarly to the case reported in the previous paper.¹⁾ The concurrent reduction of C₂- and C₄-esters in (-)-9 was achieved by using LiBH₄ in tetrahydrofuran (THF) at room temperature to afford the tetraol ((-)-10) in 84% yield, $[\alpha]_D^{27} -19.2^\circ$ ($c=1.84$, MeOH). The 1,2-diol in (-)-10 was protected as an acetonide by treatment with acetone/camphorsulfonic acid (CSA), and the acetonide (-)-11 was obtained in 89% yield, $[\alpha]_D^{26} -19.2^\circ$ ($c=1.57$, CHCl_3) (Chart 3).

To develop a method for introducing the ω -side chain into (-)-11, we preliminarily employed (±)-11. In the usual manner, the primary alcohol in (±)-11 was protected as the trityl ether by treatment with triphenylmethyl chloride (TrCl)/dimethylformamide (DMF) in the presence of 4-dimethylaminopyridine (DMAP), and the resulting 12 was converted to the pyranyl ether (13) (75% yield from 11 by treatment with 3,4-dihydro-2H-pyran (DHP)/pyridinium *p*-toluenesulfonate (PPTS) in CH_2Cl_2 . However, contrary to our expectation, the Tr-ether bond in (±)-13 was not cleaved by hydrogenolysis under reducing conditions using 10% Pd/C or PtO₂ (Chart 4). Next, we tried chemoselective oxidation of the primary alcohol of (±)-11 in the presence of the secondary alcohol with Pt black or $\text{RuCl}_2(\text{PPh}_3)_3$.⁴⁾ Oxidation of 11 with Pt black under an O₂ atmosphere for 12 h afforded the desired aldehyde-alcohol ((±)-15) in 49% yield and the diol 11 was recovered in 31% yield. Prolonged oxidation increased the production of the corresponding carboxylic acid, and the yield of 15 dropped. Ru complex-catalyzed oxidation of (-)-11 afforded 15 as a single product on thin layer chromatography (TLC).⁵⁾ The crude aldehyde 15 was subjected to Horner–Emmons reaction with dimethyl (2-oxoheptyl)phosphonate to afford the enone ((-)-16) (54%

from (-)-11, $[\alpha]_D^{26} -13.7^\circ$ ($c=0.35$, CHCl_3)). Reduction of (-)-16 by lithium thexylimonylborohydride in THF at -78°C gave (-)-17 (C_{15} - α -OH, $[\alpha]_D^{26} -13.2^\circ$ ($c=1.99$, CHCl_3)) (PG numbering) as the more polar,⁶⁾ major product in 64% yield, and (-)-18 (C_{15} - β -OH, $[\alpha]_D^{26} -23.1^\circ$ ($c=0.59$, CHCl_3)) as a less polar one in 28% yield. Even on reduction at -100°C , which required 6 h, the product ratio was almost same as at -78°C (17:18=52:25). The desired α -alcohol (-)-17 was treated with 80% aqueous acetic acid to afford the tetraol ((-)-19, 79%), $[\alpha]_D^{26} -17.0^\circ$ ($c=1.07$, MeOH). Oxidation of (-)-19 with NaIO₄ in aqueous acetone gave PGE₁ methyl ester ((-)-20) in 80% yield; this product was identical with an authentic sample in terms of the spectral data ($[\alpha]_D^{26} -53.3^\circ$ ($c=0.45$, MeOH), lit.⁷⁾ $[\alpha]_D^{23} -54.0^\circ$ ($c=1.08$, MeOH)), (Chart 5). Agreement of the sign of optical rotation supports the correctness of our estimation of the absolute configuration of (-)-5.

Experimental

IR spectra were measured on a JASCO A-202 spectrometer.¹ H-NMR spectra were measured on a JEOL JMN-PS-100 spectrometer unless otherwise stated. MS were taken on a JEOL JMS-D 300 spectrometer. Specific rotations were measured on JASCO DIP-4 and DIP-360 polarimeters. The melting points were measured with a Yanaco micro melting point apparatus and are uncorrected. For column chromatography, silica gel 70–230 mesh (Merck, Kieselgel 60) was used. TLC were performed on Silica gel 60 F₂₅₄ plates (0.25 and 2 mm) (Merck), respectively. All organic solvent extracts were washed with brine and dried over anhydrous sodium sulfate unless otherwise stated.

(±)-4-Hydroxy-2,4-bis(methoxycarbonyl)-3-(6-methoxycarbonylhexyl)-cyclopent-2-en-1-one (5) and (±)-2-Hydroxy-2,4-bis(methoxycarbonyl)-3-(6-methoxycarbonylhexyl)-cyclopent-3-en-1-one (6). K₂CO₃ (23.2 g, 167.6 mmol) was added portionwise to a stirred solution of trimethyl 2,5-dioxoundecane-1,4,11-tricarboxylate¹⁾ (20.0 g, 55.9 mmol) in MeOH (500 ml) at room temperature. After being stirred for 1 h, the reaction mixture was neutralized with 3% aqueous HCl under ice-water cooling, and the organic solvent was evaporated off *in vacuo*. The oily residue was

extracted with AcOEt (200 ml $\times$ 3), and the AcOEt extract was washed and dried. Removal of the solvent *in vacuo* gave an oily residue, which was subjected to column chromatography on silica gel (500 g), and eluted slowly with 15% AcOEt in hexane (v/v) to afford ($\pm$)-**6** (3.08 g, 15%) as a yellow oil. IR (neat): 3450, 1770, 1730—1720, 1635 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 1.1—1.9 (10H, m), 2.30 (2H, t, $J=7.2$ Hz, $\text{CH}_2\text{-COO}$), 3.30, 3.35 (1H each, t, $J=1.5$ Hz, $\text{C}_5\text{-H}$), 3.66, 3.80, 3.82 (3H each, s, COOCH_3), 4.09 (1H, br s, OH). FDMS m/z : 356 (M^+), 297. The fraction eluted with 20—40% AcOEt in hexane (v/v) afforded ($\pm$)-**5** as colorless needles, mp 63.5—64.5 $^\circ\text{C}$ (ether). IR (Nujol): 3430, 1750—1720, 1650 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.31 (2H, t, $J=7.3$ Hz, CH_2COO), 2.70, 2.89 (1H each, d, $J=18$ Hz, $\text{C}_5\text{-H}$), 3.67, 3.81, 3.87 (3H each, s, COOCH_3), 4.05 (1H, br s, OH). FDMS m/z : 357 ($\text{M}^+ + 1$), 297.

Microbial Reduction of ($\pm$)-5, (1S,4S)-1,4-Dihydroxy-2,4-bis(methoxycarbonyl)-3-(6-methoxycarbonylhexyl)cyclopent-2-ene ((+)-7 and (−)-5) In a manner similar to that described in the previous paper,¹⁾ ($\pm$)-**5** (500 mg) was subjected to microbial reduction with *Rhodotorula rubra* CCY 20-7-1 and with *Schizosaccharomyces octosporus*. The reaction mixture was shaken for 3 d at 30 $^\circ\text{C}$, then filtered with the aid of celite. In the case of *R. rubra*, hydrolysis of one of the esters was observed. The filtrate was acidified then extracted with AcOEt (200 ml $\times$ 3). Removal of the solvent *in vacuo* gave an oily residue, which was treated with CH_2N_2 . The crude products were subjected to column chromatography on silica gel (10 g). The fraction eluted with 20% AcOEt in hexane (v/v) afforded (−)-**5** (241 mg, 48%, >99% ee) as a colorless oil, $[\alpha]_{\text{D}}^{25} -119.6^\circ$ ($c=0.51$, CHCl_3). The fraction eluted with 30—40% AcOEt in hexane (v/v) afforded (+)-**7** (233 mg, 46%, >99% ee) as a colorless oil. IR (Nujol): 3470, 1735—1720, 1645, 1435, 1260, 1205, 755 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.1—2.8 (4H, m, CH_2COO , $\text{C}_5\text{-H}$), 3.66, 3.82, 3.88 (3H, each, s, COOCH_3), 3.5—3.9 (2H, br, OH $\times$ 2), 5.15 (1H, t, $J=4.8$ Hz, $\text{C}_1\text{-H}$). EIMS m/z : 357 ($\text{M}^+ - 1$), 340, 322, 308. $[\alpha]_{\text{D}}^{25} +65.4^\circ$ ($c=1.48$, CHCl_3). Reduction with *S. octosporus*; (−)-**5**: 401 mg, 80%, 8% ee, $[\alpha]_{\text{D}}^{27} -9.2^\circ$ ($c=1.54$, CHCl_3); (+)-**7**: 73 mg, 15%, >99% ee, $[\alpha]_{\text{D}}^{25} +63.1^\circ$ ($c=1.17$, CHCl_3).

(2R,3R,4R)-4-Hydroxy-3-(6-methoxycarbonylhexyl)-2,4-dimethoxycarbonylcyclopentanone ((−)-8) A solution of (−)-**5** (6.65 g, 18.7 mmol) in MeOH (600 ml) was hydrogenated in the presence of 5% Pd/C (6 g) under an H_2 atmosphere at room temperature. The catalyst was filtered off and the filtrate was concentrated *in vacuo* to afford an oily residue, which was purified by column chromatography on silica gel (60 g). The fraction eluted with 20% AcOEt in hexane (v/v) afforded (−)-**8** as a colorless oil. (Recemate: colorless needles, mp 100.5—101.5 $^\circ\text{C}$ (AcOEt-hexane)). IR (Nujol): 3450, 1765, 1745, 1730, 1125 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.30 (2H, t, $J=7.3$ Hz, CH_2COO), 2.78 (2H, s, $\text{C}_5\text{-H}$), 2.97 (1H, m, $\text{C}_3\text{-H}$), 3.30 (1H, d, $J=11.7$ Hz, $\text{C}_2\text{-H}$), 3.67, 3.78, 3.83 (3H, each, s, COOCH_3), 3.88 (1H, s, OH). EIMS m/z : 358 (M^+), 340, 326, 313, 308. $[\alpha]_{\text{D}}^{25} -71.1^\circ$ ($c=1.89$, CHCl_3).

(1R,2R,3R,4R)-2,4-Bis(methoxycarbonyl)-3-(6-methoxycarbonylhexyl)-cyclopentane-1,4-diol ((−)-9) A solution of (−)-**8** (4.00 g, 11.2 mmol) in MeOH (10 ml) was added dropwise to a stirred solution of NaBH_4 (423 mg, 11.2 mmol) in MeOH (50 ml) at -20°C . After 30 min, the reaction mixture was quenched with 3% aqueous HCl (10 ml) and concentrated *in vacuo*. The residue was extracted with AcOEt (50 ml $\times$ 4) and the combined extracts were dried. The solvent was removed *in vacuo* to give an oily residue, which was chromatographed on silica gel (40 g). The fraction eluted with AcOEt afforded (−)-**9** (4.00 g, 98%) as a colorless oil. IR (neat): 3470, 1740—1720 (br), 1435 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.66 (1H, dd, $J=11$, 6.6 Hz, $\text{C}_2\text{-H}$), 3.01 (2H, br s, OH $\times$ 2), 3.66, 3.74, 3.85 (3H each, s, COOCH_3), 4.52 (1H, dt, $J=8.0$, 6.6 Hz, $\text{C}_1\text{-H}$). EIMS m/z : 360 (M^+), 342, 328, 310, 301, 260, 251. $[\alpha]_{\text{D}}^{24} -10.8^\circ$ ($c=2.54$, CHCl_3).

(1R,2S,3R,4R)-2,4-Bis(hydroxymethyl)-3-(6-methoxycarbonylhexyl)-cyclopentane-1,4-diol ((−)-10) A solution of (−)-**9** (3.26 g, 9.05 mmol) in THF (10 ml) was added dropwise to a stirred suspension of LiBH_4 (296 mg, 13.6 mmol) in THF (40 ml) at room temperature. After 30 min, H_2O (5 ml) was added slowly and the solvent was removed *in vacuo*. The residue was chromatographed on silica gel (30 g). The fraction eluted with 10% MeOH in CHCl_3 (v/v) afforded (−)-**10** (2.75 g, 99%) as a colorless oil. IR (neat): 3370, 1735, 1435 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 0.8—1.3 (14H, m), 2.31 (2H, t, $J=7.0$ Hz, CH_2COO), 3.67 (3H, s, COOCH_3), 3.3—4.3 (8H, m, $\text{CH}_2\text{-O} \times 2$, OH $\times$ 4), 4.29 (1H, m, $\text{C}_1\text{-H}$). FDMS m/z : 343 ($\text{M}^+ + 1$), 342 (M^+), 327, 305. $[\alpha]_{\text{D}}^{25} -19.2^\circ$ ($c=1.84$, MeOH).

(5R,6R,7S,8R)-8-Hydroxy-7-hydroxymethyl-6-(6-methoxycarbonylhexyl)-2,2-dimethyl-1,3-dioxaspiro[4.4]nonane ((−)-11) A catalytic amount of CSA was added to a stirred solution of (−)-**10** (121 mg, 0.40 mmol) in acetone (2 ml) at room temperature. After 1 h, the solvent was removed *in vacuo* to leave an oily residue, which was subjected to column chromatog-

raphy on silica gel (4 g). The fraction eluted with 10% MeOH in CHCl_3 (v/v) afforded (−)-**11** (122 mg, 89%) as a colorless oil. IR (neat): 3420, 1735, 1435, 1370 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.31 (2H, t, $J=7.3$ Hz, CH_2COO), 2.96 (2H, br, OH $\times$ 2), 3.4—4.0 (2H, m, $\text{CH}_2\text{-OH}$), 3.67 (3H, s, COOCH_3), 3.83, 4.02 (1H each, d, $J=7.3$ Hz, OCCH_2O), 4.26 (1H, m, $\text{C}_1\text{-H}$). EIMS m/z : 344 (M^+), 329, 326, 311, 268. $[\alpha]_{\text{D}}^{26} -19.2^\circ$ ($c=1.57$, CHCl_3).

(5R*,6R*,7S*,8R*)-8-Hydroxy-6-(6-methoxycarbonylhexyl)-7-triphenylmethoxymethyl-2,2-dimethyl-1,3-dioxaspiro[4.4]nonane (12) A mixture of the diol ($\pm$)-**11** (47 mg, 0.14 mmol) and TrCl (42 mg, 0.15 mmol) in DMF (3 ml) was stirred at room temperature for 6 h in the presence of a catalytic amount of DMAP. The reaction mixture was diluted with brine (5 ml) and extracted with AcOEt (10 ml $\times$ 3). The AcOEt extract was washed and dried. Removal of the solvent *in vacuo* afforded an oily residue, which was subjected to column chromatography on silica gel (2 g). The fraction eluted with 15% AcOEt in hexane (v/v) afforded ($\pm$)-**12** (62 mg, 77%) as a colorless oil. IR (neat): 3460, 1735, 1590, 750, 700, 625 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 3.66 (3H, s, COOCH_3), 3.6—4.4 (3H, m, $\text{C}_8\text{-H}$, OCCH_2O), 7.1—7.6 (15H, m, Ar-H). FDMS m/z : 586 (M^+), 427, 243.

(5R*,6R*,7S*,8R*)-6-(6-Methoxycarbonylhexyl)-8-(tetrahydropyran-2-yl)oxy-7-triphenylmethoxymethyl-2,2-dimethyl-1,3-dioxaspiro[4.4]nonane ((±)-13) A mixture of ($\pm$)-**12** (50 mg, 0.09 mmol) and DHP (9 mg, 0.10 mmol) in CH_2Cl_2 (2 ml) at room temperature was stirred for 2 h in the presence of a catalytic amount of PPTS. The reaction mixture was diluted with brine (5 ml) and extracted with ether (10 ml $\times$ 3). The ether extract was washed and dried. Removal of the solvent *in vacuo* afforded an oily residue, which was subjected to column chromatography on silica gel (2 g). The fraction eluted 5% AcOEt in hexane (v/v) afforded ($\pm$)-**13** (56 mg, 98%) as a colorless oil. IR (neat): 1735, 1595, 1440, 1030, 760, 730, 700 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 2.28 (2H, t, $J=7$ Hz, $\text{CH}_2\text{-COO}$), 3.67 (3H, s, COOCH_3), 4.57, 4.86 (1H each, m, $\text{C}_8\text{-H}$, OCHO), 7.0—7.6 (15H, m, Ar-H). FDMS m/z : 670 (M^+).

(5R*,6R*,7R*,8R*)-7-Formyl-8-hydroxy-6-(6-methoxycarbonylhexyl)-2,2-dimethyl-1,3-dioxaspiro[4.4]nonane ((±)-15) A mixture of ($\pm$)-**11** (80 mg, 0.23 mmol) and Pt black (50 mg) in acetone (10 ml) was stirred for 12 h at room temperature under an O_2 atmosphere. The catalyst was filtered off, and the filtrate was concentrated *in vacuo* to afford an oily residue, which was subjected to column chromatography on silica gel (2 g). The fraction eluted with 50% AcOEt in hexane (v/v) afforded ($\pm$)-**15** (39 mg, 49%) as a colorless oil. IR (neat): 3460, 2720, 1735, 1715, 1440, 1370, 1210 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 1.1—2.5 (21H, m), 2.30 (2H, t, $J=6.7$ Hz, CH_2COO), 3.66 (3H, s, COOCH_3), 3.85, 4.06 (1H each, d, $J=9.0$ Hz, OCCH_2O), 4.57 (1H, m, $\text{C}_1\text{-H}$), 9.73 (1H, d, $J=2.2$ Hz, CHO). EIMS m/z : 342 (M^+), 327, 324, 309. The fraction eluted with AcOEt afforded the recovered diol ($\pm$)-**11** (25 mg, 31%).

9 α -Hydroxymethyl-9,9a-O-isopropylidene-15-keto-PGF_{1 β} Methyl Ester ((−)-16) A solution of the alcohol (−)-**11** (188 mg, 0.53 mmol) and $\text{RuCl}_2 \cdot (\text{PPh}_3)_3$ (393 mg, 0.41 mmol) in benzene (10 ml) was stirred for 5 h at room temperature. Removal of the solvent *in vacuo* gave a tarry residue, which was subjected to the next reaction without further purification. A solution of dimethyl (2-oxoheptyl)phosphonate (607 mg, 2.73 mmol) in ether (5 ml) was added to a stirred suspension of NaH (60% in oil; 109 mg, 2.73 mmol) in ether (5 ml) at 0 $^\circ\text{C}$. After 30 min, a solution of the above crude aldehyde **15** in CH_2Cl_2 (5 ml) was added and the whole was stirred for 5 h at room temperature. The reaction mixture was quenched with 3% aqueous HCl (5 ml) and extracted with AcOEt (20 ml $\times$ 3). The AcOEt extract was washed and dried. Removal of the solvent *in vacuo* afforded a tarry residue, which was subjected to column chromatography on silica gel (10 g). The fraction eluted with ether afforded (−)-**16** (130 mg, 54%) as a colorless oil. IR (neat): 3450, 1735, 1670, 1620, 1050 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 0.89 (3H, t, $J=5.9$ Hz, CH_3), 2.29 (2H, t, $J=7.6$ Hz, CH_2COO), 2.54 (2H, t, $J=7.6$ Hz, COCH_2), 3.66 (3H, s, COOCH_3), 3.6—4.3 (3H, m, $\text{C}_{11}\text{-H}$, OCCH_2O), 6.17 (1H, d, $J=15.7$ Hz, $=\text{CHCO}$), 6.77 (1H, dd, $J=15.7$, 8.5 Hz, $\text{CH}=\text{}$). FDMS m/z : 439 ($\text{M}^+ + 1$), 438 (M^+), 423, 393. $[\alpha]_{\text{D}}^{26} -13.7^\circ$ ($c=0.35$, CHCl_3).

9 α -Hydroxymethyl-9,9a-O-isopropylidene-PGF_{1 β} Methyl Ester ((−)-17) and Its C₁₅-Epimer ((−)-18) Lithium hexyllimonon borohydride (0.5 M in THF, 0.6 ml, 0.3 mmol) was added to a stirred solution of (−)-**16** (122 mg, 0.28 mmol) in ether (3 ml) at -78°C . The mixture was stirred for 2 h, H_2O (2 ml) was added, and the whole was extracted with AcOEt (5 ml $\times$ 4), then the AcOEt extract was washed and dried. Removal of the solvent *in vacuo* gave an oily residue, which was subjected to column chromatography on silica gel (3 g). The fraction eluted with 80% AcOEt in hexane (v/v) afforded a mixture of C₁₅-epimers (120 mg), which was subjected to

preparative TLC to afford a less polar β -alcohol (–)-**18** (34 mg, 28%) and a more polar α -alcohol (–)-**17** (80 mg, 66%), each as a colorless oil.

(–)-**18**: IR (neat): 0.88 (3H, t, $J=5.0$ Hz, CH_3), 1.0–2.4 (30H, m), 2.30 (2H, t, $J=7.6$ Hz, CH_2COO), 3.66 (3H, s, COOCH_3), 3.8–4.2 (2H, m, $\text{CH-O} \times 2$), 3.77, 4.03 (1H each, d, $J=8.8$ Hz, OCCH_2O), 5.5–5.8 (2H, m, vinyls). EIMS m/z : 422 ($\text{M}^+ - \text{H}_2\text{O}$), 404, 391, 378, 346. $[\alpha]_{\text{D}}^{26} -23.1^\circ$ ($c=0.59$, CHCl_3).

(–)-**17**: IR (neat): 0.89 (3H, t, $J=5.7$ Hz, CH_3), 0.9–2.4 (30H, m), 2.29 (2H, t, $J=7.4$ Hz, CH_2COO), 3.66 (3H, s, COOCH_3), 3.6–4.3 (4H, m, $\text{CH-O} \times 2$, OCCH_2O), 5.4–5.6 (2H, m, vinyls). EIMS m/z : 440 (M^+), 422, 404, 391, 378, 364. $[\alpha]_{\text{D}}^{26} -13.2^\circ$ ($c=1.99$, CHCl_3).

9 α -Hydroxymethyl-PGF_{1 β} Methyl Ester ((–)-19**)** The acetone (–)-**17** (18 mg, 0.04 mmol) in 80% AcOH (1 ml) was stirred at room temperature for 12 h. Removal of the solvent *in vacuo* afforded an oily residue, which was subjected to column chromatography on silica gel (2 g). The fraction eluted with 10% MeOH in CHCl_3 (v/v) afforded (–)-**19** (13 mg, 79%) as a colorless oil. IR (neat): 3370, 1740, 1630, 1560, 1440, 970 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 0.88 (3H, t, $J=4.6$ Hz, CH_3), 1.0–2.2 (24H, m), 2.29 (2H, t, $J=7.3$ Hz, CH_2COO), 3.66 (3H, s, COOCH_3), 3.8–4.2 (4H, m, $\text{CH-O} \times 2$, CH_2O), 5.4–5.7 (2H, m, vinyls). EIMS m/z : 382 ($\text{M}^+ - \text{H}_2\text{O}$), 364, 346, 328. $[\alpha]_{\text{D}}^{26} -17.0^\circ$ ($c=1.07$, MeOH).

PGE₁ Methyl Ester ((–)-20**)** A solution of NaIO_4 (11 mg, 0.05 mmol) in H_2O (1 ml) was added to a solution of (–)-**19** (10 mg, 0.025 mmol) in acetone (1 ml). The whole was stirred at room temperature for 1.5 h, and the resulting precipitate was filtered off. The filtrate was concentrated *in vacuo* to afford an oily residue, which was diluted with brine (2 ml) and extracted with AcOEt (5 ml $\times$ 3). The AcOEt extract was washed and

dried. The solvent was removed *in vacuo* to leave an oily residue, which was subjected to column chromatography on silica gel (2 g). The fraction eluted with 70% AcOEt in hexane (v/v) afforded (–)-**20** (7.4 mg, 80%) as a colorless oil. IR (neat): 3400, 1740, 1460, 1435, 1165, 1070, 965, 725 cm^{-1} . $^1\text{H-NMR}$ (CDCl_3) δ : 0.89 (3H, t, $J=5.4$ Hz, CH_3), 1.1–2.2 (18H, m, $\text{CH}_2 \times 9$), 1.8–2.6 (7H, m, CH_2COO , CHCO , $\text{OH} \times 2$), 2.73 (1H, dd, $J=18.2, 7.4$ Hz, $\text{C}_{10}\text{-H}$), 3.66 (3H, s, COOCH_3), 3.8–4.4 (2H, m, CH-O), 5.4–5.9 (2H, m, vinyls). EIMS m/z : 368 (M^+), 350, 332. $[\alpha]_{\text{D}}^{26} -53.3^\circ$ ($c=0.45$, MeOH), lit.⁷⁾ $[\alpha]_{\text{D}}^{23} -54.0^\circ$ ($c=1.08$, MeOH).

References and Notes

- 1) K. Okano, H. Suemune and K. Sakai, *Chem. Pharm. Bull.*, **36**, 1379 (1988), and references cited therein.
- 2) Hydrolysis of ester **5** was observed during the screening of yeasts (see Experimental).
- 3) M. Nakazaki, H. Chikamatsu, K. Naemura and M. Masao, *J. Org. Chem.*, **45**, 4432 (1980).
- 4) H. Tomioka, K. Takai, K. Oshima and H. Nozaki, *Tetrahedron Lett.*, **22**, 1065 (1981).
- 5) The aldehyde **15** obtained by oxidation with Ru complex was contaminated by a trace amount of by-product, but the low yield made it necessary to use the Ru complex.
- 6) In synthetic research on PGs, the polar fraction and the less polar fraction on TLC were tentatively assigned α -configuration and β -configuration, respectively.
- 7) M. Suzuki, T. Kawagishi, A. Yanagisawa, T. Suzuki, N. Okamura and R. Noyori, *Bull. Chem. Soc. Jpn.*, **61**, 1299 (1988).