

NOVEL AND CHIRAL HANTZSCH-TYPE 1,4-DIHYDROPYRIDINES HAVING A *p*-TOLYLSULFINYL GROUP. SYNTHESIS AND BIOLOGICAL ACTIVITIES AS CALCIUM CHANNEL ANTAGONISTS




Kazuyuki MIYASHITA, Masahiro NISHIMOTO, Tetsuya ISHINO, Satoshi OBIKA, and Takeshi IMANISHI*

Faculty of Pharmaceutical Sciences, Osaka University, Yamadaoka 1-6, Suita, Osaka 565, Japan

Both C-4 stereoisomers of novel Hantzsch-type 4-aryl- and 4-methyl-1,4-dihydropyridines **3** having a *p*-tolylsulfinyl group at C-5 were efficiently synthesized in optically pure forms starting from the α -sulfinyl enones **6** which could be easily obtained from (–)-menthyl (*S*)-*p*-tolylsulfinate (**4**). The stereochemistry at C-4 was found to be largely responsible for the biological activities as calcium channel antagonists of these compounds.

KEY WORDS Hantzsch-ester; 1,4-dihydropyridine; *p*-tolylsulfinyl group; synthesis; optically active form; calcium channel antagonist

The Hantzsch-esters, 1,4-dihydro-2,6-dimethyl-3,5-pyridinedicarboxylates **1**, are very attractive compounds from the standpoint of their outstanding biological activities. The 4-aryl-1,4-dihydropyridine derivatives are well known to be highly effective calcium channel antagonists.¹⁾ In addition, it was recently reported that the 4-methyl-1,4-dihydropyridines interfered with platelet aggregation, without any cardiovascular effects.²⁾ In these 4-substituted-1,4-dihydropyridines **1**, the C-4 position becomes stereogenic when R¹ and R³ are different. Since configuration is known to be so closely related to biological activities,^{1b, 3)} it is of great value to synthesize each enantiomer in a stereoselective fashion.^{4,5)}



- 3Aa:** R¹ = Ph, R² = H
b: R¹ = *o*-Cl-Ph, R² = H
c: R¹ = Me, R² = H
3Ba: R¹ = H, R² = Ph
b: R¹ = H, R² = *o*-Cl-Ph
c: R¹ = H, R² = Me

In our previous study of the NADH model compound **2**, the extraordinarily high effectiveness of a chiral sulfinyl group for asymmetric reduction has become clear.⁶⁾ Now, by combination of compounds **1** and **2**, we have designed the novel classes of optically active Hantzsch-type compounds, 4-aryl- and 4-methyl-1,4-dihydropyridines **3**, which are very interesting from the viewpoint of both biological and chemical properties. In this paper, we wish to describe the versatile syntheses of **3**, and also report their activities as calcium channel antagonists.

The keto sulfoxide **5**,⁷⁾ easily obtained from (–)-menthyl (*S*)-*p*-tolylsulfinate (**4**),⁸⁾ was condensed with arylaldehydes under modified Knoevenagel conditions⁹⁾ to give the enones **6a,b** in good yields. Enone **6c** was also prepared *via* an alternative route. The reaction of **4** with 1-propenylmagnesium bromide in THF afforded **7** as a mixture of two isomers (*E*:*Z* = *ca.* 1:3) in 66% yield.¹⁰⁾ Lithiation of **7** and the subsequent treatment with an excess of acetaldehyde gave **8** as a 1:1 diastereoisomeric mixture in 74% yield.¹¹⁾ On oxidation with the Dess-Martin periodinane,¹²⁾ **8** produced the enone **6c** in an excellent yield.

The reaction of **6a,b** with methyl 3-aminocrotonate in the presence of a catalytic amount of magnesium perchlorate in 2,2,2-trifluoroethanol at room temperature (Route A) gave the corresponding 4-aryl-1,4-dihydropyridines **3Aa,b** both as single diastereomers.^{5,13)} A similar treatment of **6c** in methanol without the catalyst also gave a satisfactory result (Table 1).^{14, 15)}

On the other hand, **6a-c** were treated with methyl acetoacetate in the presence of sodium hydride to give the corresponding diketones **9a-c**, which were condensed with ammonium acetate in methanol to afford the desired products **3Aa-c** and **3Ba-c** (Route B).^{15,16)} It is quite noteworthy that the stereoselectivity of the products (**3Aa,b** / **3Ba,b**) in Route B is opposite to that in Route A. This phenomenon means that both diastereomers in the case of 4-aryl derivatives are obtainable selectively by use of Route A or Route B, although there has been no reasonable explanation for the reverse stereoselectivity.

* To whom correspondence should be addressed.

The stereochemistries of these products **3A** and **3B** were confirmed on the basis of X-ray crystallographic analysis and ^1H -NMR data.¹⁶⁾ As shown in Fig. 1, the X-ray structure of **3Bb** indicates that the lone pair on the sulfur atom of the sulfinyl group is proximal to the methyl group at C-6 as expected from consideration of the allylic 1,3-strain.¹⁸⁾ The differences in the chemical shifts of the hydrogens at C-4 between **3Aa,b,c** (4.86, 5.47 and 3.83 ppm) and **3Ba,b,c** (4.51, 4.92 and 3.37 ppm) are attributable to the deshielding effect of the neighboring oxygen atom of the sulfinyl group in **3A** and the shielding effect of the neighboring tolyl group in **3B**, respectively.




Table 1. Cyclization of **6** to 1,4-Dihydropyridines **3**

Compd.	Route A		Route B	
	Conditions	Yield (%)	Yield ^{a)} (%)	3A:3B
a: R = Ph	b)	55	1 : 0	43
b: R = <i>o</i> -Cl-Ph	b)	45	1 : 0	41
c: R = Me	c)	80	2.5 : 1	54

a) Overall yield from **6**. b) In the presence of 0.5eq of $\text{Mg}(\text{ClO}_4)_2$ in 2,2,2-trifluoroethanol at room temperature. c) In methanol at room temperature.

Table 2. Relative Vasorelaxant Activities of 4-Aryl-1,4-dihydropyridines

Compd.	Conc. (nM)	Relative activity (%)
Nifedipine	1	100
3Aa	10	10
3Ab	10	29
3Ba	1	24
3Ba	10	69
3Bb	1	50
3Bb	10	69

Relative vasorelaxant activities of thus obtained 1,4-dihydropyridines are summarized in Table 2, which shows that the **3B**-type stereoisomers are much more effective than the **3A**-type ones and compound **3Bb** has the highest activity (1/2 of nifedipine) among these compounds. These results indicate the potential effectiveness of this type of compound having a sulfinyl group as a calcium channel antagonist.

Application of these novel 1,4-dihydropyridines for asymmetric reductions is now in progress.

ACKNOWLEDGMENT We thank Professor O. Muraoka, Kinki University, for the X-ray analysis of **3Bb**. Dr. T. Tatsuoka, Suntory Biomedical Institute, is also acknowledged for the assay of the vasorelaxant activities.

REFERENCES AND NOTES

- 1) a) Bossert F., Meyer H., Wehinger E., *Angew. Chem. Int. Ed. Engl.*, **20**, 762 (1981); Janis R.A., Triggle D.J., *J. Med. Chem.*, **26**, 775 (1983); (b) Goldmann S., Stoltefuss J., *Angew. Chem. Int. Ed. Engl.*, **30**, 1559 (1991).
- 2) Ortega M.P., Garcia M.D.C., Gijon M.A., Casa-Juana M.F.D., Priego J.G., Crespo M.S., Sunkel C., *J. Pharmacol. Exp. Ther.*, **255**, 28 (1990); Sunkel C.E., Casa-Juana M.F.D., Cillero F.J., Priego J.G., Ortega M.P., *J. Med. Chem.*, **31**, 1886 (1988).
- 3) Kajino M., Wada Y., Nagai Y., Nagaoka A., Meguro K., *Chem. Pharm. Bull.*, **37**, 2225 (1989); Ashimori A., Uchida T., Ohtaki Y., Tanaka M., Ohe K., Fukaya C., Watanabe M., Kagitani M., Yokoyama K., *ibid.*, **39**, 108 (1991).
- 4) Meyers A.I., Oppenlaender T., *J. Am. Chem. Soc.*, **108**, 1989 (1986); Enders D., Müller S., Demir A.S., *Tetrahedron Lett.*, **29**, 6437 (1988); Ebike H., Maruyama K., Achiwa K., *Chem. Pharm. Bull.*, **40**, 1083 (1992); *idem*, *Tetrahedron Asymmetry*, **3**, 1153 (1992).
- 5) Davis R., Kern J.R., Kurz L.J., Pfister J.R., *J. Am. Chem. Soc.*, **110**, 7873 (1988).
- 6) Imanishi T., Hamano Y., Yoshikawa H., Iwata C., *J. Chem. Soc., Chem. Commun.*, **1988**, 473.
- 7) Kunieda N., Suzuki A., Kinoshita M., *Bull. Chem. Soc. Jpn.*, **54**, 1143 (1981).
- 8) Solladié G., Hutt J., Girardin A., *Synthesis*, **1987**, 173.
- 9) Tanikaga R., Konya N., Tamura T., Kaji A., *J. Chem. Soc., Perkin Trans I*, **1987**, 825.
- 10) Abbott D.J., Colonna S., Stirling C.J.M., *J. Chem. Soc., Perkin Trans I*, **1976**, 492.
- 11) Fawcett J., House S., Jenkins P.R., Lawrence N.J., Russell D.R., *J. Chem. Soc., Perkin Trans I*, **1993**, 67.
- 12) Dess D.B., Martin J.C., *J. Org. Chem.*, **48**, 4155 (1983).
- 13) Davis and his co-workers⁵⁾ previously reported a similar reaction in refluxing methanol, in which the predominant formation of a **3A**-type stereoisomer was observed. In the present reactions under the same conditions, however, **3Aa,b** were found to be obtained along with **3Ba,b** in rather low yields (**3Aa/3Ba**=1.5, 22% yield, **3Ab/3Bb**=1.5, 27% yield).
- 14) The reaction of **6c** with methyl 3-aminocrotonate in the presence of magnesium perchlorate in 2,2,2-trifluoroethanol gave only a complex mixture of products.
- 15) Optical purities of these compounds were determined on the basis of ¹H-NMR analysis by use of a chiral shift reagent, Eu(tfc)₃ or (*R*)-*N*-3,5-dinitrobenzoyl- α -phenylethylamine¹⁷⁾ and were almost 100%.
- 16) **3Aa**, mp 212-213°C (MeOH). [α]_D²² +140.2° (c=0.54, CHCl₃). ¹H-NMR (CDCl₃) δ : 2.19, 2.29, 2.44, 3.57 (each 3H, s), 4.86 (1H, s), 5.73 (1H, br s), 6.72-7.07 (9H, m). **3Ba**, mp 176-177°C (benzene-hexane). [α]_D²¹ +518.8° (c=0.63, CHCl₃). ¹H-NMR (CDCl₃) δ : 2.23, 2.30, 2.40, 3.47 (each 3H, s), 4.51 (1H, s), 6.56 (1H, br s), 7.10-7.43 (9H, m). **3Ab**, mp 199-200°C (MeOH). [α]_D²² +141.5° (c=0.51, CHCl₃). ¹H-NMR (CDCl₃) δ : 2.17, 2.31, 2.39, 3.53 (each 3H, s), 5.47 (1H, s), 6.68-7.17 (9H, m). **3Bb**, mp 139-141°C (ethyl acetate). [α]_D²² +464.5° (c=0.64, CHCl₃). ¹H-NMR (CDCl₃) δ : 2.18, 2.25, 2.42, 3.46 (each 3H, s), 4.92 (1H, s), 6.95-7.51 (9H, m). **3Ac**, mp 186-187°C (Et₂O-CHCl₃). [α]_D²³ +542.5° (c=1.03, MeOH). ¹H-NMR (CDCl₃) δ : 0.25 (3H, d, *J*=6 Hz), 2.25, 2.31, 2.39, 3.65 (each 3H, s), 3.83 (1H, q, *J*=6 Hz), 7.21 (1H, br s), 7.28, 7.51 (4H, AA'BB', *J*=8 Hz). **3Bc**, mp 173-174°C (ethyl acetate). [α]_D²³ +473.3° (c=1.07, MeOH). ¹H-NMR (CDCl₃) δ : 1.11 (3H, d, *J*=6 Hz), 2.25, 2.33, 2.38 (each 3H, s), 3.37 (1H, q, *J*=6 Hz), 3.53 (3H, s), 6.76 (1H, br s), 7.25, 7.43 (4H, AA'BB', *J*=8 Hz).
- 17) Deshmukh M., Duñach E., Juge S., Kagan H.B., *Tetrahedron Lett.*, **25**, 3467 (1984).
- 18) Hoffmann R.W., *Chem. Rev.*, **89**, 1841 (1989) and references cited therein.

(Received January 20, 1995; accepted February 20, 1995)