

Amino Acids and Peptides. XLVII. Facile Synthesis of Flavacol, Deoxymuta-aspergillilic Acid and Optically Active Deoxyaspergillilic Acid from Dipeptidyl Aldehydes^{1–3)}

Yoshio OKADA,* Hiroaki TAGUCHI, and Toshio YOKOI

Faculty of Pharmaceutical Sciences, Kobe Gakuin University, Nishi-ku, Kobe 651-21, Japan.

Received June 24, 1996; accepted July 18, 1996

2(1H)-Pyrazinone ring-containing natural products, flavacol, deoxymuta-aspergillilic acid and finally optically active deoxyaspergillilic acid, were easily synthesized by a newly developed procedure for preparation of 2(1H)-pyrazinone derivatives from dipeptidyl aldehyde hydrochlorides. The absolute configuration of natural deoxyaspergillilic acid was synthetically determined as S.

Key words 2(1H)-pyrazinone; optically active deoxyaspergillilic acid; total synthesis; dipeptidyl aldehyde

Various 3,6-disubstituted 2(1H)-pyrazinones, such as flavacol **1**, deoxymuta-aspergillilic acid **2** and deoxyaspergillilic acid **3** (Fig. 1) have been found in secondary metabolites of *Aspergillus*^{4,5)} and *Streptomyces*.^{6,7)} Because of their structural relationship to aspergillilic acid, which exhibits antibiotic and smooth muscle relaxation activity, they have been of interest to both organic chemists and biochemists.⁸⁾ Therefore, extensive efforts have been directed toward the development of synthetic methods for 2(1H)-pyrazinone derivatives. Various synthetic procedures have been developed, but some inherent problems remain to be resolved.^{9,10)}

Previously, **1** was synthesized through D,L-leucyl-D,L-leucyl anhydride according to the reported method¹⁰⁾ and **2**^{11,12)} was synthesized by condensation of α -oxiimino-isovaleraldehyde with α -amino- γ -methylvaleronitrile (D,L-leucyl nitrile), followed by reduction with sodium hydrosulfide.¹³⁾ Several attempts to synthesize **3** have been made.^{14–16)} However, the methods are complicated, and the product was racemic.

Here we present a facile synthesis of **1**, **2** and optically active **3** by a newly developed and convenient method and the determination of the absolute configuration of natural deoxyaspergillilic acid.

Previously, we reported a simple and convenient synthetic procedure for 2(1H)-pyrazinone derivatives from dipeptidyl chloromethyl ketones.^{17–19)} Various kinds of dipeptidyl chloromethyl ketones were prepared and converted in good yields to the corresponding 2(1H)-pyrazinone derivatives by brief refluxing in MeOH. The corresponding dipeptidyl methyl ketones were also easily converted to 2(1H)-pyrazinone derivatives through a different cyclization mechanism from that of the chloromethyl ketones.^{19,20)} The above novel method can afford 2(1H)-pyrazinone derivatives in which desired substituent(s) can be introduced at position(s) 3 and/or 6.




Fig. 1. Structures of Flavacol **1**, Deoxymuta-aspergillilic Acid **2** and Deoxyaspergillilic Acid **3**

* indicates an asymmetric carbon atom.

* To whom correspondence should be addressed.

However, position 5 of the 2(1H)-pyrazinone derivatives is substituted with a methyl group. We considered that dipeptidyl aldehydes might afford the 2(1H)-pyrazinone derivatives without a substituent at position 5 by a similar cyclization mechanism to that of the dipeptidyl methyl ketones. Therefore, we attempted to prepare the 2(1H)-pyrazinone derivatives from the dipeptidyl aldehydes. According to Chart 1, Boc-Leu-N,O-dimethylhydroxamate (DMA) was prepared from Boc-Leu-OH and DMA using the 1,3-dicyclohexylcarbodiimide (DCC)/DMAP procedure.²¹⁾ After removal of the Boc group, the resultant amine was coupled with Boc-Phe-OH by a mixed anhydride procedure to give Boc-Phe-Leu-DMA (**5**). This amide was treated with LiAlH₄ to afford Boc-Phe-Leu-H. After removal of the Boc group of this aldehyde with 5.0N HCl-dioxane, the resultant amine hydrochloride was cyclized to 3-benzyl-6-isobutyl-2(1H)-pyrazinone (**4**) by stirring at room temperature in various solvents (DMF, dioxane, MeCN and MeOH) for 13 h. Among them, MeCN was the best solvent. Although the yield was not high, we did not study side reactions. The results showed that, as expected, 2(1H)-pyrazinone derivatives could be easily obtained from the dipeptidyl aldehyde




Chart 1.

Reagents and conditions: (i) DCC, DMAP; (ii) HCl-dioxane; (iii) isobutyl chloroformate, Et₃N, Boc-AA-OH (AA: Phe, Leu); (iv) LiAlH₄, THF, -15°C; (v) MeCN, 13 h, r.t.

Table 1. mp, t_R , $[\alpha]_D$ and MS of Natural and Synthetic Deoxyaspergillic Acid

	Natural 3	3a	3b
mp (°C)	83	80–81	80–80.5
t_R (min) ^{a)}	15.46	15.46	15.46
$[\alpha]_D^{21b)}$	+11.8°	+11.3°	–11.2°
MS (M ⁺)	208	208	208

a) See Experimental for HPLC conditions. b) $c = 1.0$, CHCl₃.

hydes.

This method was applied to the synthesis of naturally occurring 2(1*H*)-pyrazinone derivatives, because our novel method can easily afford 2(1*H*)-pyrazinone derivatives under slightly acidic conditions. It was also considered that this method might be suitable for the synthesis of optically active **3**, because **3** possesses one asymmetric carbon atom (Fig. 1) and this position is easily racemized during heating with alkali.¹¹⁾

First of all, this method was applied to the synthesis of **1** and **2**. Boc-Leu-OH was coupled with H-Leu-DMA by a mixed anhydride procedure to afford Boc-Leu-Leu-DMA (**6**), which was converted to Boc-Leu-Leu-H by treatment with LiAlH₄. After removal of the Boc group, the resultant HCl·H-Leu-Leu-H in MeCN was stirred at room temperature for 13 h to afford **1**. The structure of this compound was confirmed by NMR and MS analysis and the melting point was identical with that of natural **1**. Similarly, Boc-Leu-OH was coupled with H-Val-DMA by a mixed anhydride method to give Boc-Leu-Val-DMA (**7**), which was treated with LiAlH₄ to give Boc-Leu-Val-H. After removal of the Boc group, the resultant HCl·H-Leu-Val-H in MeCN was stirred at room temperature for 13 h to afford **2**. The structure of this compound was confirmed by NMR and MS analysis and the melting point was identical with that of natural **2**.¹²⁾

Finally, this new method was applied to the synthesis of optically active **3**. According to Chart 1, Boc-Leu-Ile-DMA (**8**) and Boc-Leu-allo-Ile-DMA (**9**) were prepared. These amides were converted to **3a** and **3b** (*S* and *R* at the asymmetric center of the *sec*-butyl group, respectively). The synthetic **3a** and **3b** exhibited a single peak at the same retention time on HPLC analysis as that of natural **3**. Their physicochemical data are summarized in Table 1. From these results, the absolute configuration of the natural deoxyaspergillic acid could be confirmed as *S*. From the NMR data for **3a** and **3b**, we can deduce that deoxyaspergillic acid might exist as a pyrazinone, rather than pyrazinol, form.¹⁹⁾ Thus, we were able to develop a facile and racemization-free synthetic procedure for the 2(1*H*)-pyrazinone derivatives (**1**, **2** and **3**) from dipeptidyl aldehydes and have succeeded in the first total synthesis of optically active **3**, as well as determination of the absolute configuration of natural **3** as *S*. Our convenient synthetic procedure should contribute to studies on the structure-antibacterial activity relationship of these compounds.

Experimental

Melting points were determined on a Yanagimoto micro-melting point apparatus without correction. Optical rotations were measured with an

automatic polarimeter, model DIP-360 (Japan Spectroscopic Co.). ¹H- (400 MHz) and ¹³C- (100 MHz) NMR spectra were recorded on a Bruker AM400 spectrometer. Chemical shift values are expressed as ppm downfield from tetramethylsilane used as an internal standard (δ -value). The *J* values are given in Hz. The ¹³C signals were assigned with the aid of distortionless enhancement by polarization transfer (DEPT) experiments, and multiplicities are indicated by p (primary), s (secondary), t (tertiary) or q (quaternary). Mass spectra were measured with a Hitachi M-2000 mass spectrometer using the EI technique. For HPLC analysis, the following conditions were employed; solvent: eluent A, 0.05% TFA/H₂O; eluent B, 0.05% TFA/MeCN; 80% A for 5 min, to 20% A for 20 min, 20% A for 5 min, to initial conditions for 10 min; column: Waters Nova-Pak C18 (3.9 × 150 mm), flow rate: 1 ml/min; retention time is reported as t_R . On TLC (Kieselgel G, Merck), R_f^1 , R_f^2 and R_f^3 values refer to the systems of CHCl₃, AcOEt and EtOH (30:19:1), AcOEt and hexane (1:2), and CHCl₃, MeOH and AcOH (90:8:2), respectively.

Boc-Phe-Leu-DMA (5) A mixed anhydride [prepared from Boc-Phe-OH (2.9 g, 11.0 mmol), Et₃N (1.7 ml, 12.1 mmol) and isobutyl chloroformate (1.44 ml, 11 mmol) as usual] in THF (80 ml) was added to an ice-cold solution of H-Leu-DMA·HCl [prepared from Boc-Leu-DMA²¹⁾ (3.0 g, 11.0 mmol) and 5.0*N* HCl-dioxane (13.2 ml, 66.0 mmol) as usual] in DMF (80 ml) containing Et₃N (1.7 ml, 12.1 mmol). The reaction mixture was stirred at 0°C for 1 h and then at room temperature overnight. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and recrystallized from AcOEt, yield 3.6 g (77.6%), mp 141–145°C, $[\alpha]_D^{25} -20.1^\circ$ ($c = 1.0$, MeOH), R_f^2 0.22. Anal. Calcd for C₂₂H₃₅N₃O₅: C, 62.7; H, 8.37; N, 9.97. Found: C, 62.8; H, 8.43; N, 9.85.

3-Benzyl-6-isobutyl-2(1*H*)-pyrazinone (4) Lithium aluminium hydride (89.8 mg, 2.37 mmol) was added to a stirred solution of **5** (0.5 g, 1.18 mmol) in tetrahydrofuran (THF) (20 ml). Reduction was complete within 30 min. The product was hydrolyzed with a solution of potassium hydrogen sulfate in water. The resultant aldehyde was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration. The Boc group was removed by treatment with 5.0*N* HCl-dioxane (1.4 ml, 7.1 mmol) and the resultant amine was dissolved in MeCN (150 ml). This solution was stirred at room temperature for 13 h. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and purified by silica-gel column chromatography (eluent, CHCl₃:AcOEt:EtOH = 30:19:1), yield 59.8 mg (20.9%), mp 105–107°C, R_f^1 0.52. ¹H-NMR (CDCl₃, 400 MHz) δ : ppm 13.1 (1H, br, NH), 7.39–7.17 (6H, m, 3-CH₂Ph + 5-H), 4.09 (2H, s, 3-CH₂Ph), 2.37 (2H, d, *J* = 7.3 Hz, 6-CH₂CH(CH₃)₂), 2.06 (1H, m, *J* = 6.8 Hz, 6-CH₂CH(CH₃)₂), 0.98 (6H, d, *J* = 6.6 Hz, 6-CH₂CH(CH₃)₂). ¹³C-NMR (CDCl₃, 100 MHz) δ : ppm 158.2 (q, C-2), 155.5 (q, C-3), 138.5 (q, C-1'), 137.9 (q, C-6), 129.4 (t, C-2'), 128.3 (t, C-3', 5'), 126.4 (t, C-4'), 123.4 (t, C-5), 39.6 (s, 6-CH₂CH(CH₃)₂), 39.4 (s, 3-CH₂Ph), 28.1 (t, 6-CH₂CH(CH₃)₂), 22.2 (p, 6-CH₂CH(CH₃)₂). Anal. Calcd for C₁₅H₁₈N₂O·0.1H₂O: C, 73.8; H, 7.51; N, 11.5. Found: C, 73.7; H, 7.43; N, 11.4.

Boc-Leu-Leu-DMA (6) A mixed anhydride [prepared from Boc-Leu-OH (0.72 g, 3.13 mmol), Et₃N (0.48 ml, 3.45 mmol) and isobutyl chloroformate (0.41 ml, 3.13 mmol) as usual] in THF (60 ml) was added to an ice-cold solution of H-Leu-DMA·HCl [prepared from Boc-Leu-DMA²¹⁾ (0.86 g, 3.13 mmol) and 5.4*N* HCl-dioxane (3.5 ml, 18.8 mmol) as usual] in DMF (60 ml) containing Et₃N (0.48 ml, 3.45 mmol). The reaction mixture was stirred at 0°C for 1 h and then at room temperature overnight. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and recrystallized from EtOH, yield 0.84 g (69.3%), mp 173–174°C, $[\alpha]_D^{25} -56.8^\circ$ ($c = 1.0$, MeOH), R_f^1 0.56, R_f^2 0.29. Anal. Calcd for C₁₉H₃₇N₃O₅: C, 58.9; H, 9.62; N, 10.8. Found: C, 58.7; H, 9.79; N, 10.9.

3-Isobutyl-6-isobutyl-2(1H)-pyrazinone (Flavacol) (1) Lithium aluminum hydride (47.0 mg, 1.3 mmol) was added to a stirred solution of **6** (0.25 g, 0.65 mmol) in THF (15 ml). Reduction was complete within 30 min. The product was hydrolyzed with a solution of potassium hydrogen sulfate in water. The resultant aldehyde was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration. The Boc group was removed by treatment with 7.2 N HCl-dioxane (0.9 ml, 6.5 mmol) and the resultant amine was dissolved in MeCN (75 ml). This solution was stirred at room temperature for 13 h. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and purified by flash silica-gel column chromatography (eluent, CHCl₃:AcOEt:EtOH=30:19:1), yield 14.5 mg (10.7%), mp 145–146 °C (natural, 144 °C), *R*_f¹ 0.62, *R*_f² 0.46, *t*_R 16.06 min (natural, 16.06 min). ¹H-NMR (CDCl₃ 400 MHz) δ: ppm 12.9 (1H, brs, NH), 7.18 (1H, s, 5-H), 2.65 (2H, d, *J*=7.1 Hz, 3-CH₂-CH(CH₃)₂), 2.39 (2H, d, *J*=7.3 Hz, 6-CH₂-CH(CH₃)₂), 2.21 (1H, nonet, 3-CH₂-CH(CH₃)₂), 2.09 (1H, nonet, 6-CH₂-CH(CH₃)₂), 0.98 (6H, d, *J*=6.6 Hz, 3-CH₂-CH(CH₃)₂), 0.97 (6H, d, *J*=6.7 Hz, 6-CH₂-CH(CH₃)₂). ¹³C-NMR (CDCl₃ 100 MHz) δ: ppm 158.3 (q, C-2), 156.4 (q, C-3), 137.7 (q, C-6), 123.0 (t, C-5), 41.5 (s, 3-CH₂-CH(CH₃)₂), 39.5 (s, 6-CH₂-CH(CH₃)₂), 28.2 (t, 3-CH₂-CH(CH₃)₂), 27.0 (t, 6-CH₂-CH(CH₃)₂), 22.7 (p, 3-CH₂-CH(CH₃)₂), 22.2 (p, 6-CH₂-CH(CH₃)₂). MS *m/z*: 208 M⁺.

Boc-Leu-Val-DMA (7) A mixed anhydride [prepared from Boc-Leu-OH (0.89 g, 3.84 mmol), Et₃N (0.59 ml, 4.23 mmol) and isobutyl chloroformate (0.5 ml, 4.23 mmol) as usual] in THF (60 ml) was added to an ice-cold solution of H-Val-DMA·HCl [prepared from Boc-Val-DMA²¹] (1.0 g, 3.84 mmol) and 5.4 N HCl-dioxane (4.27 ml, 23.1 mmol) as usual in DMF (60 ml) containing Et₃N (0.59 ml, 4.23 mmol). The reaction mixture was stirred at 0 °C for 1 h and then at room temperature overnight. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and recrystallized from EtOH, yield 0.81 g (57.1%), mp 155–157 °C, [α]_D²⁵ –65.6° (*c*=1.0, MeOH), *R*_f¹ 0.77, *R*_f³ 0.3. Anal. Calcd for C₁₈H₂₅N₃O₅: C, 57.9; H, 9.45; N, 11.3. Found: C, 57.6; H, 9.67; N, 11.2.

3-Isobutyl-6-isopropyl-2(1H)-pyrazinone (Deoxymuta-aspergillilic Acid) (2) Lithium aluminum hydride (50.8 mg, 1.34 mmol) was added to a stirred solution of Boc-Leu-Val-DMA (7) (0.25 g, 0.67 mmol) in THF (10 ml). Reduction was complete within 30 min. The product was hydrolyzed with a solution of potassium hydrogen sulfate in water, and the resultant aldehyde was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration. The Boc group was removed by treatment with 7.2 N HCl-dioxane (0.93 ml, 6.7 mmol) and the resultant amine was dissolved in MeCN (75 ml). This solution was stirred at room temperature for 13 h. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and recrystallized from acetone, yield 16.0 mg (11.8%), mp 104–105 °C (lit.⁹) 111 °C, *R*_f² 0.44, *t*_R 11.11 min. ¹H-NMR (CDCl₃ 400 MHz) δ: ppm 13.0 (1H, brs, NH), 7.22 (1H, s, 5-H), 2.85 (1H, m, 6-CH(CH₃)₂), 2.66 (2H, d, *J*=7.1 Hz, 3-CH₂-CH(CH₃)₂), 2.21 (1H, m, 3-CH₂-CH(CH₃)₂), 1.34 (6H, d, *J*=7.0 Hz, 6-CH(CH₃)₂), 0.97 (6H, d, *J*=6.7 Hz, 3-CH₂-CH(CH₃)₂). ¹³C-NMR (CDCl₃ 100 MHz) δ: ppm 158.4 (q, C-2), 157.1 (q, C-3), 143.6 (q, C-6), 120.2 (t, C-5), 41.5 (s, 3-CH₂-CH(CH₃)₂), 30.1 (t, 6-CH(CH₃)₂), 27.0 (t, 3-CH₂-CH(CH₃)₂), 22.7 (p, 3-CH₂-CH(CH₃)₂), 21.1 (p, 6-CH(CH₃)₂). MS *m/z*: 194 M⁺.

Boc-Leu-Ile-DMA (8) A mixed anhydride [prepared from Boc-Leu-OH (0.8 g, 3.4 mmol), Et₃N (0.52 ml, 3.7 mmol) and isobutyl chloroformate (0.45 ml, 3.7 mmol) as usual] in THF (50 ml) was added to an ice-cold solution of H-Ile-DMA·HCl [prepared from Boc-Ile-DMA²¹] (0.92 g, 3.4 mmol) and 6.5 N HCl-dioxane (2.6 ml, 17.0 mmol) as usual in DMF (50 ml) containing Et₃N (0.52 ml, 3.7 mmol). The

reaction mixture was stirred at 0 °C for 1 h and then at room temperature overnight. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and recrystallized from EtOH, yield 0.8 g (61.3%), mp 132–134 °C, [α]_D²⁵ –56.2° (*c*=1.0, MeOH), *R*_f³ 0.55. Anal. Calcd for C₁₉H₂₇N₃O₅: C, 58.9; H, 9.62; N, 10.8. Found: C, 58.9; H, 9.78; N, 10.9.

3-Isobutyl-6-(S)-sec-butyl-2(1H)-pyrazinone (3a) Lithium aluminum hydride (94.0 mg, 2.6 mmol) was added to a stirred solution of **8** (0.5 g, 1.3 mmol) in THF (30 ml). Reduction was complete within 30 min. The product was hydrolyzed with a solution of potassium hydrogen sulfate in water. Then, the resultant aldehyde was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration. The Boc group was removed by treatment with 5.4 N HCl-dioxane (2.4 ml, 13.0 mmol) and the resultant amine was dissolved in MeCN (150 ml). This solution was stirred at room temperature for 13 h. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and recrystallized from acetone, yield 72.2 mg (26.7%), mp 80–81 °C, [α]_D²⁵ +11.3° (*c*=1.0, CHCl₃), *R*_f² 0.5, *R*_f³ 0.63, *t*_R 15.46 min. ¹H-NMR (CDCl₃ 400 MHz) δ: ppm 12.8 (1H, brs, NH), 7.19 (1H, s, 5-H), 2.65 (2H, d, *J*=6.8 Hz, 3-CH₂-CH(CH₃)₂), 2.56 (1H, m, 6-CH(CH₃)CH₂CH₃), 2.21 (1H, m, 3-CH₂-CH(CH₃)₂), 1.82–1.6 (2H, m, 6-CH(CH₃)CH₂CH₃), 1.33 (3H, d, *J*=7.0 Hz, 6-CH(CH₃)CH₂CH₃), 0.97 (6H, d, *J*=6.7 Hz, 3-CH₂-CH(CH₃)₂), 0.91 (3H, d, *J*=7.4 Hz, 6-CH(CH₃)CH₂CH₃). ¹³C-NMR (CDCl₃ 100 MHz) δ: ppm 158.4 (q, C-2), 157.0 (q, C-3), 142.6 (q, C-6), 121.2 (t, C-5), 41.6 (s, 3-CH₂-CH(CH₃)₂), 37.3 (t, 6-CH(CH₃)CH₂CH₃), 28.5 (s, 6-CH(CH₃)CH₂-CH₃), 27.0 (t, 3-CH₂-CH(CH₃)₂), 22.7 (p, 3-CH₂-CH(CH₃)₂), 18.7 (p, 6-CH(CH₃)CH₂CH₃), 11.8 (p, 6-CH(CH₃)CH₂CH₃). MS *m/z*: 208 M⁺. Anal. Calcd for C₁₂H₂₀N₂O: C, 69.2; H, 9.68; N, 13.5. Found: C, 69.0; H, 9.88; N, 13.3.

Boc-Leu-allo-Ile-DMA (9) A mixed anhydride [prepared from Boc-Leu-OH (1.15 g, 5.0 mmol), Et₃N (0.84 ml, 6.0 mmol) and isobutyl chloroformate (0.66 ml, 5.0 mmol) as usual] in THF (80 ml) was added to an ice-cold solution of H-allo-Ile-DMA·HCl [prepared from Boc-allo-Ile-DMA (0.92 g, 3.4 mmol) and 5.4 N HCl-dioxane (5.6 ml, 30.0 mmol) as usual] in DMF (80 ml) containing Et₃N (0.84 ml, 6.0 mmol). The reaction mixture was stirred at 0 °C for 1 h and then at room temperature overnight. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and recrystallized from EtOH, yield 0.94 g (48.5%), mp 135–137 °C, [α]_D²⁵ –47.3° (*c*=1.0, MeOH), *R*_f¹ 0.78. Anal. Calcd for C₁₉H₂₇N₃O₅: C, 58.9; H, 9.62; N, 10.8. Found: C, 58.8; H, 9.89; N, 10.7.

3-Isobutyl-6-(R)-sec-butyl-2(1H)-pyrazinone (3b) Lithium aluminum hydride (47.0 mg, 1.3 mmol) was added to a stirred solution of **9** (0.25 g, 0.65 mmol) in THF (15 ml). Reduction was complete within 30 min. The product was hydrolyzed with a solution of potassium hydrogen sulfate in water. The resultant aldehyde was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration. The Boc group was removed by treatment with 5.4 N HCl-dioxane (1.2 ml, 6.5 mmol) and the resultant amine was dissolved in MeCN (150 ml). This solution was stirred at room temperature for 13 h. After removal of the solvent, the residue was extracted with AcOEt. The extract was washed with 10% citric acid, 5% NaHCO₃ and water and dried over Na₂SO₄, then the solvent was evaporated. Petroleum ether was added to the residue to afford crystals, which were collected by filtration and recrystallized from acetone, yield 55.7 mg (41.2%), mp 80–80.5 °C, [α]_D²⁵ –11.2° (*c*=1.0, CHCl₃), *R*_f² 0.5, *R*_f³ 0.63, *t*_R 15.46 min. ¹H-NMR (CDCl₃ 400 MHz) δ: ppm 12.8 (1H, brs, NH), 7.19 (1H, s, 5-H), 2.65 (2H, d, *J*=6.8 Hz, 3-CH₂-CH(CH₃)₂), 2.56 (1H, m, 6-CH(CH₃)CH₂-CH₃), 2.21 (1H, m, 3-CH₂-CH(CH₃)₂), 1.82–1.6 (2H, m, 6-CH(CH₃)CH₂CH₃), 1.33 (3H, d, *J*=7.0 Hz, 6-CH(CH₃)CH₂CH₃), 0.97 (6H, d, *J*=6.7 Hz, 3-CH₂-CH(CH₃)₂), 0.91 (3H, d, *J*=7.4 Hz,

6-CH(CH₃)CH₂CH₃). ¹³C-NMR (CDCl₃, 100 MHz) δ: ppm 158.3 (q, C-2), 157.1 (q, C-3), 142.6 (q, C-6), 121.2 (t, C-5), 41.6 (s, 3-CH₂CH(CH₃)₂), 37.3 (t, 6-CH(CH₃)CH₂CH₃), 28.4 (s, 6-CH(CH₃)-CH₂CH₃), 26.9 (t, 3-CH₂CH(CH₃)₂), 22.6 (p, 3-CH₂CH(CH₃)₂), 18.6 (p, 6-CH(CH₃)CH₂CH₃), 11.7 (p, 6-CH(CH₃)CH₂CH₃). MS *m/z*: 208 M⁺. *Anal.* Calcd for C₁₂H₂₀N₂O: C, 69.2; H, 9.68; N, 13.5. Found: C, 68.9; H, 9.97; N, 13.2.

Acknowledgements The authors are grateful to Dr. M. Sasaki of the Central Research Institute of Kikkoman Shoyu Co., Ltd., for providing natural flavacol and deoxyaspergillic acid. This work was supported in part by a grant from The Science Research Promotion Fund of the Japan Private School Promotion Foundation.

References and Notes

- 1) Part XLVI: Taguchi H., Yokoi T., Okada Y., *Chem. Pharm. Bull.*, **44**, 2037—2041 (1996).
- 2) The customary L indication for the amino acid residues is omitted. Standard abbreviations for amino acids, peptides and their derivatives are those recommended by the IUPAC-IUB Commission on Biochemical Nomenclature: *Biochemistry*, **5**, 2485—2489 (1966); *ibid.*, **6**, 362—364 (1967); *ibid.*, **11**, 1726—1732 (1972). Other abbreviations used are: Boc, *tert*-butoxycarbonyl; Et₃N, triethylamine; DMF, dimethylformamide; AcOEt, ethyl acetate; THF, tetrahydrofuran; Bzl, benzyl.
- 3) Preliminary communication, Okada Y., Taguchi H., Yokoi T., *Tetrahedron Lett.*, **37**, 2249—2252 (1996).
- 4) Yokotsuka T., Sasaki M., Kikuchi K., Asao Y., Nobuhara A., *Nippon Nogeikagaku Kaishi*, **41**, 32—38 (1967).
- 5) Sasaki M., Asao Y., Yokotsuka T., *Nippon Nogeikagaku Kaishi*, **42**, 288—293 (1968).
- 6) Tatsuka K., Tsuchiya T., Umeyama S., *J. Antibiot.*, **25**, 674—676 (1972).
- 7) Tatsuka K., Fujimoto K., Yamashita M., Tsuchiya T., Umeyama S., Umeyama H., *J. Antibiot.*, **26**, 606—608 (1973).
- 8) Dutcher J. D., *J. Biol. Chem.*, **171**, 321—339 (1947).
- 9) Jones R. G., *J. Amer. Chem. Soc.*, **71**, 78—81 (1949).
- 10) Baxter R. A., Spring F. S., *J. Chem. Soc.*, **1947**, 1179—1183.
- 11) MacDonald J. C., Bishop G. G., Mazurek M., *Tetrahedron*, **32**, 655—660 (1976).
- 12) Dunn G., Newbold G. T., Spring F. S., *J. Chem. Soc.*, **1949**, 2586—2589.
- 13) Nakamura S., *Agr. Biol. Chem.*, **25**, 658—664 (1961).
- 14) Newbold G. T., Sharp W., Spring F. S., *J. Chem. Soc.*, **1951**, 2679—2682.
- 15) Gallagher J. J., Newbold G. T., Sharp W., Spring F. S., *J. Chem. Soc.*, **1952**, 4870—4874.
- 16) Chigira Y., Masaki M., Ohta M., *Bull. Chem. Soc. Jpn.*, **39**, 1014—1020 (1966).
- 17) Taguchi H., Yokoi T., Kasuya F., Nishiyama Y., Fukui M., Okada Y., *J. Chem. Soc., Chem. Commun.*, **1994**, 247.
- 18) Okada Y., Taguchi H., Nishiyama Y., Yokoi T., *Tetrahedron Lett.*, **35**, 1231—1234 (1994).
- 19) Taguchi H., Yokoi T., Tsukatani M., Okada Y., *Tetrahedron*, **51**, 7361—7372 (1995).
- 20) Taguchi H., Yokoi T., Tsukatani M., Okamoto S., Wanaka K., Yasuda M., Horie N., Hijikata-Okunomiya A., Okada Y., "Peptide Chemistry," ed. by N. Nishi, 1995, Protein Research Foundation, Osaka, 1996, pp. 289—292.
- 21) Fehrentz J. A., Castro B., *Synthesis*, **1983**, 676—678.