

SYNTHESIS OF 1,4-DIKETONES BY MICHAEL ADDITION OF *O*-AROYLMANDELO-NITRILES INVOLVING REARRANGEMENT OF AROYL GROUP AND DECYANATION

Akira MIYASHITA,* Yoshiyuki MATSUOKA, Atsushi NUMATA, and Takeo HIGASHINO

School of Pharmaceutical Sciences, University of Shizuoka, 52-1 Yada, Shizuoka 422, Japan

The anions derived from *O*-aroylmandelonitriles **1** reacted with Michael addition acceptors such as acrylonitrile (**7**) and methyl acrylate (**10**) to give the corresponding 1,4-diketones **12**, **13**, and **15** in moderate to good yields. Under acidic conditions, the 1,4-diketones **12**, **13**, and **15** were converted into the furans **17**, **18** and **19** in good yields.

KEY WORDS *O*-aroylmandelonitrile; anion; Michael addition; 1,4-diketone; decyanation; furan

The anions derived from mandelonitriles behave as aroyl anions,¹⁾ and *O*-aroylmandelonitriles (**1**),^{1a)} *O*-trimethylsilylmandelonitriles (**2**),^{1a)} and *O*-(α -ethoxyethyl)mandelonitriles (**3**)^{1b)} can be used as aroyl anion equivalents. In connection with our studies on reactions promoted by the catalytic action of cyanide ion or the electron-accepting effect of the cyano group,²⁾ we therefore became interested in the mandelonitriles. Further, *O*-aroylmandelonitriles (**1**) are easily prepared.

The structure of the *O*-aroylmandelonitriles **1** is similar to that of Reissert compounds **4**. It was reported that the isoquinoline Reissert compound (**4a**) reacts with acrylonitrile (**7**) to give the ketone **5** in low yield together with the pyrrole derivative **6**.³⁾ The formation of **5** and **6** was explained in terms of Michael addition followed by rearrangement of the benzoyl group and decyanation. This result suggested that Michael addition of the anions derived from *O*-aroylmandelonitriles **1** with acrylonitrile (**7**) would give 1,4-diketones by rearrangement of the aroyl group and decyanation in the same way as Reissert compounds. 1,4-Diketones are starting compounds for preparation of five-membered heterocycles such as furans.









Chart 1

When *O*-benzoylmandelonitrile (**1a**) was treated with acrylonitrile (**7**) in the presence of NaNH₂ in DMF at room temperature for 2 h, only one product was formed.⁴⁾ It was identified as 2,3-dibenzoylpropionitrile (**12a**) from the spectral data and elemental analysis.⁵⁾ As assumed, the reaction appears to proceed through rearrangement of the benzoyl group and decyanation. Strong bases such as NaH and NaNH₂ are required to produce the anion from *O*-benzoylmandelonitrile (**1a**). DMF and DMSO were effective solvents, as shown in Chart 2.

Various *O*-aroylmandelonitriles **1a-f** were used in this Michael addition with acrylonitrile (**7**), and the expected 1,4-diketones **12a-f** were obtained in moderate to good yields. The reaction could be extended to methyl acrylate (**10**) and chalcone (**11**), and the corresponding 1,4-diketones **15a**, **15d**, and **16a** were formed in moderate yields. But, it failed to afford 1,4-diketones **14** from *O*-aroylmandelonitriles **1** and cinnamonnitrile (**9**), presumably because of steric hindrance. Attempts to synthesize 1,4-diketones **12g**, **12h**, **15g**, and **15h** by the use of *O*-benzoylmandelonitriles **1g** and **1h** having not only a strong electron-withdrawing substituent but also a strong electron-donating substituent were unsuccessful. Similarly, the 1,4-diketone **12i** with an aliphatic group could not be obtained by treatment of **1i** with **7**. However, the scope and limitations of the reaction remain to be fully established.

* To whom correspondence should be addressed.

$\text{Ar}^1-\text{CH}(\text{CN})-\text{COAr}^2 + \text{R}-\text{CH}=\text{CH}-\text{X} \xrightarrow[\text{r.t., 2 h}]{\text{base}} \text{Ar}^1-\text{C}(=\text{O})-\text{CH}(\text{R})-\text{CH}(\text{X})-\text{C}(=\text{O})-\text{Ar}^2$		1a-i		7-11	12-16					
Ar =										
		A	B	C	D	E	F			
Ar ¹	Ar ²	R	X	solvent	base ¹⁾	Yield (%) 1,4-diketone	(Recovery)			
1a	A	A	7	H	CN	DMSO	NaH	12a	62	(6)
1a	A	A	7	H	CN	DMF	NaNH ₂	12a	63	
1a	A	A	7	H	CN	DMF	DBU	12a	—	(55)
1a	A	A	7	H	CN	DMF	K ₂ CO ₃	12a	—	(72)
1a	A	A	7	H	CN	THF	NaNH ₂	12a	—	(85)
1a	A	A	8	Me	CN	DMF	NaH	13a	50	
1a	A	A	10	H	COOMe	DMF	NaNH ₂	15a	84	(10)
1a	A	A	11	Ph	COPh	DMSO	NaH	16a	48	
1b	A	B	7	H	CN	DMF	NaNH ₂	12b	41	(32)
1b	A	B	8	Me	CN	DMF	NaNH ₂	13b	63	(19)
1b	A	B	9	Ph	CN	DMF	NaNH ₂	14b	—	(98)
1c	A	C	7	H	CN	DMF	NaNH ₂	12c	30	
1d	B	A	7	H	CN	DMF	NaNH ₂	12d	55	(14)
1d	B	A	8	Me	CN	DMF	NaNH ₂	13d	42	(24)
1d	B	A	9	Ph	CN	DMF	NaNH ₂	14d	—	(95)
1d	B	A	10	H	COOMe	DMF	NaNH ₂	15d	74	(8)
1e	C	A	7	H	CN	DMF	NaNH ₂	12e	30	
1f	C	C	7	H	CN	DMF	NaNH ₂	12f	48	
1g	D	A	7	H	CN	DMSO	NaH	12g	—	(3) ²⁾
1g	D	A	10	H	COOMe	DMF	NaNH ₂	15g	—	(61)
1h	E	A	7	H	CN	DMF	NaNH ₂	12h	—	(—) ²⁾
1h	E	A	10	H	COOMe	DMF	NaNH ₂	15h	—	(—) ²⁾
1i	F	A	7	H	CN	DMF	NaNH ₂	12i	—	(28) ²⁾

1) DBU; 1,8-Diazabicyclo[5,4,0]-7-undecene. 2) Separation of the reaction products was not easy.

Chart 2

The proposed reaction pathway is illustrated in Chart 3. Michael addition proceeds between the anion derived from *O*-benzoylmandelonitrile (**1a**) and acrylonitrile (**7**) to give the intermediate (a). Then, rearrangement of the benzoyl group with decyanation results in the formation of the 1,4-diketone **12a**. The reaction process is similar to that of the Reissert compound **4a**. We considered that the rearrangement of the aroyl group proceeds through intramolecular reaction by formation of the intermediate (b) having a five-membered ring. To clarify the reaction pathway, cross-reaction was carried out. When a mixture of *O*-benzoylmandelonitrile (**1a**) and *O*-(*p*-toluoyl)-*p*-methylmandelonitrile (**1f**) was treated with acrylonitrile (**7**) in DMF, only 2,3-dibenzoylpropionitrile (**12a**) and 2,3-di(*p*-toluoyl)propionitrile (**12f**) were obtained. Formation of the cross-reaction products **12c** and **12e** could not be detected by HPLC analysis based on the retention times of authentic samples.⁶⁾ This result proves that the rearrangement of the aroyl group proceeds intramolecularly, as shown in Chart 3.

1,4-Diketones are useful compounds in organic synthesis. It has been reported that five-membered heterocycles such as furans can be synthesized by ring-closure of these compounds.⁷⁾ This led us to synthesize




Chart 3




Chart 4

several furans by use of the 1,4-diketones obtained in this study. In refluxing benzene under acidic conditions, 2,3-dibenzoylpropionitrile (**12a**) was converted into 2,5-diphenyl-3-furancarbonitrile (**17a**) in good yield.⁸⁾ The furans **17b**, **18b**, **18d**, and **19a** were obtained by similar treatments of the 1,4-diketones **12b**, **13b**, **13d** and **15a**. This is of interest, since a number of furans found in nature have biological activities.⁹⁾

REFERENCES AND NOTES

- 1) a) Deuchert K., Hertenstein U., Hunig S., Wehner G., *Chem. Ber.*, **112**, 2045-2061 (1979); b) Stork G., Maldonado L., *J. Am. Chem. Soc.*, **96**, 5272-5275 (1974); c) Jacobson R. M., Lahm G. P., Clader J. W., *J. Org. Chem.*, **45**, 395-405 (1980); d) Stork G., Maldonado L., *J. Am. Chem. Soc.*, **93**, 5286-5287 (1971); e) Hamana M., Endo T., Saeki S., *Tetrahedron Lett.*, **1975**, 903-906.
- 2) a) Miyashita A., Matsuoka Y., Iwamoto K., Higashino T., *Chem. Pharm. Bull.*, **42**, 1960-1962 (1994); b) Higashino T., Sato S., Miyashita A., Katori T., *Chem. Pharm. Bull.*, **34**, 4569-4576 (1986); c) *idem, ibid.*, **35**, 4078-4086 (1987); d) Higashino T., Goi M., Hayashi E., *ibid.*, **22**, 2493-2501 (1974).
- 3) a) Giridhar V., McEwen W. E., *J. Heterocycl. Chem.*, **8**, 121-123 (1971); b) Uff B. C., Budhram R. S., *Synthesis*, **1978**, 206; c) Uff B. C., Budhram R. S., Consterdine M. F., Hicks J. K., Slingsby B. P., Pemblington J. A., *J. Chem. Soc., Perkin 1*, **1977**, 2018-2022.
- 4) A typical procedure: Sodium amide (86 mg, 2.2 mmol) was slowly added to a mixture of *O*-benzoyl-mandelonitrile (**1a**, 474 mg, 2 mmol) and acrylonitrile (**7**, 212 mg, 4 mmol) in 10 ml of DMF, and the resulting solution was stirred at room temperature for 2 h. The reaction mixture was poured into 100 ml of H₂O, neutralized with AcOH, and extracted with AcOEt. The organic layer was washed with H₂O, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography on SiO₂ with benzene to give 1,3-dibenzoylpropionitrile (**12a**, 332 mg, 63%), colorless needles (hexane), mp 65-66°C.
- 5) **12a**: *Anal.* Calcd for C₁₇H₁₃NO₂: C, 77.55; H, 4.98; N, 5.32. Found: C, 77.68; H, 4.81; N, 5.31. IR (KBr) cm⁻¹: 2236 (CN), 1678 (CO). ¹H-NMR (CDCl₃) δ (ppm): 3.53 (1H, q, Ha, Jab = 19.8, Jac = 9.6), 4.07 (1H, q, Hb, Jba = 19.8, Jbc = 6.0), 5.00 (1H, q, Hc, Jcb = 6.0, Jac = 9.6), 7.13-7.63 (6H, m, aromatic H), 7.80-8.17 (4H, m, aromatic H).

$$\text{PhCO}-\text{C}(\text{H}_a)-\text{C}(\text{H}_b)(\text{CN})-\text{C}(\text{H}_c)-\text{COPh}$$
- 6) HPLC conditions: Column; Develosil Packed Column (DEVELOASIL C8-5), 4.6 × 250 mm. Flow; 1.0 ml/min. Solvent; acetonitrile / H₂O (7:3). Detector; UV 240 nm. The chromatogram of the cross-reaction mixture was compared with that of authentic samples based on the retention times. The retention times of the authentic samples were as follows; **12a** (8.04 min), **12c** (9.69 min), **12e** (9.71 min), and **12f** (11.93 min).
- 7) a) Jones R. G., Kornfeld E. C., *J. Org. Chem.*, **19**, 1671-1680 (1954); b) *idem*, *J. Am. Chem. Soc.*, **77**, 4069-4074 (1955).
- 8) A typical procedure: A mixture of 1,4-dibenzoylpropionitrile (**12a**, 418 mg, 1.56 mmol) and a catalytic amount of *p*-toluenesulfonic acid (50 mg, 0.29 mmol) in 30 ml of benzene was refluxed for 5 h. The reaction mixture was concentrated under reduced pressure and the residue was purified by column chromatography on SiO₂ with benzene. The first fraction gave 2,5-diphenyl-3-furancarbonitrile (**17a**, 298 mg, 78%), colorless needles (hexane), mp 117-118 °C. *Anal.* Calcd for C₁₇H₁₁NO: C, 83.25; H, 4.52; N, 5.71. Found: C, 83.25; H, 4.33; N, 5.65. IR (KBr) cm⁻¹: 2222 (CN). ¹H-NMR (CDCl₃) δ (ppm): 6.80 (1H, s, furan C⁴-H), 7.23-7.83 (8H, m, aromatic H), 7.84-8.20 (2H, m, aromatic H).
- 9) Dean F. M., Sargent M. V., "Comprehensive Heterocyclic Chemistry" Vol. 4, ed. by Katritzky A., Rees C. W., Pergamon Press, Oxford, 1984, pp. 531-712.

(Received September 25, 1995; accepted January 19, 1996)