

A NOVEL SYNTHESIS OF (+)-AZIMIC ACID

Toshiko KIGUCHI, Mitsuko SHIRAKAWA, Ichiya NINOMIYA, and Takeaki NAITO*


Kobe Pharmaceutical University, Motoyamakita, Higashinada, Kobe 658, Japan

A combination of 1,3-dipolar cycloaddition of *Z*-nitron (2) to the chiral dipolarophile (3) and subsequent ring transformation of the resulting adducts (4 and 6) to piperidinol (17) has provided a new practical synthesis of 2,3,6-trisubstituted piperidine alkaloid, (+)-azimic acid (1).

KEY WORDS azimine; azimic acid; 1,3-dipolar cycloaddition; nitron; enantioselective synthesis; isoxazolidine

Piperidine alkaloids are abundant in nature and many of them exhibit important biological activity. Azimine and carpaine are two of the macrocyclic dilactones¹⁾ consisting of two molecules of characteristic 2,3,6-trisubstituted piperidines with carboxyl and hydroxyl groups, designated as azimic and carpamic acids. In addition, their interesting biological activities have attracted the interest of chemists as their synthetic targets as seen from several total synthetic studies.²⁾


Asymmetric cycloadditions of nitrones have been successfully applied to the synthesis of natural products by many groups.³⁾ We now provide a potential method for synthesis of (+)-azimic acid (1) *via* the 1,3-dipolar cycloaddition of the nitron (2) to the chiral dipolarophile (3) and the subsequent ring transformation of the resulting adducts to the piperidinols and their deoxygenation reaction.




Our study⁴⁾ on the 1,3-dipolar cycloaddition of *C,N*-dialkyl nitron to the allyl alcohols established that the reaction rate and the stereoselectivities are influenced by the form of the hydroxyl group, of which the rate was accelerated and the *trans*-selectivity was improved when the magnesium alkoxide (OMgBr)⁵⁾ is used. For the synthesis of (+)-azimic acid (1), we first investigated the magnesium-induced cycloaddition of *Z*-nitron (2),⁶⁾ prepared from *N*-benzylhydroxylamine and methyl 7-oxoheptanoate,⁷⁾ to the allyl alcohol (3) which was prepared from methyl (-)-lactate according to the known procedure.⁸⁾ The allyl alcohol (3) was treated with methylmagnesium bromide and the resulting magnesium alkoxide with nitron (2) in methylene dichloride under reflux to give a 1:4 mixture of two adducts (4) and (5) in 17% combined yield with recovery of a large amount of 3. Thus, magnesium-induced cycloaddition of *Z*-nitron (2) took place with *trans*-stereoselectivity to give a *trans-threo* isomer (5), which was not suited for the synthesis of (+)-azimic acid (1). On the other hand, cycloaddition of *Z*-nitron (2) to the free allyl alcohol (3) in toluene under reflux proceeded smoothly to give a mixture of three adducts (4—6) in 83% combined yield which was readily separated by chromatography. Their stereochemistries were deduced from the structures of the

* To whom correspondence should be addressed.

corresponding ring-transformed piperidinols as shown below. Of three adducts (4—6), the *cis-threo* (4) and *trans-erythro* (6) were employed for the synthesis of (+)-azimic acid (1). The *threo*-selectivity observed in both cycloadditions to magnesium alkoxide and to free alcohol was in accordance with the known results.⁹⁾



Protection of the hydroxyl group in the adducts (4—6) with MEM chloride and deprotection of the silyl group with TBAF gave the alcohols (7—9) in 54—76% yields. Upon treatment with methanesulfonyl chloride-pyridine, the alcohols (7—9) gave the corresponding mesylates which were then treated with hydrogen in the presence of Pearlman's catalyst to yield a mixture of the ring-transformed piperidinols (13—15) and the acyclic mesylates (10—12). For the completion of the cyclization reaction of the acyclic mesylates, the crude hydrogenated products were treated with NaHCO₃ in MeOH-H₂O to give the piperidinols (13—15)¹⁰⁾ in 62—93% yields *via* 3 steps which were characterized with the ¹H-NMR spectra. Deoxygenation of the piperidinols (13) and (15) *via* Barton's ester¹¹⁾ gave the identical piperidine (16) in 93% yield from 13 and 81% yield from 15, respectively. Deprotection of the MEM group in 16 was achieved by treatment with TiCl₄ to give the piperidinol (17)¹⁰⁾ in 81% yield. The hydrolysis of the ester group in 17 with KOH-MeOH gave (+)-azimic acid (1), which was found to be identical with the authentic sample of (+)-1 upon comparisons of their spectral data^{2ab)} including optical rotation [colorless crystals (MeOH), m.p. 217°C (dec.), [α]_D +7.4° (c=0.52, MeOH), lit. m.p. 210-214°C (dec.),^{2c)} 214-215°C (EtOH),^{2b)} [α]_D +7.9° (c=1, MeOH)^{2b)}].



Reagent; (a) MEMCl, (*i*-Pr)₂NEt, CH₂Cl₂, Δ; (b) (*n*-Bu)₄NF, THF, Δ; (c) MsCl, Py., 0°C; (d) 20% Pd(OH)₂-C, MeOH, H₂; (e) NaHCO₃, H₂O-MeOH; (f) Im₂CS, THF; (g) Bu₃SnH, Toluene, Δ; (h) TiCl₄, CH₂Cl₂, 0°C; (i) MeOH, HCl

ACKNOWLEDGMENTS We wish to thank the Ministry of Education, Science and Culture (Japan) and the Science Research Promotion Fund of the Japan Private School Promotion Foundation for research grants. Thanks are also extended to Professor S. Hanessian, University of Montreal (Canada), for sending us the spectral data of (+)-azimic acid.

REFERENCES AND NOTES

- 1) Smalberger T. M., Rall G. J. H., De Waal H. L., *Tetrahedron*, **24**, 6417-6421 (1968).
- 2) a) Brown E., Dhal R., *J. Chem. Soc., Perkin Trans. 1*, **1976**, 2190-2193; b) Hanessian S., Frenette R., *Tetrahedron Lett.*, **1979**, 3391-3394; c) Lu Z. -H., Zhou W. -S., *Tetrahedron*, **49**, 4659-4664 (1993) and references therein.
- 3) Tufariello J. J., "1,3-Dipolar Cycloaddition Chemistry," Vol. 2, ed. by Padwa A., John Wiley & Sons, New York, 1988, pp. 83-168; Confalone P. N., Huie E. M., "Organic Reactions," Vol. 36, ed. by Kende A. S., John Wiley & Sons, New York, 1988, pp. 3-173; Breuer E., "Nitrones, Nitronates and Nitroxides," in *The Chemistry of Functional Groups*, eds. by Patai S., Rappoport Z., John Wiley & Sons, New York, 1989, pp. 139-312; Naito T., Ikai M., Shirakawa M., Fujimoto K., Ninomiya I., Kiguchi T., *J. Chem. Soc., Perkin Trans. 1*, **1994**, 773-775.
- 4) Naito T., Shirakawa M., Ikai M., Ninomiya I., Kiguchi T., *Recl. Trav. Chim. Pays-Bas*, **115**, 13 (1996).
- 5) Kanemasa S., Tsuruoka T., Wada E., *Tetrahedron Lett.*, **34**, 87-90 (1993); Kanemasa S., Tsuruoka T., *Chem. Lett.*, **1995**, 49-50.
- 6) The Z configuration of nitrone (**2**) was firmly deduced by comparison of ^1H -NMR spectrum with those of the related nitrones.⁴⁾ **2**: Colorless plates. m.p. 65-67°C (Et₂O). IR (CHCl₃) cm^{-1} : 1731 (CO₂), 1600 (C=N). ^1H -NMR (200 MHz, CDCl₃) δ : 6.63 (1H, t, $J=6$ Hz, HC=N⁺), 4.87 (2H, s, CH₂Ph). *Anal.* Calcd for C₁₆H₁₇NO: C, 68.41; H, 8.04; N, 5.32. Found: C, 68.15; H, 8.10; N, 5.16.
- 7) Ballini R., Marcantoni E., Petrini M., *Synth. Commun.*, **21**, 1075-1081 (1991).
- 8) Ley S. V., Armstrong A., Díez-Martín D., Ford M. J., Grice P., Knight J. G., Kolb H. C., Madin A., Marby C. A., Mukherjee S., Shaw A. N., Slawin A. M. Z., Vile S., White A. D., Williams D. J., Woods M., *J. Chem. Soc., Perkin Trans. 1*, **1991**, 667-692. The stereostructure of **3** was not determined in the literature but was characterized by the empirical rule on ^1H -NMR data reported: Iio H., Mizobuchi T., Tsukamoto M., Tokoroyama T., *Tetrahedron Lett.*, **27**, 6372-6376 (1986).
- 9) Kanemasa S., Nishiuchi M., Kamimura A., Hori K., *J. Am. Chem. Soc.*, **116**, 2324-2339 (1994); Shimazaki M., Okazaki F., Nakajima F., Ishikawa T., Ohta A., *Heterocycles*, **36**, 1823-1836 (1993); Houk K. N., Moses S. R., Wu Y.-D., Rondan N. G., Jäger V., Schohe R., Fronczek F. R., *J. Am. Chem. Soc.*, **106**, 3880-3882 (1984).
- 10) **13**: Colorless oil. IR (CHCl₃) cm^{-1} : 3402 (OH), 1731 (CO₂). ^1H -NMR (500 MHz, CDCl₃) δ : 4.12 (1H, q, $J=3$ Hz, 4-H), 3.36 (1H, br t, $J=2$ Hz, 3-H), 3.27 (1H, br q, $J=6$ Hz, 2-H), 2.98 (1H, m, 6-H), 2.30 (2H, t, $J=7$ Hz, 5'-H₂), 1.67 (1H, br dt, $J=14, 2.5$ Hz, 5-Heq), 1.57 (1H, ddd, $J=14, 11, 3$ Hz, 5-Hax), 1.18 (3H, d, $J=6.5$ Hz, 2-Me). $[\alpha]_{\text{D}}^{-11^\circ}$ ($c=1.43$, MeOH). **14**: Colorless oil. IR (CHCl₃) cm^{-1} : 3441 (OH), 1731 (CO₂). ^1H -NMR (500 MHz, CDCl₃) δ : 3.71 (1H, ddd, $J=11, 8.5, 5$ Hz, 4-H), 3.42-3.37 (2H, m, 2, 3-H), 2.82 (1H, m, 6-H), 2.27 (2H, t, $J=7$ Hz, 5'-H₂), 2.03 (1H, ddd, $J=13, 5, 3$ Hz, 5-Heq), 1.11 (3H, d, $J=7$ Hz, 2-Me) 1.06 (1H, br dt, $J=13, 11.5$ Hz, 5-Hax). $[\alpha]_{\text{D}}^{-46^\circ}$ ($c=0.89$, MeOH). **15**: Colorless oil. IR (CHCl₃) cm^{-1} : 3443 (OH), 1732 (CO₂); ^1H -NMR (500 MHz, CDCl₃) δ : 3.53 (1H, ddd, $J=11, 5, 4$ Hz, 4-H), 3.48 (1H, br s, $W_{1/2}=5.5$ Hz, 3-H), 2.73 (1H, qd, $J=6.5, 1$ Hz, 2-H), 2.54 (1H, m, 6-H), 2.30 (2H, t, $J=7.5$ Hz, 5'-H₂), 1.76 (1H, br dt, $J=12, 3$ Hz, 5-Heq), 1.23 (1H, br q, $J=12$ Hz, 5-Hax), 1.12 (3H, d, $J=6.5$ Hz, 2-Me). $[\alpha]_{\text{D}}^{+7^\circ}$ ($c=1.02$, MeOH). **17**: Colorless oil. ^1H -NMR (300 MHz, CDCl₃) δ : 3.61 (1H, m, $W_{1/2}=6.5$ Hz, 3-H), 2.84 (1H, qd, $J=6.5, 1.5$ Hz, 2-H), 2.61 (1H, m, 6-H), 2.31 (2H, t, $J=7$ Hz, 5'-H₂), 1.94 (1H, br dq, $J=10.5, 3$ Hz, 4-Heq), 1.69—1.24 (11H, m, 4-Hax, 5-H₂, 1"—4"-H₂), 1.18 (3H, d, $J=6.5$ Hz, 2-Me).
- 11) Barton D. H. R., McCombie S. W., *J. Chem. Soc., Perkin Trans. 1*, **1975**, 1574-1585; Rasmussen J. R., Slinger C. J., Kordish R. J., Newman-Evans D. D., *J. Org. Chem.*, **46**, 4843-4846 (1981).

(Received April 19, 1996; accepted May 14, 1996)