

A CONVENIENT SYNTHESIS OF BENZO[C]PHENANTHRIDINE ALKALOID, CHELERYTHRINE, BY THE PALLADIUM-ASSISTED INTERNAL BIARYL COUPLING REACTION


Takashi HARAYAMA,* Toshihiko AKIYAMA, and Kazuko KAWANO

Faculty of Pharmaceutical Sciences, Okayama University, Tsushima-naka 1-1-1, Okayama 700, Japan

Total synthesis of chelerythrine, a benzo[*c*]phenanthridine alkaloid, was accomplished *via* the internal aryl-aryl coupling reaction of halo-amide (4) by the palladium-assisted cyclization reaction.


KEY WORDS internal biaryl coupling; Heck type reaction; halo-amide; benz[*c*]phenanthridine alkaloid; chelerythrine

Benzo[*c*]phenanthridine alkaloids have attracted much attention because of their potent pharmacological activities.¹⁾ Recently, it was reported that chelerythrine (1)²⁾ and isofagaridine (2)³⁾ inhibited protein kinase C and DNA topoisomerase I, respectively. Although several reports on the total synthesis of chelerythrine have been published,⁴⁾ the methods had some disadvantages such as many steps, low yield, and/or no generality. Therefore, we planned to develop a convenient synthetic method for 1.



The cross-coupling reaction with palladium catalyst has been an extremely useful tool in organic synthesis.⁵⁾ We designed the synthetic plan for 1 involving an internal biaryl cyclization by palladium as a key reaction,⁶⁾ as shown in Chart 1. Since the coupling product (3), oxychelerythrine, had already been converted into chelerythrine (1),^{4e)} the synthesis of 3 indicates a formal synthesis of 1. It was reported that an internal coupling reaction of bromo-amide (7b) with a Pd reagent proceeded in 50% yield.^{6a)} In order to improve the yield, the reaction was re-examined using purified Pd(OAc)₂,⁷⁾ phosphine ligand and base. The results are summarized in Table 1. On

* To whom correspondence should be addressed.

**Table 1.** Results of Cyclization Reaction of 2-Halo-*N*-methyl-*N*-phenylbenzamide (**7**)^{a)}

X	Run	Pd(OAc) ₂ (eq.)	Ligand	Base	Solvent	Temp.	Time	Yield (%)	
								8	S.M.
I	1	0.05	PPh ₃	Ag ₂ CO ₃	DMF	Refl.	40 min	79	—
	2	0.2	PPh ₃	Ag ₂ CO ₃	DMF	Refl.	15 min	93	—
	3	0.2	POT	Ag ₂ CO ₃	DMF	Refl.	15 min	93	—
	4	0.2	PPh ₃	Ag ₂ CO ₃	DMF	30-35°C	35 h	85	—
	5	0.2	PPh ₃	Ag ₂ CO ₃	Xylene	30-35°C	23 h	93	—
	6	0.2	PPh ₃	Ag ₂ CO ₃	Benzene	Refl.	10 min	98	—
	7	0.2	PPh ₃	Ag ₂ CO ₃	CH ₃ CN	Refl.	15 min	95	—
	8	0.2	PPh ₃	<i>i</i> Pr ₂ NEt	DMF	Refl.	4.5 h	21	7
	9	0.2	PPh ₃	<i>i</i> Pr ₂ NEt	Benzene	Refl.	6 h	45	14
Br	10	1.0	PPh ₃	Ag ₂ CO ₃	DMF	Refl.	60 h	75	7
	11	0.2	POT	Ag ₂ CO ₃	DMF	Refl.	1.5 h	99	—


a) All reactions were carried out using Pd(OAc)₂ and ligand in a ratio of 1 : 2 and 2 mol equivalent of base.

Chart 2

Table 2. Results of Cyclization Reaction of 6-Halo-2,3-dimethoxy-*N*-methyl-*N*-(6,7-methylenedioxy-1-naphthyl)benzamide (**4**) to Oxachelerythrine (**3**)^{a)}

X	Run	Pd(OAc) ₂ (eq.)	Ligand	Base	Solvent	Temp.	Time	Yield (%)
								3
I	1	0.2	PPh ₃	Ag ₂ CO ₃	DMF	Refl.	20 min	85
	2	0.2	POT	Ag ₂ CO ₃	DMF	Refl.	20 min	94
Br	3	0.2	PPh ₃	Ag ₂ CO ₃	DMF	Refl.	2 h	79
	4	0.2	POT	Ag ₂ CO ₃	DMF	Refl.	3 h	96

a) All reactions were carried out in the presence of ligand (0.4 eq) and base (2 eq).

using 0.2 eq of $\text{Pd}(\text{OAc})_2$, PPh_3 , and Ag_2CO_3 , the solvent had no crucial effect on the internal coupling reaction of iodo-amide (**7a**), although Ag_2CO_3 was superior to Hünig base. On the other hand, the coupling reaction of bromo-amide (**7b**) proceeded slowly even when using a stoichiometric amount of $\text{Pd}(\text{OAc})_2$ in the presence of PPh_3 as a ligand in DMF (see run 10 in Table 1). However, on using tris(2-methylphenyl)phosphine (POT) as ligand, the reaction proceeded smoothly using 0.2 eq of $\text{Pd}(\text{OAc})_2$ and gave phenanthridone (**8**) in an excellent yield (see run 11 in Table 1).

Next, we applied these methods to the synthesis of oxychelerythrine (**3**) from halo-amides (**4**), which were prepared from iodo-acid (**5a**)⁸ or bromo-acid (**5b**)⁹ and naphthylamine (**6**)¹⁰ as shown in Chart 2. The results are summarized in Table 2. The coupling reaction of both halo-amides (**4**) with $\text{Pd}(\text{OAc})_2$, PPh_3 or POT and Ag_2CO_3 in DMF under reflux afforded oxychelerythrine (**3**) in an excellent yield, although iodo-amide (**4a**) was more reactive than bromo-amide (**4b**).

In conclusion, the present method using the Pd reagent is very convenient and effective for preparing benzo[c]phenanthridine alkaloids. We are now investigating the generality of this method.

REFERENCES AND NOTES

- 1) a) Simanek V., "The Alkaloids," Vol. 26, ed. by Brossi A., Academic Press. Inc., New York, 1983, pp185-240; b) Dostal J., Potacek M., *Collect. Czech. Chem. Commun.*, **55**, 2840-2873 (1990).
- 2) Herert J. M., Augereau J. M., Gleye J., Maffrand J. P., *Biophys. Res. Commun.*, **172**, 993-999 (1990).
- 3) Fang S.-D., Wang L.-K., Hecht S. M., *J. Org. Chem.*, **58**, 5025-5027 (1993).
- 4) a) Bailey A. S., Warthing C. R., *J. Chem. Soc.*, **1956**, 4535-4543; b) Ninomiya I., Yamamoto O., Naito T., *J. Chem. Soc., Perkin Trans. I*, **1983**, 2171-2174; c) Kesser S. V., Gupta Y. P., Barakrishnan P., Sawal K. K., Mohamamad T., Dutt M., *J. Org. Chem.*, **53**, 1708-1713 (1988); d) Smidrkal J., *Collect. Czech. Chem. Commun.*, **49**, 1412-1420 (1984); e) Hanaoka M., Motonishi T., Mukai C., *J. Chem. Soc., Perkin Trans. I*, **1986**, 2253-2256; Ishii H., Takeda S., Ogata K., Hanaoka M., Harayama T., *Chem. Pharm. Bull.*, **39**, 2712-2714 (1991); f) Ishii H., Ishikawa T., Takeda S., Suzuki M., Harayama T., *Chem. Pharm. Bull.*, **40**, 2002-2006 (1992); g) Martin G., Guitian E., Castedo L., *J. Org. Chem.*, **57**, 5907-5911 (1992).
- 5) a) Tsuji J., "Palladium Reagents and Catalysts," John Wiley & Sons Inc., New York, 1995, pp 125-252; b) Knight D.W., "Comprehensive Organic Synthesis," Vol. 3, ed by Trost B. M. Fleming I., Pergamon Press, Oxford, 1991, pp 481-520.
- 6) a) Ames D. E., Opaeko A., *Tetrahedron*, **40**, 1919-1925 (1984); b) Bringmann G., Walter R., Weirich R., *Angew. Chem. Int. Ed. Engl.*, **29**, 977-991 (1990) and references cited therein; c) Hosoya T., Takashiro E., Matsumoto T., Suzuki K., *J. Am. Chem. Soc.*, **116**, 1004-1015 (1994) and references cited therein; d) Deshpande P., Martin O. R., *Tetrahedron Lett.*, **31**, 6313-6316 (1990); e) Hosoya T., Takashiro E., Matsumoto T., Suzuki K., *Tetrahedron Lett.*, **35**, 4591-4594 (1994).
- 7) Ohrai K., Kondo K., Sodeoka M., Shibasaki M., *J. Am. Chem. Soc.*, **116**, 11737-11747 (1994).
- 8) Dyke S. F., Tiley E. P., *Tetrahedron*, **31**, 561-568 (1975).
- 9) Auerbach J., Weisman S. A., Blacklock T. J., Angeles M. R., Hoogsteen K., *Tetrahedron Lett.*, **34**, 931-934 (1993).
- 10) Naphthylamine (**6**) was prepared from 2,3-dimethanesulfonyloxy-5-nitro-naphthalene¹¹) via three steps in a total yield of 35%; i) hydrolysis with 2% NaOH, ii) methylenation with CH_2Br_2 in the presence of CsF in DMF, iii) catalytic reduction with Pd-C/H₂ in THF.
- 11) Stermitz F. R., Gillespie J. P., Amoros L. G., Romero R., Stermitz T. A., *J. Med. Chem.*, **18**, 708-713 (1975).

(Received June 13, 1996; accepted July 15, 1996)