

Heterobimetallic Templates for Carbon–Carbon Bond Formation by Migratory Insertion Reactions Involving CO, Isonitriles or Olefins


Pierre Braunstein, Michael Knorr and Thomas Stährfeldt

Laboratoire de Chimie de Coordination, URA CNRS 0416, Université Louis Pasteur, 4 rue Blaise Pascal, F-67070 Strasbourg Cédex, France


New patterns for heterobimetallic reactivity are presented that combine electronic effects (of the metal–metal bond) and steric control (by a bridging, hemilabile trialkoxysilyl ligand) in organometallic insertion reactions involving isocyanides, CO and olefins.

The insertion of small molecules, such as CO, organic isocyanides and olefins, into metal–carbon bonds has been recognized as an elementary step of considerable importance in organometallic chemistry and catalysis.¹ Several mononuclear Pd^{II} complexes have been described recently which allow stepwise successive insertion reactions (*e.g.* leading to polyketone chain growth).^{2–4} We report here on the first studies performed with a heterometallic complex designed (i) to possess the migrating group *trans* to a metal–metal bond of high *trans*-influence and (ii) to contain an intramolecular donor which, by successive ‘opening and closing’, provides in the opened state the coordination site required for the incoming substrate and, in the closed state, the stabilisation of the inserted product (Scheme 1).

Relatively little has been reported on insertion reactions with heterometallic alkyls, perhaps because of the limited number of such complexes and/or their instability.^{5,6} We have now prepared the stable bimetallic alkyls [(Me)M(μ-dppm)Fe{Si(OMe)₃}(CO)₃] (**1**, M = Pd; **2a**, M = Pt; dppm = Ph₂PCH₂PPh₂) by reaction of [M(Me)Cl(1,5-cod)]⁷ with [Fe{Si(OMe)₃}(CO)₃(dppm-P)][–] in THF, whereas [(Et)Pt(μ-dppm)Fe{Si(OMe)₃}(CO)₃] **2b** was obtained by an oxidative addition–insertion reaction between [HFe{Si(OMe)₃}(CO)₃(dppm-P)]⁸ and [Pt(C₂H₄)₃] in toluene. These complexes contain an η²-μ₂-Si–O bridge (1H




Scheme 1 Substrate coordination/migratory insertion sequence resulting from the opening and bridging behaviour of the alkoxy-silyl ligand (M = Pd, Pt)




Scheme 2

and ³¹P{¹H} NMR evidence) that is dynamic on the ¹H NMR time scale in the case of palladium, the exchange between the OMe groups resulting most likely from rapid rotation of the –Si(OMe)₃ ligand about the Fe–Si bond.⁹ The donor interaction SiO → Pt in complexes **2** is displaced by CO (CH₂Cl₂, 20 °C) to give **3** in which this ligand occupies the suitable *cis* position with respect to the alkyl group. These complexes represent rare examples in which a CO and an alkyl ligand bound *cis* to each other in a square planar environment have been observed under ambient conditions.^{7a,10a,c} However, after *ca.* 30 min, acyl formation has occurred which liberates a coordination site and the dative O → Pt intramolecular interaction is restored, resulting in complexes **4**. The latter bind CO reversibly, since bubbling of CO through their CH₂Cl₂ solution quantitatively affords **5**, which lose their terminal Pt-bound CO on purging (but not the acyl CO) (Scheme 2). All the successive ‘opening and closing’ steps involving the trialkoxysilyl ligand may thus be easily identified in the Fe–Pt systems. They will also form the basis for the behaviour of analogous Fe–Pd complexes, although the much higher reactivity of the latter has precluded characterization of all the individual steps.


Under a CO atmosphere, the bimetallic alkyl **1** in CH₂Cl₂ afforded within 1 h the acyl complex **6** in good yields. In contrast to most palladium acyl complexes, **6** is stable with respect to CO de-insertion. Reaction of **1** with 1 equiv. C≡N-(*o*-anisyl) yielded the imino acyl complex **7**, analogous to **6** (Scheme 3). Reaction with two equivalents C≡N-Bu¹ selectively gave **8**, whereas with the more reactive C≡N-(2,6-xylyl), bis-insertion occurred readily to give **9** selectively. In the presence of excess C≡N-(2,6-xylyl), an equilibrium was established with **10** which contains a Pd-bound terminal isocyanide [ν(CN) = 2173 cm^{–1}] in place of the dative O → Pd bond. The reaction of **1** with two equiv. C≡N-(*o*-anisyl) gave a mixture of mono-, di- and tri-insertion products. Selective tri-insertion of isocyanide leading to **11** was achieved when 3 equiv. C≡N-(*o*-anisyl) were used. Further addition of isocyanide



Scheme 3 Reactions in CH₂Cl₂, –10 to 25 °C. i, 1 equiv. C≡NR¹ [R¹ = C₆H₄ (*o*-OMe)]; ii, 2 equiv. C≡NR² [R² = Bu¹]; iii, 2 equiv. C≡NR³ [R³ = C₆H₃Me₂-2,6]; iv, 3 equiv. C≡NR¹



Scheme 4



opens again the dative $O \rightarrow Pd$ interaction to give **12** [$\nu(CN) = 2186\text{ cm}^{-1}$], en route to the next migratory insertion.¹¹

Olefin insertion into the Pd-acyl bond of **6** occurred quantitatively after 3 h in the presence of norbornene (nbe) to give **13** whose 1H NMR data indicate that *cis* addition of Pd-C(O)Me across the exo face of norbornene has occurred (Scheme 4). The IR data confirm the expected coordination of the acyl group to palladium, resulting in a five-membered chelate rather than the less favoured $Fe-Si-O \rightarrow Pd$ four-membered ring.^{12,13}

Further CO insertion into the newly formed Pd-C bond of the norbornadiene (nbd) analogue of **13** took place under a CO atmosphere and this reaction was accelerated in the presence of excess nbd. It led to stable **14** which contains an oligomeric chain with alternating CO and nbe units whose exact length is not known at the moment (mononuclear complexes related to **14** with $n = 1-3$ have been characterized by IR, 1H , $^{13}C\{^1H\}$ and $^{31}P\{^1H\}$ NMR).^{10b} This indicates that olefin insertion into the Pd-acyl bond is more facile than CO insertion into a Pd-alkyl bond. No olefin insertion was observed in the case of the platinum complex **4a** (ethylene or nbd, CH_2Cl_2 , $20^\circ C$).

The mechanism of these insertion reactions obeys the following pattern: opening of the labile $SiO \rightarrow Pd$ bond, coordination of the substrate, *cis* migration, isomerization and closing of the $Fe-Si-O \rightarrow Pd$ ring. The unique behaviour of the $-Si(OMe)_3$ ligand in these systems extends to bimetallic chemistry the concept of hemilabile or 'dangling' chelating ligands, which has been successfully used in mononuclear chemistry.¹⁴ The bimetallic nature of all the complexes presented here is readily ascertained in solution by $^{31}P\{^1H\}$ NMR where an AX pattern is always observed. The $^{2+3}J(PP)$ values are diagnostic for the presence or absence of an $\eta^2-\mu-Si-O$ bridge: in the former case, this coupling constant is *ca.* 40 Hz smaller than in the latter. Although pentacoordinated intermediates have not been detected and need not be invoked in this mechanism, their involvement in the course of the displacement of the $O \rightarrow M$ bond cannot be excluded.

The favourable conditions for (i) *cis* coordination of the entering substrate and (ii) *cis* migration of the alkyl group are built in the design of our bimetallic system since (i) the thermodynamically preferred *trans*-P-Fe-Si arrangement forces the labile, dative $SiO \rightarrow M$ bond to be *cis* to the M-R bond and (ii) the high *trans* influence of the metal-metal bond, which is always *trans* to the M-R bond, promotes migration of the R group. These features emphasize the interesting chemo- and stereochemical control resulting from the appropriate interplay between metal-metal and metal-ligand interactions in heterometallic systems.

We thank the Centre National de la Recherche Scientifique and the Deutsche Forschungsgemeinschaft for support and an Associate Research position for M. K. and the Commission of the European Communities for a PhD grant to T. S. (contract 913014).

Received, 5th May 1994; Com. 4/02665I

Footnote

† Selected NMR spectroscopic data at 298 K. $^{31}P\{^1H\}$ NMR (121.5 MHz) spectra were referenced to external H_3PO_4 . J values given in Hz. Complex **1**, 1H NMR (300 MHz, C_6D_6): δ 0.82 (s, 3H, PdMe), 3.49 [t, 2H, PCH₂P, $^2J(PH)$ 10.1], 3.91 (s, vbr, 9H, OMe); $^{31}P\{^1H\}$ NMR (C_6D_6): δ 42.2 [d, $^{2+3}J(PP)$ 51, P(Pd)], 63.5 [d, P(Fe)]. Complex **3a** was precipitated as a yellowish powder by rapid addition of pentane to its solution. IR ν (CH_2Cl_2)/ cm^{-1} : 2084m (PtCO); $^{31}P\{^1H\}$ NMR ($CDCl_3$): δ 23.2 [d, P(Pt), $^{2+3}J(PP)$ 81, $^1J(PtP)$ 3319], 62.8 [d, P(Fe), $^2J(PtP)$ 34]. Complex **4a**, IR ν (CH_2Cl_2)/ cm^{-1} : 1634m (acyl); $^{31}P\{^1H\}$ NMR ($CDCl_3$): δ 3.2 [d, P(Pt), $^{2+3}J(PP)$ 44, $^1J(PtP)$ 5564], 63.3 [d, P(Fe), $^2J(PtP)$ 84]. Complex **5a**, IR ν (CH_2Cl_2)/ cm^{-1} : 2086m (PtCO), 1676w (acyl); $^{31}P\{^1H\}$ NMR ($CDCl_3$): δ 11.8 [d, P(Pt), $^{2+3}J(PP)$ 81, $^1J(PtP)$ 3687], 62.3 [d, P(Fe), $^{2+3}J(PtP)$ 32]. Complex **6**, $^{31}P\{^1H\}$ NMR ($CDCl_3$): δ 26.7 [d, P(Pd), $^{2+3}J(PP)$ 56], 65.2 [d, P(Fe)]. Complex **7**: $^{31}P\{^1H\}$ NMR (C_6D_6): δ 28.9 [d, P(Pd), $^{2+3}J(PP)$ 53], 62.7 [d, P(Fe)]. Complex **8**, IR ν (CH_2Cl_2)/ cm^{-1} : 2200m (PdCNR), 1648m (iminoacyl); $^{31}P\{^1H\}$ NMR (C_6D_6): δ 17.2 [d, P(Pd), $^{2+3}J(PP)$ 83], 63.4 [d, P(Fe)]. Complex **9**, IR ν (CH_2Cl_2)/ cm^{-1} : 1644m, 1610m (iminoacyl); 1H NMR (C_6D_6): δ 1.71 (s, 3H, CH₃), 2.13 (s, 6H, xylyl-Me), 2.18 (s, 6H, xylyl-Me), 3.44 [t, 2H, PCH₂P, $^2J(PH)$ 10.1], 3.86 [s, 9H, (MeO)₃Si]. Complex **11**, 1H NMR (C_6D_6 , 308 K): δ 1.74 (s, 3H, MeC), 4.00 [s, 9H, (MeO)₃Si]. Complex **12**, $^{31}P\{^1H\}$ NMR (C_6D_6): δ 20.9 [d, P(Pd), $^{2+3}J(PP)$ 87], 63.1 [d, P(Fe)]. Complex **13**, IR ν (CH_2Cl_2)/ cm^{-1} : 1645m ($C=O \rightarrow Pd$); 1H NMR ($CDCl_3$): δ 2.36 [s, 3H, C(O)Me], 3.64 [s, 9H, (MeO)₃Si]. Complex **14**, $n = 2$: IR ν (KBr)/ cm^{-1} : 1708m (ketonic C=O), 1635m ($C=O \rightarrow Pd$); $^{31}P\{^1H\}$ NMR ($CDCl_3$): δ 35.0 [d, P(Pd), $^{2+3}J(PP)$ 63], 63.9 [d, P(Fe)]; $^{13}C\{^1H\}$ NMR ($CDCl_3$): δ 29.3 [s, C(O)Me], 44.1 [t, CH₂P, $^1J(PC)$ 18 Hz], 231.5 (s, $C=O \rightarrow Pd$).

References

- (a) G. K. Anderson and R. J. Cross, *Acc. Chem. Res.*, 1984, **17**, 67; (b) A. Yamamoto, *Organotransition Metal Chemistry*, Wiley, New York, 1986; (c) L. D. Durfee and I. P. Rothwell, *Chem. Rev.*, 1988, **88**, 1059.
- T. W. Lai and A. Sen, *Organometallics*, 1984, **3**, 866.
- M. Brookhart, F. C. Rix and J. M. DeSimone, *J. Am. Chem. Soc.*, 1992, **114**, 5894.
- E. Drent, J. A. M. v. Broekhoven and M. J. Doyle, *J. Organomet. Chem.*, 1992, **417**, 235.
- (a) D. M. Antonelli and M. Cowie, *Organometallics*, 1991, **10**, 2550; (b) F. Antwi-Nsiah and M. Cowie, *Organometallics*, 1992, **11**, 3157; (c) M. Ferrer, O. Rossell, M. Seco and P. Braunstein, *J. Chem. Soc., Dalton Trans.*, 1989, 379.
- (a) B. Longato, J. R. Norton, J. C. Huffman, J. A. Marsella and K. G. Caulton, *J. Am. Chem. Soc.*, 1981, **103**, 209; (b) L. W. Arndt, B. T. Bancroft, M. Y. Daresbourg, C. P. Janzen, C. M. Kim, J. Reibenspies, K. E. Varner and K. A. Youngdahl, *Organometallics*, 1988, **7**, 1302.
- (a) G. P. C. M. Dekker, A. Buijs, C. J. Elsevier, K. Vrieze, P. W. N. M. van Leeuwen, W. J. J. Smeets, A. L. Spek, Y. F. Wang and C. H. Stam, *Organometallics*, 1992, **11**, 1937; (b) F. T. Lapido and G. K. Anderson, *Organometallics*, 1994, **13**, 303.
- P. Braunstein, M. Knorr, U. Schubert, M. Lanfranchi and A. Tiripicchio, *J. Chem. Soc., Dalton Trans.*, 1991, 1507.
- (a) P. Braunstein, M. Knorr, H. Piana and U. Schubert, *Organometallics*, 1991, **10**, 828; (b) P. Braunstein, M. Knorr, A. Tiripicchio and M. Tiripicchio-Camellini, *Angew. Chem., Int. Ed. Engl.*, 1989, **28**, 1361.
- (a) I. Tóth and C. J. Elsevier, *J. Am. Chem. Soc.*, 1993, **115**, 10388; (b) R. v. Asselt, E. E. C. G. Gielens, R. E. Rülke, K. Vrieze and C. J. Elsevier, *J. Am. Chem. Soc.*, 1994, **116**, 977; (c) G. K. Anderson and R. J. Cross, *J. Chem. Soc., Dalton Trans.*, 1979, 1246.
- Y. Yamamoto and H. Yamazaki, *Inorg. Chem.*, 1974, **13**, 438.
- J. S. Brumbaugh, R. R. Whittle, M. Parvez and A. Sen, *Organometallics*, 1990, **9**, 1735.
- B. A. Markies, M. H. P. Rietveld, J. Boersma, A. L. Spek and G. van Koten, *J. Organometal. Chem.*, 1992, **424**, C12.
- (a) G. K. Anderson and G. J. Lumetta, *Organometallics*, 1985, **4**, 1542; (b) E. Lindner, J. Dettinger, R. Fawzi and M. Steimann, *Chem. Ber.*, 1993, **126**, 1347; (c) H. Jin and K. J. Cavell, *J. Chem. Soc., Dalton Trans.*, 1994, 415.