

Chemical Reviews

Volume 92, Number 8

December 1992

Four-Membered Ring Compounds Containing Bis(methylene)cyclobutene or Tetrakis(methylene)cyclobutane Moieties. Benzocyclobutadiene, Benzodicyclobutadiene, Biphenylene, and Related Compounds

Fumio Toda*

Department of Industrial Chemistry, Faculty of Engineering, Ehime University, Matsuyama 790, Japan

Peter Garratt*

Christopher Ingold Laboratories, Department of Chemistry, University College London, 20 Gordon Street, London WC1H 0AJ, UK


Received September 9, 1988 (Revised Manuscript Received May 15, 1992)

Contents

1 Introduction	1685
2 <i>n</i> -kis(methylene)cyclobutene Derivatives	1686
2.1 3,4-Bis(methylene)cyclobutene and Nonbenzannelated Derivatives	1686
2.1.1 Preparative Methods	1686
2.1.2 Structure and Spectroscopic Properties	1688
2.1.3 Reactions	1688
2.2 Cyclobutene-1,2-dione, 2,4-Bis(methylene)cyclobutane-1,3-dione, and Nonbenzannelated Derivatives	1689
2.2.1 Preparative Methods	1689
2.2.2 Structure and Spectroscopic Properties	1690
2.2.3 Reactions	1691
2.3 Tetrakis(methylene)cyclobutane and Nonbenzannelated Derivatives	1692
2.3.1 Preparative Methods	1692
2.3.2 Structure and Spectroscopic Properties	1693
2.3.3 Reactions	1693
3 Benzocyclobutadiene and Related Systems	1694
3.1 Preparative Methods	1694
3.2 Structure and Spectroscopic Properties	1695
3.3 Reactions	1696
4 Benzodicyclobutadiene Derivatives	1696
4.1 Reactions	1697
4.1.1 Oxidation	1697
4.1.2 Addition	1698
4.1.3 Complexes with Fe(CO) ₃	1699
4.1.4 Reaction with Dienophiles and Dienes	1699
4.1.5 Addition of Dichlorocarbene	1700
4.1.6 Insertion of Heteroatoms	1701
5 Biphenylene and Related Compounds	1702
5.1 Preparative Methods	1702
5.2 Structure and Spectroscopic Properties	1704
5.3 Reactions	1705
6 Conclusions	1705

1 Introduction

The 3,4-bis(methylene)cyclobutene group occurs in many chemical species. It is a strained entity in that sp^2 hybridization would require bond angles of 120° at each of the four carbons of the 4-membered ring. Nevertheless, 3,4-bis(methylene)cyclobutene (**1a**) is itself formed by a thermal rearrangement (section 2.1) and is a stable, well characterized compound. Formally introducing a double bond to join the methylene atoms gives benzocyclobutene (**2**),[‡] a compound of considerably less stability. The bis(methylene)cyclobutene structure **2a** is the most important of the contributing canonical structures, since structure **2b** incorporates a cyclobutadiene moiety. Annelating a benzene ring to




1a gives bis(methylene)benzocyclobutane **3**, and annelation to benzocyclobutadiene gives biphenylene (**4**). Both **3** and **4** are more stable than **2**, and in both of these systems canonical forms containing the dimethylenecyclobutene are important but not exclusive, structures containing tetrakis(methylene)cyclobutane ([4]radialene) moieties (**3b**, **4c**) making a major contribution.

[‡] Benzocyclobutene is the correct, but confusing, name for **2**, annelated systems ending in "ene" indicating complete unsaturation, unlike nonannelated nonaromatic systems. We prefer the name benzocyclobutadiene for **2** and will use it throughout this review.




Fumio Toda received his Bachelor's, Master's, and Doctor's degrees from Osaka University at Osaka city, Japan in 1956, 1958, and 1960, respectively. After staying at Osaka University for four years as an Instructor and in United States for two years as a post-doctoral fellow, he obtained an Associate Professorship at Ehime University in 1966. He was promoted to full Professor in 1970. His research interests are in the area of strained small-ring compounds, host-guest inclusion compounds, solid-state chemistry, and reaction control in crystals.


Extending the number of bis(methylene)cyclobutene structures from biphenylene can be done in two modes, linearly or angularly, as in 5 and 6, respectively. In 5 at least one 3,4-bis(methylene)cyclobutene structure must be retained (5a,c), whereas in 6 a structure with only tetrakis(methylene)cyclobutane moieties can be accommodated (6a).



4-Methylenecyclobuten-3-one (7), cyclobutene-3,4-dione (8), 3,4-bis(methylene)cyclobutane-1,2-dione (9), and 2,4-bis(methylene)cyclobutane-1,3-dione (10) have related structures with similar strain effects. Again, these entities can participate in benzannelated structures such as 11 and 12, but as with bis(methylene)-benzocyclobutane, other canonical structures make important contributions.



This review will examine the behavior of compounds possessing the bis(methylene)cyclobutene moiety, or a



Peter Garratt obtained a B.Sc. in Biological Chemistry from Bristol University in 1956 and, after a period in teaching and a year with J. K. N. Jones in Canada, obtained a Ph.D. with T. J. Katz at Columbia University in 1964. He then returned to the UK as a NATO Fellow to work with F. Sondheimer at Cambridge University and was appointed a University Demonstrator and Fellow of Fitzwilliam College in 1966. He moved to University College London in 1967 and was made a Reader in 1974. His current research interests are in strained compounds, organic conductors, inhibitors of enzymes early in the pyrimidine biosynthetic pathway, ion channel blockers, and probing the structure of the melatonin receptor.

related structure, and will examine the limits that can be placed on these structural entities and the effects that they engender. Compounds of this type were reviewed in the classic text of Cava and Mitchell,¹ and certain aspects of the chemistry of such compounds may be found in other reviews.^{2,3}


2 *n*-kis(methylene)cyclobutene Derivatives

2.1 3,4-Bis(methylene)cyclobutene and Nonbenzannelated Derivatives


2.1.1 Preparative Methods

3,4-Bis(methylene)cyclobutene (1a), although experiencing considerable bond-angle strain, is well known, being initially prepared, albeit in only 1% yield, by Blomquist and Maitlis in 1961 through the Hofmann degradation of 3,4-bis[(trimethylammonio)methyl]cyclobutene dihydroxide.⁴ Currently, the main synthetic route to 1a and substituted derivatives involves the electrocyclic rearrangement of hexa-1,2,4,5-tetraene or a substituted derivative. Thus 1a can be derived thermally from 13a.⁵ The tetramethyl derivative 1b can be similarly prepared from the corresponding tetramethylhexatriene 13b, a higher temperature being required (250 °C, cf. 150 °C).^{6,7} The *meso*-diallene 14 rearranged slowly at 150 °C but much more rapidly in boiling THF in the presence of copper(I) chloride,⁸ in both cases to give 15 by conrotatory ring closure. The same catalytic effect of copper(I) was also observed in the formation of 1a.⁷

The first, and still probably the best, preparative route to 3,4-bis(methylene)cyclobutene (1a) was by the thermal rearrangement of hexa-1,5-diyne (16), the initial [3,3] sigmatropic rearrangement to 13a being much slower than the subsequent electrocyclic ring closure of 13a to 1a.⁹⁻¹¹




Thermolysis of the dibromotetraene 18, readily prepared from the diol 17 by treatment with HBr, gave the dibromobis(diphenylmethylene)cyclobutene (19a) in almost quantitative yield.¹² The dichloride 19b could be similarly prepared, and the cyclobutenyl halides were reactive, the bromine in 19a being readily replaced by hydrogen to give 19c by treatment with LiAlH₄. Coupling of the bromoallene 20 with copper(I) chloride gave a mixture of the bis(methylene)cyclobutenes 22 and 23, presumably via the bisallene dimer 21.¹³




Pasto and co-workers^{14,15} have investigated the mechanism of the bisallene to bis(methylene)cyclobutene rearrangement and have confirmed the original observations of Skattebol and co-workers.⁸ In solution the reaction is catalyzed by Cu(I) salts, and the transformation proceeds at around 60 °C, whereas in the gas-phase temperatures of ca. 300 °C are required. Thermolysis of the *threo*-bisallene 24 gave a mixture of the two cyclobutenes 25 and 26 in the ratio 10.4:89.6

in the gas-phase reaction and 24.0:76.0 in solution. Both products are formed in a conrotatory process, compound 26 being much less sterically crowded than compound 25.




The mechanism of the rearrangement of 1,5-hexadiene was studied by Henry and Bergman,¹⁶ who showed that, after rearrangement, the deuterium atoms of 1,6-dideuteriohexa-1,5-diyne appeared in the 1,2-positions of 3,4-bis(methylene)cyclobutene (1a). At higher temperatures 1a rearranged to benzene and fulvene.


Treatment of the propargyl chlorides 27 with tetrakis(triphenylphosphoryl)nickel in benzene under argon gave the 2-dipropenylcyclobutenes (28).¹⁷ 2,7-Dimethylocta-2,3,5,6-tetraene (29) gave a trimer under these conditions but a mixture of 3,4-di-2-propylenecyclobutene (28, R = H) and 3,4,7,8-tetra-2-propylenecycloocta-1,5-diene (30) on reaction with dicyclooctadienenickel.¹⁸



Bis(methylene)cyclobutene (1a) can also be prepared by the retro-Diels-Alder cleavage of 31 at 425 °C, the latter being prepared in three steps from the Diels-Alder adduct of quadricyclane with maleic anhydride.¹⁹



Thermolysis of 1,2-diethynylcyclopropane (31a) at atmospheric pressure gave bicyclo[3.2.0]hepta-1,4,6-triene (33), presumably via the highly strained bisallene 32.²⁰ *cis*-1,2-Diethynylcyclobutane on thermolysis gives, by contrast, mainly 1,2-dihydropentalene and very little bicyclo[4.2.0]octa-1,5,7-triene.^{21,22} Bicyclo[6.2.0]deca-1,7,9-triene (36) is obtained when the tetrabromide 34 is treated with methyllithium at -30 °C, presumably through the intermediacy of the bisallene 35.²³



2.1.2 Structure and Spectroscopic Properties

A detailed microwave analysis has been carried out on **1a**.²⁴ The structure is shown in Figure 1. The molecule is flat, the C1–C4 and C2–C3 bonds are single, the bond angles in the ring are close to 90°, and the exocyclic bond angles are ca. 135°.


An earlier electron diffraction study by Skancke²⁵ had given values of values 135.7 pm for C₁–C₂, 148.8 pm for C₁–C₂ (C₃–C₄), and 151.6 for C₂–C₃. The compound has a large dipole moment (0.617 D) for an alternant hydrocarbon.^{26,24} In the ¹H NMR spectrum the cyclobutene protons are at δ 6.84 and the two types of methylene protons at δ 4.59 and 4.70.¹⁰ The detailed vapor infrared spectrum has been analyzed,²⁷ and the electronic spectrum is broad, showing a band at 240 nm ($\log \epsilon$ 4.2) and at 294 nm.^{4,27}

In 3,4-bis(diphenylmethylene)cyclobutene, the protons on the aromatic ring appear as two bands centered on δ 6.8 and 8.2 in the ratio 1:1. The higher field δ 6.8 band is attributed to the inner benzene rings which are mutually shielding (Figure 2), and this phenomena is observed in related derivatives.²⁸


A variety of calculations have been carried out for **1a**. The calculated enthalpy of formation is in good agreement with the experimental value determined from hydrogenation²⁹ and indicates that the system is nonaromatic. Earlier calculations had suggested that **1a** had a small, negative resonance energy per electron (REPE), comparable to that of benzocyclobutene (**2**) (**1a**, -0.028β , **2**, -0.027β).³⁰

2.1.3 Reactions

Although **1a** is sterically ideally set up as a Diels–Alder diene, this type of reaction does not occur, presumably because the ensuing product has to have a cyclobutadiene ring. Addition of diazomethane to **1a** gave the spiro derivatives **37–39**.³¹



Halogens add to the tetraphenyl derivatives **19a,b** on the cyclobutene double bond to give the corresponding tetrahalocyclobutanes **44a,b**.³² The endo double bond of the tetraphenyl derivatives **19a,c** is also photosensitive. Irradiation of a benzene solution of **19a** under an oxygen atmosphere gave **42a** as crimson prisms.³² The corresponding irradiation of **19c** gave 2,3-bis(diphenylmethylene)butane-1,2-dione (**43a**).^{33,34} This reaction probably proceeds via a dioxetane in-



Bond	Length (pm)	Angle	Degrees
C ₁ - C ₂	136.80	< C ₁ - C ₂ - C ₃	93.54
C ₂ - C ₃	147.92	< C ₂ - C ₃ - C ₄	86.46
C ₃ - C ₄	149.68	< C ₃ - C ₄ - C ₅	135.75
C ₄ - C ₅	133.79	< C ₄ - C ₅ - H ₅ (E)	133.68
C ₁ - H ₁	107.99	< H ₁ - C ₁ - C ₂	121.01
C ₅ - H ₅ (E)	108.59	< C ₄ - C ₅ - H ₅ (Z)	120.33
C ₅ - H ₅ (Z)	108.41		

Figure 1. Structure of 3,4-bis(methylene)cyclobutene by microwave spectroscopy.





Figure 2. Structure of **19** showing overlap of inner benzene rings.

intermediate whereas the photooxidation of **19a** may proceed through the radical intermediate **40**, which then undergoes a 1,2-shift of a bromine atom to give the radical **41**. Unlike the bromo derivative, the dichloride **19b** is inert to photooxidation but can be readily oxidized by dioxygen in the presence of a catalytic amount of diethyl ether peroxide to **43b**.




A comparison of the electronic spectra of the dihalocyclobutenes **19a–c** with the saturated derivatives, obtained after the addition of halogen, is revealing. The unsaturated compounds are colorless, with absorption bands in the ultraviolet, whereas the saturated derivatives are colored, with absorption bands in the visible region (Table 1). This finds a satisfying explanation in that the canonical contributions to **19** to allow conjugation require antiaromatic cyclobutadiene containing structures whereas those for **44** do not (Scheme 1).³⁵


Reaction of **19a** with AgClO₄ in the presence of water gave the hydroxycyclobutene derivative **45a**, while the same reagent in the presence of methanol gave the corresponding methoxy derivative **45b**. These compounds have very similar electronic spectra to **19**, presumably because of a similar dipolar contribution to the ground state (Scheme 1).

Table 1. Electronic Spectra of Bis(diphenylmethylene)-cyclobutene and -cyclobutane Derivatives


compd	solvent	λ_{\max} , nm ($10^{-2}\epsilon_{\max}$)
19a	CH ₃ Cl	290 (364)
19b	CH ₃ Cl	285 (364)
19c	CH ₃ Cl	280 (308)
44a	CH ₃ Cl	295 (135), 400 (162)
44b	CH ₃ Cl	290 (126), 388 (159)
44c	CH ₃ Cl	260 (235), 357 (209)
45a	EtOH	275 (131)
45b	EtOH	271 (158)
46a	EtOH	277 (103)
46b	EtOH	280 (154)

Scheme 1

The first derivative of 4-methylenecyclobut-2-enone, 3-(diphenylhydroxymethyl)-4-(diphenylmethylene)cyclobut-2-enone (**46a**), was obtained by treatment of **42b** with aqueous AgClO_4 .³⁵ Similar treatment of the dibromide **42a** gave the corresponding monobromo derivative **46b**.³⁵ The UV spectra of **46a** and **46b** are both similar to that of 3,4-bis(diphenylmethylene)cyclobutene (**19c**), again indicating that cyclobutadiene-like structures do not greatly contribute to the ground state. The IR spectra of **46a** and **46b** show the $\text{C}=\text{O}$ stretching bands at ca. 1770 cm^{-1} (**46a**, 1765 cm^{-1} ; **46b**, 1790 cm^{-1}), the higher frequency suggesting that the carbonyl group is not highly polarized, again attributable to little contribution from **46''**.



The corresponding ethers could not be prepared by substituting MeOH for H_2O in the above reactions. In the case of **42b**, treatment with AgClO_4 in the presence of MeOH gave **42d** by a simple substitution of bromine. The attempted preparation of **47a** by methanolysis of **46a** with MeOH in the presence of H_2SO_4 also gave **42d**, presumably through an acid catalyzed rearrangement. Treatment of **46a** with bromine led to the reformation of **42b** rather than substitution of the hydroxyl group. Presumably, whereas the tautomer **46a** is preferred over the nonconjugated form, the bromide and methoxyl tautomers **42b** and **42d** are more

Scheme 2


stable than the conjugated systems. This may reflect a subtle balance between ring strain (there are four $\text{sp}^3\text{-C}$ atoms in **47** and only three in **42d**) and conjugation (Scheme 2).

2.2 Cyclobutene-1,2-dione, 2,4-Bis(methylene)-cyclobutane-1,3-dione, and Nonbenzannulated Derivatives

2.2.1 Preparative Methods


The majority of methods for preparing cyclobutenediones rely on the hydrolysis of geminal dihalides or geminal hydroxy halides to form the carbonyl groups. The methods have been reviewed by Schmidt and Ried.³⁶ Methods for the construction of the 4-membered ring include photochemical and thermal $[2+2]$ -cycloaddition,³⁷⁻⁴⁰ dimerization of tetrahaloethene with subsequent nucleophilic addition, elimination,⁴¹⁻⁴³ thermal $[2+2]$ -cycloadditions between electron-rich and electron-poor ethenes,⁴⁴ intramolecular cyclization of hexatrienes,^{45,46} rearrangement of cyclopropene derivatives,^{47,48} and the addition of terakis[arylisocyanide] nickel to diphenylethyne.⁴⁹ The conversion of one cyclobutenedione, particularly the 2-halo or 2,3-dihalo derivatives, into another has also been a fruitful method of formation (see Schmidt and Ried, ref 36). General examples of some of these methods are shown in Scheme 3.

3,4-Bis(diphenylmethylene)cyclobutane-1,3-dione (**48**) was obtained in almost quantitative yield by treatment of **44a** with AgBF_4 in wet MeCN . Interestingly, increasing the water content to 10% gave **49** as the major product (44%) together with some **45a** (27%).³² Substituting AgClO_4 for AgBF_4 in wet MeCN gave 3,4-bis(diphenylmethylene)succinic anhydride (**50**) in 85% yield. In this latter reaction, **45a** is initially formed but is then oxidized by the HClO_4 produced in the course of the reaction. This is substantiated by the ready oxidation of **48** with aqueous HClO_4 , as well as with aqueous H_2O_2 and HNO_3 .³²




4-Methylenecyclobut-2-enone (**7**) was first prepared in 1973 by Trahanovsky and Park by the flash pyrolysis of furfuryl benzoate (**51**) at $640\text{--}700\text{ }^\circ\text{C}$.^{50,51} Martin and co-workers have prepared **7** in a similar manner to

Scheme 3




their preparations of 3,4-bis(methylene)cyclobutene (1a) and cyclobutene-1,3-dione (8) by the pyrolysis of the adduct **52**.^{19,52} Methylene cyclobutene was observed as a minor component in the pyrolysis of 2-propynyl propiolate.⁵³




2,4-Bis(methylene)cyclobutane-1,3-dione (**10**) has been prepared by the flash photolysis of **54**, which in turn was prepared by dimerization of the ketene **53**.^{54,55}

The dimer **54** reacted with methylenetriphenylphosphorane to give the monomethylene and bis(methylene) derivatives **55** and **56** and these in turn underwent flash

Scheme 4




pyrolysis to give tris(methylene)cyclobutanone (**57**) and tetrakis(methylene)cyclobutane (**58**) (Scheme 4).⁵⁵ Earlier, 2,4-bis(diphenylmethylene)cyclobutane-1,3-dione (**61**) had been prepared by the decomposition of the mixed anhydride **59**, presumably by way of the transient intermediate propadienone **60**.^{56,57}



2-(Diphenylmethylene)-4-benzylidenecyclobutane-1,3-dione (**63**) was prepared by base condensation of 2-(diphenylmethylene)cyclobutane-1,3-dione with benzaldehyde.⁵⁸ The dione **62** was prepared from the bromide **46b** by base-catalyzed hydrolysis followed by elimination of HBr and benzophenone and subsequent protonation of the resultant enolate anion. Condensation of **62** with diphenylcyclopropenone rather than benzaldehyde gave the methylenecyclopropene **64**.⁵⁸ Both **63** and **64** added methoxide under basic conditions to give the cyclobutenyl enolate ions **65** and **66** (Scheme 5).

2.2.2 Structure and Spectroscopic Properties


The ¹H NMR spectrum of 3-cyclobutene-1,2-dione (**8**) shows one signal at δ 9.73, the low-field value indicating a considerable contribution from dipolar forms. The IR spectrum shows a band at 1792 cm⁻¹, a considerably higher frequency than that of a simple ketone.³⁷ The IR spectrum of the bis(diphenylmethylene)cyclobutane-1,2-dione (**48**) also shows the carbonyl band at high frequency (1750 cm⁻¹). The microwave structure of 4-methylenecyclobut-2-enone (**7**) shows that it has a substantial dipole moment (3.39 \pm 0.05 D) with the parameters shown in Figure 3. The ¹H NMR spectrum shows four doublets of triplets at δ 8.66 (H³), 7.16 (H²), 5.01 (H^{5-Z}), and 4.78 (H^{5-E}). The



Bond	Length (pm)	Angle	Degrees
C1 - C2	147.5	O - C1 - C4	134.5
C2 - C3	137.5	C1 - C2 - H2	132.3
C3 - C4	147.5	C2 - C1 - C4	86.1
C1 - C4	156.6	C1 - C4 - C3	86.1
C4 - C5	134	C4 - C5 - H5	121.5
C1 - O	119		
C2 - H2 (C3 - H3)	108.3		
C5 - H5	108.6		

Figure 3. The structure of 4-methylenecyclobut-2-enone from the microwave spectrum.

Scheme 5




IR spectrum has bands at 1783 (C=O), 1688 (C=C), and 823 cm^{-1} (C=CH₂).

2,4-Bis(diphenylmethylene)cyclobutane-1,3-dione (**61**) has the carbonyl and methylene groups bent slightly away from each other out of the ring plane, the angle between the carbonyl carbon and the substituted carbon of the phenyl ring being 6.2°. 2,4-Bis(phenylmethylene)cyclobutane-1,3-dione is centrosymmetric and nearly planar, the phenyl groups being slightly twisted out of the plane of the 4-membered ring.⁶⁰


The IR spectrum of the 1,3-dione **64** shows a band at 1655 cm^{-1} for the C=O stretch, at a considerably lower frequency than those of **63** (1680 cm^{-1}) and **62** (1710 cm^{-1}) but at a comparable frequency to that of the methylenecyclopropene **67**,⁶¹ indicating a major contribution from the dipolar form. All of these carbonyl stretching frequencies are significantly lower than those of the cyclobutene-1,2-diones or methylenecyclobutane-1,2-diones (see 8 and 48). This is probably due in part to the smaller dipolar contribution to the 1,2-diones because of dipole-dipole interaction, but it also probably reflects the interaction between the two carbonyl groups in the latter systems. The ortho protons on the phenyl rings of **63** (δ 8.23–8.33), **64** (δ 8.47–8.60), and **67** (δ 8.65–8.80) are deshielded, largely through the diamagnetic anisotropy of the carbonyl group.⁶¹

In the UV spectrum the absorption bands for the 1,2-diones are at a much longer wavelength than those for the 1,3-diones (Table 2) and this may also reflect a smaller contribution of the dipolar form to the ground state in the 1,2-diones.



2.2.3 Reactions


The reactions of substituted cyclobutene-1,2-diones have been reviewed.⁶² The double bond is reactive and, for example, the proton of 3-phenylcyclobutene-1,3-dione (**68**) can be substituted by treatment with halogen to give the corresponding 4-halo derivative **69**.³⁹



Nitrogen and phosphorus ylides act as nucleophiles to displace bromine from **66**,⁶³ but Wittig reactions can also be carried out on the carbonyl groups.^{64–66}


1,4-Addition to the double bond can also occur to give keto enols (**68** → **70**), the latter being stabilized by hydrogen bonding. Besides acids and thiols, carbanions

Scheme 6





will add readily, giving the enol on protonation, and aromatic systems will add in the presence of Lewis acids (**68** → **71**). Ring-opening reactions can occur with anions and amines. For example, **68** gives the hydroquinone **72** with the anion derived from dibenzyl ketone and quinoxaline **73** with *o*-phenylenediamine (Scheme 6).⁶²


The bis(diphenylmethylene)butane-1,3-dione (**48**) reacts with *o*-phenylenediamine **75** in CCl₄ containing AcOH to give the quinoxaline **75** in 98% yield. If the reaction is carried out in CCl₄ containing CBrCl₃ as a source of radicals, then the diazocine **76** was obtained in 85% yield together with a little **75**.⁶⁷ In both cases addition of *o*-phenylenediamine to **48** gives the adduct **74**, which dehydrates to **75** or rearranges to **76** by loss of hydrogen atoms.



Photooxidation of diphenylcyclobutene-1,2-dione (77) in benzene or MeCN gave a mixture of 78 and 79.⁶⁸ Photoirradiation of 48 in 99% methanol gave 81 in 72% yield through initial ring opening to the bisketene 80 which oxidatively recycled to the anhydride 50 and this in turn photocyclized.⁶⁹



The diazo ketone 82, formed from the monotosyl hydrazone of 48 by treatment with basic alumina in $CHCl_3$, on irradiation in benzene containing a nucleophile gave the corresponding bis(diphenylmethylene)cyclopropane derivative. Thus with MeOH, 2-propanol, and aniline as nucleophiles the cyclopropanes 85a–c were obtained. The reaction presumably proceeds via a Wolff rearrangement of the initially formed ketocarbene 83 to the ketene 84.^{70–72} This is the first observation of a Wolff rearrangement of a diazo ketone in a 4-membered ring.




2.3 Tetrakis(methylene)cyclobutane and Nonbenzannelated Derivatives

2.3.1 Preparative Methods

Tetrakis(methylene)cyclobutane (58), [4]radialene, was first prepared and identified by Griffin and Peterson in 1962,^{73,74} but these authors noted that it had probably been prepared earlier by others as a component of mixtures. Their route to 58 involved as the last step a multiple 1,2-elimination from a number of symmetrical tetrasubstituted cyclobutanes, and the best precursors were found to be the tetrahalo derivatives 88. The synthetic route is shown in Scheme 7.

A number of other methods for the preparation of 58 have been reported, one of which was illustrated in section 2.2.1.⁵⁵ Hopf and co-workers^{75,76} adapted Hunts-


Scheme 7



man and Wristers rearrangement of hexadiyne to the protected octa-2,6-diyne-1,8-diol 89 which, on thermolysis, gave the substituted bis(methylene)cyclobutene 90. Hydrolysis, conversion of the resulting diol to the dihalide and elimination gave 58.

At the same time that Griffin and Peterson reported the preparation of 58, a report on the synthesis of tetrakis(diphenylmethylene)cyclobutane (102) appeared involving the photodimerization of tetraphenylbutatriene.⁷⁷ The structure derived for 102 was largely based on the products of ozonolysis (see below).


A series of phenyl-substituted derivatives of 58 could be prepared from 3,4-bis(diphenylmethylene)cyclobutane-1,2-dione (48).⁷⁸ Addition of MeMgI to 48 gave a 2:3 mixture of the *cis*- and *trans*-diols 91 in 57% yield. The mixture, on treatment with PBr_3 in pyridine, gave the [4]radialene 92. This is not a very stable compound, and it dimerizes readily at 60 °C in $CHCl_3$ to give 93.



Tris(diphenylmethylene)cyclobutanone (96) was prepared by the sequence of reactions shown in Scheme 8.⁷⁸ Treatment of 42c with benzaldehyde in the presence of copper powder and cyclohexyl isocyanide gave the condensation product 94 in 52% yield. Treatment of 48 with diphenyl ketene gave 96 in 43% yield, presumably via 95 and loss of CO_2 . Both 94 and 96 are stable compounds and serve as precursors to other tetrakis(methylene)cyclobutane derivatives.


Treatment of 94 with MeMgI followed by phosphorus tribromide in pyridine gave the thermally stable radialene 97. A similar treatment of 96 gave the corresponding radialene 98. Substituting $PhCH_2MgCl$ for MeMgI in these reactions gave the radialenes 99 and 100, respectively.⁷⁸ Both the monomethylene

Scheme 8




derivatives were obtained as yellow needles while the hexa- and heptaphenyl derivatives were red prisms.

Tetrakis(diphenylmethylene)cyclobutane (102) was prepared as deep red needles by the copper-assisted tetramerization of 1,1-dibromo-2,2-diphenylethene (101) and was, as were the radialenes 97–100, a stable substance.⁷⁹



The original report⁷⁷ that 102 could be formed from tetraphenylbutatriene (103) was shown to be incorrect,⁸⁰ the product being the asymmetric dimer 104. The products reportedly obtained by ozonolysis^{77,81} must also be incorrectly identified, and they differ in properties from the compounds above with the same assigned structure. Other compounds collated by Cava and Mitchell¹ in the chapter "Methylene Analogs of Cyclobutadienequinone" should also be considered to have different structures to those given.



An attempt to make 102 from the bromo ketone 42b failed. Reaction of 42b with Ph_3P gave the salt 105, which could be readily converted into the ylide 106

Table 2. Electronic Spectra of Phenyl-Substituted Tetraradialenes as Solutions in CHCl_3

compd	λ_{max} , nm ($10^{-2}\epsilon_{\text{max}}$)
92	262 (326), 320 sh (94), 405 (57)
97	280 (248), 350 sh (93), 425 (53)
99	306 (402), 373 (181), 450 sh (54)
98	284 (283), 340 sh (196), 480 sh (28)
100	298 (319), 400 sh (138), 490 (25)

with NaOMe. The ylide did not react with benzophenone, however, but it did react with ketene or phenyl isocyanate. The desired tris(methylene)cyclobutanes 107a,b cyclized spontaneously to the naphthalene derivatives 108a,b (Scheme 9).

2.3.2 Structure and Spectroscopic Properties


The ^1H NMR spectrum of tetrakis(methylene)cyclobutane (58) shows a singlet at δ 5.19, and in the IR spectrum there are absorption bands at 3095, 1760, 1710, 1400, and 880 cm^{-1} , the three high frequency bands being weak.⁷⁴ The IR and Raman spectra of 58 and its perdeuterated derivative are consistent with it having a planar ring with D_{4h} symmetry.⁸² The photoelectron spectrum indicates that the ground state of the molecular ion is $^2\text{B}_{1u}$ and the ionization energy of a π -electron is 8.35 eV.⁸³ Numerous papers have appeared involving calculations on 58.

As would be expected, in the phenylsubstituted derivatives the UV absorption bands move to longer wavelengths with increasing number of phenyl groups (Table 2).⁷⁸ An X-ray crystallographic structural analysis of the heptaphenyl derivative 100 shows that it has a puckered structure with the seven phenyl groups in an irregular arrangement.⁸⁴ In the octaphenyl derivative 102 the X-ray crystallographic structure shows it has the propeller-II rather than the propeller-I type structure, as shown in Figure 2,⁷⁹ and the 4-membered ring is very puckered (34.7°). The ^1H NMR spectrum of 102 shows a broad singlet at δ 6.80, and in the ^{13}C NMR spectrum the ring carbon is at δ 139.3 and the methylene carbon at δ 132.0.⁷⁹


2.3.3 Reactions

Tetrakis(methylene)cyclobutane (58) adds carbenes to form spiro derivatives. Dichlorocarbene initially adds twice to give mainly the 1,3-dimethylene derivative 109 rather than 110, and further treatment gives the tris and tetrakis adducts 111 and 112.⁷⁶ Dibromocarbene

Scheme 9



gives only the bis adduct and diazomethane gives a complex mixture.




Dienophiles react only once to give bis(methylene)cyclobutene derivatives, the second addition being precluded because cyclobutadienes would be formed.⁷⁴ Gas-phase pyrolysis at 880 °C gave styrene and *o*-xylene.⁷⁴ Addition of bromine gave 3,4-dibromotetrakis-(bromomethyl)cyclobutene, and addition of HBr gave 1-(bromomethyl)-3,4-dibromo-2,3,4-trimethylcyclobutene.⁷⁴ Hydrogenation gave *cis,syn,cis*-tetramethylcyclobutane.⁷⁴

3 Benzocyclobutadiene and Related Systems

Benzocyclobutadiene (2) is a highly reactive compound that readily undergoes dimerization, and for many years its occurrence could only be demonstrated by trapping experiments,¹ including the preparation of an iron tricarbonyl complex.⁸⁵ Its spectroscopic properties have now been determined using either matrix isolation^{86,87} or fast-flow techniques.⁸⁸ Naphthocyclobutadiene derivatives (e.g. 122) and other annelated systems are much more stable and can be isolated.¹


3.1 Preparative Methods

The majority of the methods of preparing benzocyclobutadiene (2) involve elimination from 1,2-disubstituted benzocyclobutenes. Early methods involved elimination of halogen from 1,2-dihalobenzocyclobutadienes (113a,b) with zinc,¹ and these methods were modified to produce 2 in the gas phase before cooling to 20 K in a matrix.⁸⁶ α,α -Dibromo-*o*-xylene (114) can also be dehalogenated to 2 over magnesium at a higher temperature (440 °C). Trahanovsky and Fischer⁸⁸ produced 2 in their flow NMR experiment by desilating




the derivative 115 in CD_3CN by mixing with a CD_3CN solution of $\text{Bu}_4\text{N}^+\text{F}^-$.


Alternative syntheses of 2 involve the rearrangement of octa-3,5-diene-1,7-diyne. Reaction of (*Z,Z*)-1,8-bis-(trimethylsilyl)octa-3,5-diene-1,7-diyne (116) with NaOH in ethanol gave the dimer 117 of 2, presumably by cleavage of the silyl groups, electrocyclicization to cycloocta-1,2,4,6,7-pentaene followed by a second electrocyclicization.⁸⁹ Thermal rearrangement of 116 gave 1,2-bis(trimethylsilyl)cyclobutadiene (118), the simplest benzocyclobutadiene derivative so far isolated.⁹⁰




Thermolysis of substituted 1,2,3,4-tetramethyl-3,4-ethynylcyclobutenes 119 gave substituted benzocyclobutadienes 120, the rearrangement occurring via octadienediynes, as above.⁹¹ Some of these compounds (R = *t*-Bu, mesityl) are stable systems but others dimerize thermally (R = H, Me, Ph), the nature of the product depending on the substituent.




Naphthocyclobutadiene and anthrocyclobutadiene derivatives have been prepared by dehalogenation of the corresponding 1,2-dihalides (e.g. 121 \rightarrow 122).⁹² Naphthocyclobutene itself dimerizes to dinaphtho[*a,c*]cyclooctatetraene.⁹³



The heterocyclic analogues of benzocyclobutadiene have been prepared in which the benzene ring has been replaced by a 5-membered ring, either furan or thiophene. The parent systems were obtained by thermolysis of 2,3-diethynylloxirane (123a) and its sulfur analogue (123b).⁹⁴⁻⁹⁶ In the case of thienocyclobutadiene (124b), extrusion of sulfur was a competitive process in the rearrangement.



6,7-Diphenyl-3-thiabicyclo[3.2.0]hepta-1,4,6-triene (127) was prepared by a Wittig reaction between the diketone 125 and the bisylide 126.⁹⁷



There have been numerous syntheses of benzocyclobutenedione (**12**) following the original synthesis from 1,2-diiodobenzocyclobutene by Cava and his co-workers,^{98,99} and these methods have been extensively reviewed.¹⁰⁰⁻¹⁰² Many substituted benzocyclobutenediones have been prepared by modification of these methods and some examples are shown in Scheme 10.

3.2 Structure and Spectroscopic Properties


The ¹H NMR spectrum of **2** taken at a flow rate of 45 mL min⁻¹ shows three bands at δ 6.36, 6.26, and 5.78. At a slower flow rate of 3 mL min⁻¹ the two high-field signals are partially resolved as an AA', BB' system, showing these signals are due to the protons on the 6-membered ring.⁸⁸ The ¹H NMR spectrum of silyl derivative **118** shows signals at δ 6.30 and 5.75 in CD₃CN, with little solvent shift in CCl₄.⁹⁰ Table 3 lists the ¹H NMR spectrum chemical shifts for a number of benzocyclobutenes and related compounds. The high-field position of these bands, similar to the ring protons in 5,6-bis(methylene)cyclohexadiene, can be attributed to the paratropicity of the 8 π -system.

The ¹³C NMR spectrum of the tetramethyl derivative **120** (R = ^tBu) shows ring carbons at δ 146.3 (C-1), 117.3 (C-2), 132.1 (C-3), and 156.0 (C-7).⁹¹

The matrix IR spectrum of **2** shows bands at 737 and 700 cm⁻¹, attributed to deformation of the protons on the 6- and 4-membered rings, respectively.⁸⁶

The matrix electronic spectrum of **2** shows absorption maxima at 243, 246.5, 256, 264, 270, 281.5, and 289 nm,⁸⁶ and solution spectrum of **118** shows maxima at 236, 243, 404, 417 (sh), and 472 (sh) nm, the three long wavelength bands being weak.⁹⁰ The 1,2-diphenyl derivative **120** shows maxima at 207, 218 (sh), 278, and 468 nm⁹¹ and the 1,2-diphenylthienocyclobutadiene (**127**) maxima at 266.5, 273.5, and 447 nm.⁹⁷ The latter compound is thermochromic, being deep reddish orange at room temperature and pale orange at lower temperatures.

The photoelectron spectrum of **2** has the lowest state of ²B₁ symmetry (7.87 eV) and is consistent with there being little contribution of the biradical to the ground state.⁸⁷



The X-ray crystallographic structure of **120** (R = ^tBu) shows the 4-membered ring to be rectangular, with long bonds connected to the phenyl ring (153.4 pm), and short nonconnecting bonds (135.9, 141.6 pm).⁹¹ The bond angles in the 4-membered ring are close to 90°. Similar alternation of bonds in the 4-membered ring is seen in the X-ray crystallographic structure of **127**, the connected bonds being 151.5 pm and the nonconnecting bonds 144.0 and 137.5 pm, the bond angles again being close to 90° but, as with **120**, the angle at the ring junction being slightly smaller than the other.¹⁰⁵ The phenyl rings are twisted out of the plane of the 4-membered ring by 23°. The X-ray structures of the substituted benzocyclobutadienes **175** and **177** (section 4.1.4) both show long connected bonds (175, 155.4 pm;

Scheme 10





Table 3. Proton Chemical Shifts in the ¹H NMR Spectra of Benzocyclobutadiene and Related Compounds

compound	¹ H NMR spectrum chemical shifts, δ		
	H ¹	H ²	H ³
	6.36	6.26 6.30	5.78 5.75
118 	6.33 6.75	6.11 6.01	
124b 		6.16	
		6.5	

177, 155.0 pm), short nonconnecting bonds (175, 133.6 pm; **177**, 135.0 pm) and some bond alternation in the benzene ring, the "bis(methylene)cyclobutadiene" benzene bonds being shorter (175, 136.5 pm; **177**, 136.0 pm).


Scheme 11



3.3 Reactions


Benzocyclobutadiene (2) is thermally unstable and dimerizes to the angular dimer 117.¹ Naphtho[*b*]cyclobutadiene (129) is also thermally unstable, but in this case, dimerization gives the linear dimer 130.⁹³ More highly substituted benzocyclobutadienes dimerize to various products depending on the substituents. 1-*tert*-Butylbenzocyclobutadiene is thermally unstable and the angular dimer is obtained,¹⁰⁶ while the 3,4,5,6-tetramethylbenzocyclobutadienes give angular, linear, and rearranged dimers depending on the nature of the 1,2-substituents (Scheme 11).⁹¹

Benzocyclobutadiene (2) undergoes Diels–Alder addition with dienes, and this was the original method of demonstrating its formation as a reactive intermediate.¹ Benzocyclobutadienes can also act as dienes, as in the dimerization reactions, and 120 ($R = \text{tBu}$) reacts with tetracyanoethene (TCNE) and dicyanoethene across the 6-membered ring to give 131.¹⁰⁷ The thienocyclobutadiene 127 adds TCNE to give the adduct 133, presumably by a [2 + 2] addition followed by ring opening.⁹⁷ The bisilyl derivative 118 reacts with dimethyl acetylenedicarboxylate in a Diels–Alder manner to give dimethyl 3,4-bis(trimethylsilyl)naphthalene-1,2-dicarboxylate.⁹⁰




Benzocyclobutadiene and derivatives can be stabilized by forming organometallic complexes. Rear-

angement of bis(trimethylsilyl)octa-3,5-diene-1,7-diyne with $\text{CpCo}(\text{CO})_2$ gave the sandwich compound 133 which, on treatment with $\text{Me}_4\text{N}^+\text{F}^-$, gave the parent sandwich compound 134.¹⁰⁸




The benzocyclobutadienes are paratropic 8π -electron systems and should add or lose 2π -electrons to become 10 or 6π -electron diatropic compounds. Dissolving the diol 135 in a mixture of SbF_5 and SO_2ClF at -78°C gave the cation 136,^{109,110} while treating diphenylbenzocyclobutene (137) with BuLi in TMEDA gave the dianion 138.¹¹¹ Whereas the dication appears to be a delocalized species, the dianion appears to have much of its charge localized in the substituents, presumably thus reducing electronic repulsion in the 4-membered ring.



Benzocyclobutenedione (12) and its derivatives have properties similar to dicarbonyl compounds. The 4-membered ring is, however, more readily cleaved because of the relief of ring strain that thus ensues. The carbonyl groups undergo nucleophilic addition and examples can be found in various sections of this review.

4 Benzodicyclobutadiene Derivatives


Benzodicyclobutadiene (139) has a 10π -electron periphery, but as with most polycyclic systems, this is probably not significant and the compound is best considered as an annelated benzocyclobutadiene where the annelation ensures that there is a cyclobutadiene ring in all canonical forms.



Benzodicyclobutadiene would thus be expected to be a reactive, unstable compound, and it has not been prepared. One derivative, 2,7-di-*tert*-butyl-4,5,9,10-tetraphenyltricyclo[6.2.0.0^{3,6}]deca-1,3(6),4,7,9-pentene (147) has been prepared^{113,112} as shown in Scheme 12.


Treatment of the alcohol 140, prepared from methyl pivalate and (phenylethynyl)magnesium bromide, with HBr in AcOH gave the bromoallene 141. Treatment of 141 with CuCl in DMF by a previously reported method¹¹³ gave *dl*-142 and *meso*-3,8-di-*tert*-butyl-

Scheme 12



1,5,6,10-tetraphenyldeca-3,4,6,7-tetraene-1,9-diyne (143) together with 3,6-di-*tert*-butyl-1,5,8-triphenyl-6-phenylethynyocta-3,4-diene-1,7-diyne (144), in yields of 11%, 11%, and 54%, respectively.¹¹² The same compounds could also be obtained by CuCl treatment of the acetate 148 or the Grignard reagent 149.¹¹² When 142 is heated in benzene it rearranges to the benzodicyclobutadiene 147 in 97% yield, probably by a disrotatory electrocycloization to 145 followed by a second electrocyclic ring closure and an allene dimerization, although these reactions do involve cyclobutadiene intermediates. Heating 143 in benzene provides support for this mechanism since the bis(methylene)-cyclobutene 146 is obtained by a disrotatory ring closure. Heating 146 in *o*-xylene then gives 147.

The benzodicyclobutadiene 147 was obtained as greenish blue needles which melt at 296–297 °C and shows absorption bands in the UV spectrum in CHCl_3 solution at 394 and 643 nm. The compound is air sensitive but thermally stable under N_2 , and no radical species is observed in the ESR spectrum.¹¹⁴ The mass spectrum shows a peak at m/e 271 corresponding to the dication M^{2+} in addition to a peak at m/e 542 for the molecular ion.¹¹² The formation of the dication probably represents the stability of the species with an electron removed from each 4-membered ring (151). Attempts to obtain an X-ray crystallographic structure were unsuccessful, the crystal decomposing. Since there are no ring protons, the ^1H NMR spectrum cannot be used to probe the ring current. The nature of 147 has thus to be determined largely from its chemical properties.



4.1 Reactions

4.1.1 Oxidation

Treatment of 147 with 2 mol equiv of $\text{Hg}(\text{OAc})_2$ together with 3 mol equiv of a nucleophile (NuH) in xylene at room temperature for 15–60 min gave 5,10-di-Nu-substituted 2,7-*tert*-butyl-4,5,9,10-tetraphenyltricyclo[6.2.0.0^{3,6}]deca-1,3,6,8-tetraene (155) (Scheme 13). The formation of 155 can be interpreted as a one-electron oxidation process. Electron transfer from 147 to $\text{Hg}(\text{II})$ gives the radical cation 150 which is then captured by the nucleophile to give the radical 152. Oxidation of this radical (as though through the

Scheme 13





Table 4. Reaction of 147 with Nucleophiles and Mercuric Acetate in Xylene at Room Temperature

nucleophile	reaction time, min	product	yield, %
MeOH	15	155a	68
EtOH	60	155b	31
PhOH	15	155c	46
PhNH ₂	15	155d	74
4-MeC ₆ H ₄ NH ₂	15	155e	63
AcOH	30	155f	55
PhCO ₂ H	60	155g	17

canonical hybriide 153) by a second one-electron process gives the cation 154 which is in its turn captured by the nucleophile to give 155. A considerable part of 155 formed is derived from the diradical dication 156, which itself reacts with the nucleophile through its dication form 157 to give 155. The yields and reaction times for a variety of nucleophiles are shown in Table 4.


Similar results to those tabulated were also obtained when a mercury(I) or silver(I) salt was used instead of the mercuric acetate. For example, reaction of 147 with methanol in the presence of Hg₂Cl₂ gave 155a in 63% yield, and with AgOAc 155a was obtained in 82% yield. When an equimolar amount of Hg(OAc)₂ rather than 2-fold excess was used, 155a and mercury were obtained in 68% and 82% yields, respectively.¹¹⁵

These reactions, which are effectively 1,8-additions, are paralleled by the addition of benzoyl peroxide and azobisisobutyronitrile, which give 158a and 158b, respectively.




As might be expected from the results with mercury and silver, compound 147 is readily oxidized by other reagents, the products depending on the reaction conditions. Stirring a suspension of 147 in MeOH under oxygen for 1 h at room temperature gave 161 in 96% yield.¹¹⁶ The same product was obtained with CCl₄, acetone, or petroleum ether as solvent. The oxidation probably proceeds by the sequence of reactions shown in Scheme 14. The intermediacy of the ring-cleaved dibenzoylbicyclobutadiene 159 in the sequence was supported by an alternative synthesis and its subsequent conversion to 161. Bis(methylene)cyclobutene 146 was oxidized by KMnO₄ to the octadienediyne 162, which was rearranged to 159 by heating to reflux in mesitylene. Compound 159 oxidizes immediately on exposure to air to give 161, but it can be trapped as its dibromo derivative 163 by treatment with bromine. Oxidation of 147 proceeds in the dark and triplet dioxygen must be the oxidizing agent.¹¹⁷⁻¹¹⁹

The oxirane 161 is stable in the crystalline state at room temperature, but in benzene solution oxygen is gradually absorbed and the isobenzofuran peroxide 164

Scheme 14

is obtained. Photooxidation of 161 in THF readily gives 164.¹¹⁶

Oxidation of 147 in the solid state proceeds by a different pathway to give the *o*-quinomethide derivative 165.¹¹⁶




Oxidation of either 147 or 165 with KMnO₄ gave 1,2,4,5-tetrabenzoyl-3,6-di-*tert*-butylbenzene (166).¹¹⁶

These results clearly indicate that 147 is not an aromatic 10 π -system, the ready cleavage of one of the 4-membered rings presumably occurring because a cyclobutadiene structure is thereby removed. The formation of 165 in the solid-state photooxidation may be due to a decrease in delocalization of the central benzene ring due to the annelating cyclobutadiene rings, thus allowing it to be oxidized.

4.1.2 Addition

Bromine adds readily to 147 in CHCl₃ to give the tetrabromide 167. Two equivalents of HBr add to 147 to give the dicyclobutabenzene derivative 169 in which the two bromine atoms are in the same 4-membered ring. Presumably an initial 1,8-addition of HBr occurs to give 168, which then adds the second molecule of HBr regiospecifically in a 1,6-manner. Debromination of 169 with zinc gives the benzocyclobutadiene derivative 170.¹¹⁷




4.1.3 Complexes with $\text{Fe}(\text{CO})_3$

Treatment of 147 with an equimolar amount of $\text{Fe}_2(\text{CO})_9$ in boiling xylene under N_2 for 4 h gave a mixture of the red crystalline 1:1 complex 171 and the black red crystalline 1:2 complex 172.^{118,119}

X-ray crystallographic analyses of the two complexes showed that 171 is best represented by the structure in which the iron atom is bonded to one of the 4-membered rings as a cyclobutadienyl complex, the 4-membered ring being square ($\text{C}-\text{C}$, 146.4 pm) with the iron atom centrally situated. The remaining bismethylenecyclobutene unit has alternating double and single bonds ($\text{C}-\text{C}$, 150.9 pm, $\text{C}=\text{C}$ 135.4 pm).^{118,119} In 172, both of the iron carbonyl groups are on the same side of the ring and are more strongly complexed to the nonbenzenoid double bond. The occurrence of both iron atoms on the same side may arise from simultaneous coordination of both iron atoms of a $\text{Fe}_2(\text{CO})_9$ molecule.

The distribution of the 10 π -electrons of the ring system in the complex 172 should be indicated by the X-ray crystallographic structure. If both of the 4-membered rings contributed 4 π -electrons to an iron atom, then two electrons remain and a σ -bond might be expected between the 2,7-positions. The X-ray crystallographic structure parameters do not indicate this to be the case (see Figure 4).



4.1.4 Reaction with Dienophiles and Dienes

Considering the possible canonical contributors to the structure of 147, there are a number of diene units which could be susceptible to reaction with dienophiles. In the "benzenoid" structure 147a, reaction across the 1,6-positions would give a product in which the cyclobutadiene character of both 4-membered rings is lost and there is little ring strain (pathway a, Figure 5). Reaction across the 3,4-positions in structure 147a (pathway b) and across the 3,6-positions in structure 147b (pathway d) would also remove the cyclobutadiene ring character of both 4-membered rings, but the products would be highly strained. Reaction across the 4,5-positions in 147b (pathway e) would give a benzocyclobutadiene structure and reaction across the




Figure 4. The X-ray crystallographic structure of the bis(iron tricarbonyl) complex 172.



Figure 5. Possible sites for dienophile addition to 147.

2,7-positions (pathway c) would give a product with two cyclobutadiene rings. Pathway a thus seems most probable, followed by pathway e, and pathway c seems the most unlikely.


Heating a 1:1 mixture of 147 and dimethyl fumarate (173a) in *o*-xylene under N_2 gave the 1,6-addition product 174a in 75% yield.¹²⁰ This reaction proceeds by pathway a, and the products from reaction with *trans*-dibenzoyl ethylene (173b) and *trans*-dicyanoethylene (174c) have analogous structures, 174b and 174c, respectively.^{120,121} As the simple consideration of cyclobutadiene structures suggested, pathway a is preferred in this system. Heating TCNE with 147 gives, however, mainly the product 175, together with a small amount of 176 (Scheme 15).¹²¹ Pathway e is now preferred over pathway a, presumably because steric repulsion between the CN and *tert*-butyl groups in the transition state of pathway a raises its energy above that of pathway e. An X-ray crystallographic study of the product 176 clearly indicates the serious steric crowding that occurs in this compound.¹²² A theoretical treatment of the reaction indicated that pathway a was energetically the most favorable.¹²² Calculations of the nonbonded interaction energies for various modes of approach of TCNE to 147 suggest that there is a folding of the 4-membered ring of 147 in the transition state.¹²³

Heating 175 converts it readily into 177, an analogous reaction to that which gives 132 (section 3.3).^{124,125} The X-ray structural analyses of 175¹²³ and 177¹²⁶ show the expected dominance of the dimethylenecyclobutene form as seen in other benzocyclobutadienes (section 3.2).


Interestingly, the Diels-Alder reaction between TCNE and benzocyclobutadienes in which the phenyl groups have been substituted for *p*-halogenophenyl groups, 178a and 178b, takes a different course, via pathway d, to give the propellanes 179a and 179b in 66% and 84% yield, respectively.¹²¹ The reason for this change in reaction pathway is not clear.

Heating 179a under reflux in *p*-xylene under N_2 for 30 min gave a mixture of 176a and 177a in 44% and


Scheme 15



40% yield, respectively.¹²¹ This process may occur by a retro-Diels-Alder reaction to give 178a and TCNE, followed by recombination by either pathway a or e. Some support for this mechanism is provided by heating 179a in the presence of an equimolar amount of 147 when 178a is formed in addition to 176 and 177.¹²¹




Diethyl acetylenedicarboxylate and dimethyl maleate both react with 147 by pathway b, the former dienophile giving the naphthocyclobutadiene 180.^{124,117}




The reason that dimethyl maleate takes a different reaction pathway compared to diethyl fumarate is

Scheme 16



probably the greater steric hindrance in pathway a in the maleate case.¹²⁰ Diphenylcyclopropene (181) also reacts with 147 via pathway b to give the quadricyclanone 183 via the intermediate 182.¹²⁷ Photoirradiation of 183 gave 184, and thermolysis gave 185. The former compound could be converted into 185 by thermolysis (Scheme 16).¹²⁷


With furan, 147 acted as a dienophile, reacting through one of the cyclobutene double bonds to give the benzocyclobutadiene derivative 185a.¹¹⁷



4.1.5 Addition of Dichlorocarbene

Treatment of 147 with CHCl₃ and KOH under phase-transfer conditions gave the propellane derivative 186 in 20% yield as the sole isolable product¹²⁸ (Scheme 17). Similar treatment of 178a and 178b gave the

Scheme 17




corresponding propellanes in 18% and 24% yield, respectively.¹²⁸ Although these are highly strained systems, all three propellanes were stable substances.

Hydrolysis of 186 by heating in acetone containing a small amount of aqueous NaOH for 2.5 h gave 147 in 72% yield. This reaction probably proceeds by hydrolysis of the *gem*-dichlorides to the ketone, followed by loss of CO.¹²⁸

Hydrolysis with AgClO₄ in wet THF initially gave a greenish-blue solution, the color then fades gradually to give finally a mixture of AgCl, silver, and 166. The color can be attributed to the 1:1 silver complex 188. In support of this concept, a benzene solution of equimolar amounts of AgClO₄ and 147 was the same color and had the same spectral properties (λ_{\max} 592 nm; ¹H NMR δ 1.09 (9 H, *t*-Bu), 1.24 (9 H, *t*-Bu), and decomposition of a dry benzene solution of 147 and excess AgClO₄ with water gave silver metal and 170 in 66% yield.

4.1.6 Insertion of Heteroatoms

Heteroatoms can be introduced between 4-C and 5-C of 147 to give tricyclic systems with 4-, 6-, and 5-membered rings. These compounds can be considered to be derived from the cyclobutenoheterocycles 189 and 190 (see section 3) by insertion of a benzene ring to give 191 and 192. Inspection of the structure of 192 suggests it should be more stable than 190, possessing a benzocyclobutadiene rather than a cyclobutadiene moiety, whereas 189 and 191 appear likely to have similar stabilities.




Heating 147 under reflux in DMSO under N₂ for 3 h gave the isobenzofuran derivative 194 in 54% yield. The reaction may proceed by the sequence shown (192a \rightarrow 193), addition of the DMSO followed by loss of dimethyl sulfide to form the oxirane and valence tautomerism to the furan (Scheme 18).¹²⁹

Heating 147 with tosylazide at 80 °C in *p*-xylene gave the pyrrole analogue 195 in 56% yield. The product from this latter reaction is in contrast with that from the addition of chlorocarbene and suggests that tosylnitrene is not the reactive species, probably reflecting the stability of tosyl azide.¹³⁰


Treatment of 147 with sulfur in boiling benzene under N₂ gave the thiophene analogue 196 in 43% yield.

In all of these cases the hetero atom has been introduced between the 4,5-positions of the benzocyclobutadiene. When 147 is heated with powdered selenium in boiling *p*-xylene under N₂, compound 197 is obtained in which the selenium atom has been inserted between the 3,4-positions. The derivative 197 was obtained in 89% yield as orange prisms. The reason for this change in regioselectivity may be steric, since the selenium atom is larger and in the selenium analogue of 196 the phenyl substituents on the heterocyclic ring will be crowded into the *tert*-butyl groups. In 197, there will be only one such interaction. Tellurium could not be introduced into 147, and this may reflect the greater steric interactions that would occur in both of the potential isomers. When 147 is heated with selenium dioxide in *p*-xylene then competition between insertion of oxygen and selenium occurs and a mixture of 194 (14%), 197 (32%), and 165 (22%) is formed.¹²⁹ The

Scheme 18





selenium derivative 197 is sensitive to oxygen, reacting even in the crystalline state, to give the dibenzoylselenophene 198 in quantitative yield.




When compounds 194 and 196 are treated with perchloric acid, one of the *tert*-butyl groups is lost and the perchlorate salt is formed. Thus 194 gives 199 which, on treatment with water, gives the furan 200. Similar treatment of 196 gave 201. Both 200 and 201 are stable compounds that can be reprotonated back to the corresponding salts. In the ¹H NMR spectrum, 200 shows signals at δ 0.85 (s, ¹Bu) and 6.50 (s, H²), together with signals for the phenyl groups, and 201 shows signals at δ 0.79 (s, ¹Bu) and 6.39 (s, H²). The chemical shift signals for H² can be compared to that of the similar proton in diphenylnaphthocyclobutadiene 122, H² δ 6.50.⁹² The chemical shifts of each these H² protons suggests that the 4-membered rings in these compounds sustain a paramagnetic ring current, but that the paratropicity is less than that found in the benzocyclobutadiene and related compounds (see Table 3, section 3.2). The ease of protonation of 200 and 201 may indicate that the systems should be considered as a 12 π -electron antiaromatic systems, the destabilization being lost on protonation. The nitrogen analogue 195 and the selenium derivative 197 did not form salts with perchloric acid. It is of interest that the H² protons in 200 and 201 are derived from the departing *tert*-butyl group and not the perchloric acid, as was shown by the nonincorporation of deuterium when DCIO₄ and D₂O were used.¹²⁹

Treatment of the selenophene 197 with Fe₂(CO)₉ gave the complex 171 and selenium in 85% and 98% yield, respectively.¹¹⁷ Treatment of 197 with TCNE gave the deselenated product 203, presumably via the Diels-Alder adduct 202 which then loses selenium. The highly conjugated nature of 203 is reflected in the purple color




of its crystals and the electronic spectrum [λ_{max} 260 nm (ϵ 35 400), 332 (36 500), 555 (12 300)].¹¹⁷




5 Biphenylene and Related Compounds

Biphenylene (4) was prepared by Lothrop¹³¹ in 1941 by an Ullman reaction of 2,2'-dibromo- (204a) or 2,2'-diiodobiphenyl (204b) and was, for many years, the only known derivative of cyclobutadiene fused to benzene.



There are five canonical uncharged structures for biphenylene, two of which contain a cyclobutadiene ring and three which do not; the X-ray crystallographic structure shows that biphenylene is best represented by the structure shown in Figure 6 (section 5.2).¹³² The bonds between the 6-membered rings approximate to single bonds in length and there is some bond alternation in the 6-membered rings, indicating that the canonical structure 4c containing the tetrakis(methylene)cyclobutane moiety is the major contributor.



A variety of compounds related to biphenylene have now been prepared and the formation and properties of a number of these will be described in this section.


5.1 Preparative Methods

The three main methods of forming biphenylene and its derivatives are as follows: (a) by coupling a biphenyl at the ortho positions of the two rings, (b) by dimerization of benzyne, and (c) by extrusion of nitrogen


or carbon dioxide and carbon monoxide from benzo-[c]cinnolines or phthalic anhydrides.

Method a is based on Lothrop's original synthesis and has been used to prepare a large variety of biphenylenes (see ref 133 and references therein). Copper powder appears preferable to the copper(I) oxide originally used as the cyclizing agent.¹³⁴


Method b involves the dimerization of benzyne and the yield of biphenylene depends on the method of generation of the benzyne. The oxidation of 1-aminobenzotriazoles (205) with lead tetraacetate produces high yields of biphenylenes,¹³⁵ but substituted benzotriazoles are rather inaccessible.



Method c utilizes flash vacuum pyrolysis and is only suitable for small amounts.¹³⁶⁻¹³⁹ It is a highly versatile method, however, and a variety of biphenylenes and heterobiphenylenes have been prepared by this route.




More recently, Vollhardt and his co-workers have introduced a method in which a 6-membered ring is constructed from a cobalt-catalyzed cyclization, the 4-membered ring arising because of the substrate chosen.¹⁴⁰ With bis(trimethylsilyl)acetylene (208) and *o*-diethynylbenzene (207), the 2,3-bis(trimethylsilyl)biphenylene (209a) is formed, which can be converted into biphenylene with trifluoroacetic acid.




Method a, method c and Vollhardt's method have also been used to prepare benzobiphenylenes and poly(biphenylenes) as described below.


Analogues of biphenylene in which one of the benzene ring has been replaced by another heterocyclic ring have also been prepared. Pyrolysis of the appropriate *N*-substituted benzocinnoline (e.g. 210) thus gives the pyridino biphenylene (e.g. 211).¹⁴¹⁻¹⁴⁶




For the replacement of a benzene ring by a 5-membered heterocycle, two methods have been used. 2-Thianorbiphenylene (213) was prepared by a Wittig reaction between 1,2-benzocyclobutadienedione (12) and the preformed bis-ylide 212.⁶⁴



1-Thianorindane (215) was prepared by thermal extrusion of nitrogen from 214 and is much less stable than 213, dimerizing to a number of products at 0 °C.¹⁴⁷




The oxirane 218 was prepared by oxidation of 216 with *m*-chloroperoxybenzoic acid to 217, which was then converted to 218 by bromination with NBS, followed by debromination with KO^tBu. Compound 218 shows no tendency to tautomerize to the oxepin.¹⁴⁸




Two isomeric thionin derivatives were prepared using the bis-Wittig approach described above for 2-thianorindane.¹⁴⁹

2,5-Dithiabispindane (219) and the tetraphenyl derivative (220) are the only known biphenylene derivatives in which both benzene rings are replaced by a 5-membered heterocycle,^{150a,b} both being obtained in very low yield.




A dibenzobiphenylene 221 (tetraazaphthylene) has been prepared with pyrazine rings in place of the benzene rings of the biphenylene moiety by two routes: condensation and polyphosphoric acid cyclization of squaric acid and *o*-diaminobenzene,¹⁵¹ and by method c.¹⁵²




A number of compounds have been prepared in which one of the benzene rings has been replaced by larger carbocyclic system (Scheme 19). Two tropone isomers were prepared by Wege and co-workers,¹⁵³ the first 223 by allylic oxidation of 222 followed by dehydrogenation with dichlorodicyanoquinone (DDQ), and the second 225 by bromination and debromination of 224. Compound 223 is much more stable than 225, which

Scheme 19



could only be trapped with cyclopentadiene. Three chloro derivatives of 223 have also been prepared by dichlorocarbene addition to 1- and 2-methoxybiphenylene, followed by hydrolysis.¹⁵⁴ The tropolones 226 and 227 have been prepared, and it was shown that the tautomers with the bismethylene structure were preferred to those with the benzocyclobutadiene structure (e.g. 226a).^{155,156}

Compounds with cyclooctatetraene 228 and methano-[10]annulene 229 rings were also prepared by the bis-Wittig procedure.^{157,158}




Biphenylene analogues have been prepared in which one of the benzene rings has been replaced by a charged $4n + 2$ π -electron system. The norbiphenylene anion 231 was prepared by a double Hoffman elimination from the 1,3-trimethyl ammonium salt 230 and is unstable, readily dimerizing.¹⁵⁹ The tropylium ion analogue, the homobiphenylene cation 230, has been prepared from the cycloheptatriene 232, obtained from the diene 232, by hydride removal with triphenylmethyl hexafluorophosphate and can be isolated as a salt.¹⁶⁰ A benzo analogue has also been prepared.¹⁶¹


Reduction of the cyclooctatetraene derivative 228 with alkali metals gave the corresponding dianion, equivalent to replacing one of the benzene rings by a 10 π -electron system.¹⁵⁷

The homobiphenylene anion (234), a biphenylene analogue with an 8 electron, $4n$ π -system replacing a benzene ring, has been prepared from 232 by treatment with potassium amide in liquid ammonia.¹⁶²

A large number of benzannulated biphenylene derivatives have been prepared by a variety of routes. Fusion can occur at either the a or b bond of the biphenylene to give linear or angular products, both



being known. Besides the application of a number of the methods described previously for biphenylenes and heterobiphenylenes, a method of considerable generality has been introduced by McOmie and co-workers.¹⁶³ A benzocyclobutaquinone is treated with *o*-bis(cyano-methyl)alkyne and a hindered base, DBU being preferred, in the presence of calcium hydride to remove the water formed in the reaction. An example is shown with the formation of the naphthalenobiphenylene 236 by reaction of 235 with 12.



Compounds have also been prepared in which a number of biphenylene units have been incorporated. Vollhardt's method has been particularly successful in this area, but the earlier methods have also been used, and Barton and co-workers¹⁶⁴ have adopted a Diels-Alder approach, using bis(methylene)cyclohexadienes as the diene component and benzocyclobutenes as the dienophile.

Again, angular and linear compounds can be constructed and examples of the various methods are given in Scheme 20.


5.2 Structure and Spectroscopic Properties

Biphenylene (4) is a stable, pale yellow crystalline solid. The X-ray crystallographic structure¹³² is shown in Figure 6.

The bonds between the phenylene rings are long. The X-ray crystallographic structures of a number of biphenylene derivatives and heterosubstituted biphenylenes have been obtained. These show similar bond lengths and angles to those of biphenylene itself. *N*-Ethyl-2,7-diazabiphenylene has long bonds (150 pm) between the two pyridinyl rings, the remaining bonds varying in length as in pyridine itself.¹⁶⁹ In angular [3]phenylene (6), the bonds between the phenylene rings are long (150.3 pm) and the central phenylene ring has alternating long and short bonds [134.5, 144.9 (annulated to 4-membered ring), 134.8, 144.6, 134.8, and 144.9.]¹⁶⁶ The outer benzenes also show alternation, with the bonds exocyclic to the 4-membered ring shorter, but this effect is much less pronounced, resembling the situation in biphenylene.

The X-ray crystallographic structure of hexakis(trimethylsilyl)tris(benzocyclobutadieno)benzene (251) is shown in Figure 7.¹⁶⁸ This clearly shows that the

Scheme 20



central ring is cyclohexatriene rather than benzene, the double and single bonds alternating with the bis-(methylene)cyclobutene structure dominant at each 4-membered ring.




Figure 6. X-ray crystallographic structure of biphenylene. Bond lengths are in pm, and bond angles are in degrees.




Figure 7. X-ray crystallography structure of hexakis(trimethylsilyl)tris(benzocyclobutadieno)benzene. Bond lengths are in pm, and bond angles are in degrees.


The ^1H NMR spectra of biphenylene shows the protons at a higher field than for simple benzenoid systems (H-1, δ 6.70; H-2, δ 6.60). The spectrum of 2-thianorbiphenylene (**213**) shows similar shifts (H-1, δ 6.49; H-4,5, δ 6.86), and this effect is presumed to arise from the paratropic effect of the antiaromatic 4-membered ring, since on oxidation of **213** to the sulfone, the phenyl signals move to lower field (H-4,5, δ 7.63), while the protons on the thiophene ring are virtually unaffected, indicating that the diatropic and paratropic contributions are of similar magnitude.¹⁰⁵ This effect is common in all of the biphenylene systems and becomes accentuated in annelated and polybiphenylenes.¹⁷⁰ In the polybiphenylenes, the protons on the inner benzene rings shift to still higher fields. For example, in the linear [3]phenylene **5**, the protons on the central ring resonate at δ 5.91, while in the angular system **6** they resonate at δ 6.13.¹⁷¹ Similar effects have been observed with systems containing a central naphthalene ring; in **244** the naphthalene protons resonate at δ 8.06, in **247** at δ 6.69, and in **248** at δ 6.42.¹⁶⁷ Thus, in these systems it also appears that bond localization occurs on annelation, structures with bismethylenecyclobutene moieties predominating.

In the homobiphenylene cation (**233**), the H-2,4 protons resonate at higher field than the remaining tropylium ring protons, suggesting that a greater amount of positive charge resides on these carbons, as expected for the bis(methylene)cyclobutene structure.¹⁶⁰ The benzannelated derivative shows similar shift differences, and the destabilizing effect of the benzocyclobutadiene annelation on the tropylium ion appears to be similar to that of annelation by benzene.¹⁰⁵

5.3 Reactions

Biphenylene undergoes classical electrophilic addition to give the 2-substituted derivatives. Under certain conditions, ring opening can occur to give benzocyclooctatetraene derivatives, possibly through addition followed by valence tautomerism of the tricyclo-

[6.4.0.0^{2,7}]dodecapentaene derivative.^{172,173} Thermolysis of biphenylene gives tetraphenylene, presumably via the 2,2'-biphenyl diradical.¹⁷⁴⁻¹⁷⁶ Photoirradiation of biphenylene gave the dimer **253**.¹⁷⁷




253

Reduction of biphenylene gives the monoanion and then the dianion, the latter a potential 14 π -electron system.¹⁷⁸⁻¹⁸¹ Oxidation of biphenylene gave the dication, the spectral properties of which indicate it is a delocalized 10 π -electron system.^{182,183}

The thiophene ring of 2-thianorbiphenylene (**213**) is more reactive than a normal thiophene, readily adding bromine.⁶⁴ On oxidation **213** forms the sulfoxide and sulfone, neither of which undergo Diels-Alder reactions.


Hydrogenation of the polybiphenylenes, which occurs under very mild conditions, provides considerable insight into the structure of those compounds. With linear [3]phenylene (**5**), the central ring is reduced to the cis-syn-cis isomer, and reduction of **252** occurs in a similar fashion with reduction of the central ring to give the cis-syn-cis-syn-cis hexahydro derivative.^{171,184} The ^1H NMR spectrum of the hexahydro derivative shows a singlet at δ 4.19 for the cyclohexane ring protons and the ^{13}C NMR spectrum for the compound shows only four absorption signals, substantiating the high symmetry.¹⁸⁴

The reactivity of the central ring of the tetrakis(trimethylsilyl) derivative of [3]phenylene is also illustrated by the reaction with butyllithium. Treatment with butyllithium followed by iodobutane gives the dialkylated compound **254**, presumably arising from nucleophilic addition of the butyl anion followed by trapping of the resultant anion with the iodobutane.¹⁷¹



254

Reaction with [3]phenylene (**5**) with SbF_5 in SO_2ClF gave the dication, and with potassium in THF the dianion was formed.¹⁷¹ The spectra of the dianion suggest that the two bis(methylene)benzocyclobutene structures (**255a** and **255b**) are the major canonical contributors.



255a

255b

6 Conclusions

Two factors restrict the formation of compounds containing bis(methylene)cyclobutene and tetrakis-

(methylene)cyclobutane structural groups: strain and the introduction of cyclobutadiene structures. The wealth of synthetic and structural information, and the extensive body of knowledge on the reactions of compounds containing these moieties, suggest that the first restriction is less important than the second in limiting the preparation of these compounds. Those compounds in which contributions from canonical structure containing cyclobutadiene systems are minimized are more stable, as is illustrated by the series, 3,4-bis(methylene)cyclobutene (1a), benzocyclobutadiene (2), and biphenylene (4), and is now supported by many of the findings described above, for example in the different properties of 1-thia- and 2-thianorbi-phenylene (215 and 213, respectively). Even where such structures are enforced, however, compounds can be prepared if access to reagents is prevented by suitable substitution. This is illustrated clearly in the formation of the benzodicyclobutadiene derivative 147 which, despite having one cyclobutadiene system in each of its canonical forms, is isolable. The preferences for structures without a cyclobutadiene contribution has been nicely demonstrated in the phenylenes, particularly by the work of Vollhardt and his collaborators. Phenyl rings are eventually persuaded to adopt cyclohexatriene structures as a means of limiting cyclobutadiene forms, as seen most spectacularly with tris(benzocyclobutadieno)benzene (252).

All of the compounds containing 4-membered rings described in this review do, however, involve degrees of bond angle strain and can be expected to undergo reactions in which this strain is lost or minimized. Such reactions are extensively illustrated in the sections above. Compounds containing bis(methylene)cyclobutene or tetrakis(methylene)cyclobutane moieties should thus continue to provide valuable reactive intermediates for the preparation of compounds with new, interesting structures.

References

- Cava, M. P.; Mitchell, M. J. *Cyclobutadiene and Related Compounds*; Academic Press: New York, 1967.
- Huntsman, W. D. In *The Chemistry of Ketenes, Allenes, and Related Compounds*; Patai, S., Ed.; John Wiley: Chichester, 1980; Part 2, pp 522-657.
- Lloyd, D. *Non-benzenoid Conjugated Carbocyclic Compounds*; Elsevier: Amsterdam, 1984; pp 431.
- Blomquist, A. T.; Maitlis, P. M. *Proc. Chem. Soc., London* **1961**, 332.
- Hopf, H. *Angew. Chem., Int. Ed. Engl.* **1970**, *9*, 732.
- Skattebol, L.; Solomon, S. J. *Am. Chem. Soc.* **1965**, *87*, 4506.
- Hopf, H.; Lenich, F. T. *Chem. Ber.* **1973**, *106*, 3461.
- Kleveland, K.; Skattebol, L. *J. Chem. Soc., Chem. Commun.* **1973**, 433.
- Huntsman, W. D.; Wristers, H. J. *J. Am. Chem. Soc.* **1963**, *85*, 3308.
- Huntsman, W. D.; Wristers, H. J. *J. Am. Chem. Soc.* **1967**, *89*, 342.
- Coller, B. A. W.; Heffernan, M. L.; Jones, A. J. *Aust. J. Chem.* **1968**, *21*, 1807.
- Toda, F.; Kumada, K.; Ishiguro, N.; Akagi, K. *Bull. Chem. Soc. Jpn.* **1970**, *43*, 3535.
- Toda, F.; Ohi, M. *J. Chem. Soc., Chem. Commun.* **1975**, 506.
- Pasto, D. J.; Yang, S. H. *J. Org. Chem.* **1989**, *54*, 3544.
- Pasto, D. J.; Kong, W. *J. Org. Chem.* **1989**, *54*, 4028.
- Henry, T. J.; Bergman, R. G. *J. Am. Chem. Soc.* **1972**, *94*, 5103.
- Pasto, D. J.; Mitra, D. K. *J. Org. Chem.* **1982**, *47*, 1381.
- Pasto, D. J.; Huang, N.-Z. *J. Org. Chem.* **1985**, *50*, 4465.
- Martin, H. D.; Kagabu, S.; Schwiwek, H. J. *Tetrahedron Lett.* **1975**, 3311.
- D'Amore, M. B.; Bergman, R. G.; Kent, M.; Hedeys, E. *J. Chem. Soc., Chem. Commun.* **1972**, 49.
- Eisenhuth, L.; Hopf, H. *J. Am. Chem. Soc.* **1974**, *96*, 5667.
- Eisenhuth, L.; Hopf, H. *Chem. Ber.* **1975**, *108*, 2635.
- Masamune, S.; Chin, C. G.; Hojo, K.; Seidner, R. T. *J. Am. Chem. Soc.* **1967**, *89*, 4804.
- Brown, R. D.; Godfrey, P. D.; Hart, B. T.; Ottrey, A. L.; Onda, M.; Woodruff, M. *Aust. J. Chem.* **1983**, *36*, 639.
- Skanche, A. *Acta Chem. Scand.* **1968**, *22*, 3239.
- Brown, R. D.; Burden, F. R.; Jones, A. J.; Kent, J. E. *J. Chem. Soc., Chem. Commun.* **1967**, 808.
- Griffith, D. W. T.; Kent, J. E.; O'Dwyer, M. F. *Aust. J. Chem.* **1972**, *25*, 241.
- Toda, F.; Ohi, N. *Bull. Chem. Soc. Jpn.* **1972**, *45*, 1820.
- Roth, W. R.; Lennartz, H. W.; Vogel, E.; Leiendecker, M.; Oda, M. *Chem. Ber.* **1986**, *119*, 837.
- Hess, B. A. J.; Schaad, L. J. *Aust. J. Chem.* **1972**, *25*, 2231.
- Blickle, P.; Hopf, H.; Bloch, M.; Jones, T. B. *Chem. Ber.* **1979**, *112*, 3691.
- Toda, F.; Akagi, K. *Tetrahedron* **1971**, *27*, 2801.
- Toda, F.; Higashi, M.; Akagi, K. *J. Chem. Soc., Chem. Commun.* **1969**, 1219.
- Toda, F.; Kumada, K.; Akagi, K. *Bull. Chem. Soc. Jpn.* **1970**, *43*, 2275.
- Toda, F.; Akagi, K. *Tetrahedron Lett.* **1970**, 5289.
- Schmidt, A. H.; Ried, W. *Synthesis* **1978**, 1.
- Hinschaw, J. C. *J. Chem. Soc. (D)* **1971**, 630.
- Scharf, H. D. *Angew. Chem., Int. Ed. Engl.* **1974**, *13*, 520.
- Smutny, E. J.; Caserio, M. C.; Roberts, J. D. *J. Am. Chem. Soc.* **1960**, *82*, 1793.
- Brune, H. A.; Horlbeck, G.; Röttele, H.; Tanger, Z. *Z. Naturforsch. B* **1973**, *28*, 68.
- Solomon, W. C.; Dee, L. A. *J. Org. Chem.* **1964**, *29*, 2790.
- Blomquist, A. T.; Nicholas, P. P. *J. Org. Chem.* **1967**, *32*, 863, 866.
- Park, J. D.; Snow, C. M.; Lacher, J. R. *J. Am. Chem. Soc.* **1951**, *73*, 2342.
- Ooms, P. H. J.; Scheeren, J. W.; Nivard, J. F. *Synthesis* **1975**, 260, 263, 639.
- Toda, F.; Ishihara, H.; Akagi, K. *Tetrahedron Letters* **1969**, 2531.
- Higashi, M.; Toda, F.; Akagi, K. *Chem. Ind. (London)* **1969**, 491.
- DeBoer, C. D. *J. Chem. Soc., Chem. Commun.* **1972**, 377.
- Breslow, R. *J. Am. Chem. Soc.* **1965**, *87*, 1320, 1326.
- Sizuki, Y.; Takizawa, T. *J. Chem. Soc., Chem. Commun.* **1972**, 837.
- Trahanovsky, W. S.; Park, M.-G. *J. Am. Chem. Soc.* **1973**, *95*, 5412.
- Trahanovsky, W. S.; Park, M.-G. *J. Org. Chem.* **1974**, *39*, 1448.
- Albert, B.; Heller, C.; Iden, R.; Martin, G.; Martin, H. D.; Mayer, B.; Oftring, A. *Isr. J. Chem.* **1985**, *25*, 74.
- Bilinski, V.; Dreiding, A. S.; Hollenstein, H. *Helv. Chim. Acta* **1983**, *66*, 2322.
- Ripoll, J. L.; Lasne, M. C. *Tetrahedron Lett.* **1978**, 5201.
- Lasne, M. C.; Ripoll, J. L.; Denis, J. M. *Tetrahedron* **1981**, *37*, 503.
- Taylor, G. A. *J. Chem. Soc., Chem. Commun.* **1968**, 1314.
- Taylor, G. A. *J. Chem. Soc. (C)* **1969**, 1755.
- Toda, F. *Chem. Lett.* **1972**, 621.
- Gatehouse, B. M. *Cryst. Struct. Commun.* **1982**, *11*, 369.
- Gatehouse, B. M. *Cryst. Struct. Commun.* **1982**, *11*, 365.
- Bergmann, E. D.; Agranat, I. *Tetrahedron* **1966**, *22*, 1275.
- Knorr, H.; Ried, W. *Synthesis* **1978**, 649.
- Knorr, U.; Knorr, H.; Ried, W.; Schuckmann, W. *Chem. Ber.* **1976**, *109*, 3869.
- Garratt, P. J.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1972**, *94*, 7087.
- Cava, M. P.; Firouzabadi, H.; Krieger, M. *J. Org. Chem.* **1974**, *39*, 480.
- Ried, W.; Knorr, H.; Knorr, U. *Chem. Ber.* **1976**, *109*, 1506.
- Ro, Z.; Tanaka, K.; Toda, F. *Bull. Chem. Soc. Jpn.* **1983**, *56*, 3193.
- Bird, C. W. *J. Chem. Soc., Chem. Commun.* **1968**, 1537.
- Toda, F.; Nakaoka, H.; Yuwane, m. K.; Todo, E. *Bull. Chem. Soc. Jpn.* **1973**, *46*, 1737.
- Ueda, K.; Toda, F. *Chem. Lett.* **1975**, 779.
- Ueda, K.; Igaki, M.; Toda, F. *Bull. Chem. Soc. Jpn.* **1976**, *49*, 3173.
- Nickels, H.; Dürr, H.; Toda, F. *Chem. Ber.* **1986**, *119*, 2249.
- Griffin, G. W.; Peterson, L. I. *J. Am. Chem. Soc.* **1962**, *84*, 3398.
- Griffin, G. W.; Peterson, L. I. *J. Am. Chem. Soc.* **1963**, *85*, 2268.
- Trabert, L.; Hopf, H. *Liebigs Ann. Chem.* **1980**, 1786.
- Trabert, L.; Hopf, H.; Schomburg, D. *Chem. Ber.* **1981**, *114*, 2405.
- Uhler, R. O.; Shechter, H.; Tiers, G. V. D. *J. Am. Chem. Soc.* **1962**, *84*, 3397.
- Tanaka, K.; Toda, F. *Tetrahedron Lett.* **1980**, *21*, 2713.
- Iyoda, M.; Otani, H.; Oda, M.; Kai, Y.; Baba, Y.; Kasai, N. *J. Am. Chem. Soc.* **1986**, *108*, 5371.
- Berkovitch-Yellin, Z.; Lahav, M.; Leiserowitz, L. J. *J. Am. Chem. Soc.* **1974**, *96*, 918.
- Uhler, R. O. *Dissertation Abstr.* **1960**, *21*, 765.
- Miller, F. A.; Brown, F. R.; Rhee, K. H. *Spectrochim. Acta (A)* **1972**, *28*, 1467.
- Bally, T.; Buser, U.; Haselbach, E. *Helv. Chim. Acta* **1978**, *61*, 38.
- Hart, H.; Ward, D. L.; K, T.; Toda, F. *Tetrahedron Lett.* **1982**, *23*, 2125.
- Emerson, G. F.; Watts, L.; Pettit, R. *J. Am. Chem. Soc.* **1965**, *87*, 131.
- Chapman, O. L.; Chang, C. C.; Rosenquist, N. R. *J. Am. Chem. Soc.* **1976**, *98*, 261.
- Koenig, T.; Imre, D.; Hoobler, J. A. *J. Am. Chem. Soc.* **1979**, *101*, 6446.
- Trahanovsky, W. S.; Fischer, D. R. *J. Am. Chem. Soc.* **1990**, *112*, 4971.

- (89) Mitchell, G. H.; Sondheimer, F. *J. Am. Chem. Soc.* **1969**, *91*, 7520.
(90) Vollhardt, K. P. C.; Yee, L. S. *J. Am. Chem. Soc.* **1977**, *99*, 2010.
(91) Straub, H. *Liebig's Ann. Chem.* **1978**, 1675.
(92) Cava, M. P.; Hwang, B.; van Meter, J. P. *J. Am. Chem. Soc.* **1963**, *85*, 4032.
(93) Cava, M. P.; Hsu, A. F. C. *J. Am. Chem. Soc.* **1972**, *94*, 6441.
(94) Vollhardt, K. P. C.; Bergman, R. G. *J. Am. Chem. Soc.* **1972**, *94*, 8950.
(95) Bergman, R. G.; Vollhardt, K. P. C. *J. Chem. Soc., Chem. Commun.* **1973**, 214.
(96) Vollhardt, K. P. C.; Bergman, R. G. *J. Am. Chem. Soc.* **1973**, *95*, 7538.
(97) Garratt, P. J.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1972**, *94*, 1022.
(98) Cava, M. P.; Napier, D. R. *J. Am. Chem. Soc.* **1957**, *79*, 3606.
(99) Cava, M. P.; Napier, D. R.; Pohl, R. J. *J. Am. Chem. Soc.* **1963**, *85*, 2076.
(100) Turney, T. A. In *Chemistry of Quinonoid Compounds*; Patai, S., Ed.; John Wiley: New York, 1974; Vol. 2, pp 857.
(101) Schmidt, A. H.; Ried, W. *Synthesis* **1978**, 869.
(102) Wong, H. N. C.; Chan, T.-L.; Luh, T.-Y. In *Chemistry of Quinonoid Compounds*, 2nd ed.; Patai, S., Rappaport, Z., Ed.; John Wiley: New York, 1988; Vol. 2.
(103) South, M. S.; Liebeskind, L. S. *J. Org. Chem.* **1982**, *47*, 3815.
(104) McOmie, J. F. W.; Perry, D. H. *J. Chem. Soc., Chem. Commun.* **1973**, 248.
(105) Garratt, P. J. *Pure Appl. Chem.* **1975**, *44*, 783.
(106) Müller, E.; Fettel, H.; Sauerbier, M. *Synthesis* **1970**, 82.
(107) Straub, H.; Miller, B. Z. *Naturforsch. B* **1980**, *35 B*, 1484.
(108) Duclos, R. I.; Vollhardt, K. P. C.; Yee, L. S. *J. Organomet. Chem.* **1979**, *174*, 1979.
(109) Olah, G. A.; Liang, G. *J. Am. Chem. Soc.* **1976**, *98*, 3033.
(110) Olah, G. A.; Liang, G. *J. Am. Chem. Soc.* **1977**, *99*, 6045.
(111) Boche, G.; Etzrodt, H.; Marsch, M.; Thiel, W. *Angew. Chem., Int. Ed. Engl.* **1982**, *21*, 132.
(112) Toda, F.; Takahara, Y. *Bull. Chem. Soc. Jpn.* **1976**, *49*, 2515.
(113) Toda, F.; Takehira, Y. *J. Chem. Soc., Chem. Commun.* **1975**, 174.
(114) Toda, F.; Mukai, K. *Chem. Lett.* **1975**, 777.
(115) Takehira, Y.; Tanaka, K.; Toda, F. *Chem. Lett.* **1976**, 1323.
(116) Toda, F.; Dann, N.; Tanaka, K.; Takehira, Y. *J. Am. Chem. Soc.* **1977**, *99*, 4529.
(117) Toda, F.; Tanaka, K.; Matsui, M. *Tetrahedron Letters* **1982**, *23*, 217.
(118) Butters, T.; Toda, F.; Winters, W. *Angew. Chem., Int. Ed. Engl.* **1980**, *19*, 926.
(119) Butters, T.; Winter, W.; Toda, F. *Acta Crystallogr.* **1981**, *B37*, 1532.
(120) Toda, F.; Tanaka, K. *J. Chem. Soc., Chem. Commun.* **1976**, 177.
(121) Toda, F.; Yoshioka, T. *Chem. Lett.* **1977**, 561.
(122) Tsukada, H.; Shimanouchi, H.; Sasada, Y. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 983.
(123) Tsukada, H.; Shimanouchi, H.; Sasada, Y. *Bull. Chem. Soc. Jpn.* **1978**, *51*, 985.
(124) Toda, F.; Dann, N. *J. Chem. Soc., Chem. Commun.* **1976**, 30.
(125) Toda, F.; Tanaka, K.; Matsui, M. *Tetrahedron Lett.* **1982**, *23*, 217.
(126) Tsukada, H.; Shimanouchi, Y.; Sasada, Y. *Acta Crystallogr.* **1977**, *B 33*, 2951.
(127) Toda, F.; Tanaka, K. *J. Chem. Soc., Chem. Commun.* **1976**, 1010.
(128) Toda, F.; Yoshioka, T.; Kawada, M. *Chem. Lett.* **1978**, 1451.
(129) Toda, F.; Tanaka, K. *Chem. Lett.* **1979**, 1451.
(130) Breslow, D. S.; Sloan, M. F.; Newburg, N. R.; Renfrow, W. B. *J. Am. Chem. Soc.* **1969**, *91*, 2273.
(131) Lothrop, W. C. *J. Am. Chem. Soc.* **1941**, *63*, 1187.
(132) Fawcett, J. K.; Trotter, J. *Acta Crystallogr.* **1966**, *20*, 87.
(133) Constantine, P. R.; Hall, G. E.; Harrison, C. R.; McOmie, J. F. W.; Searle, R. J. G. *J. Chem. Soc. (C)* **1966**, 922.
(134) Salfield, J. C.; Baume, E. *Tetrahedron Lett.* **1966**, 3365.
(135) Cambell, C. D.; Rees, C. W. *J. Chem. Soc. (C)* **1969**, 742.
(136) Kanokranaporn, S.; MacBride, J. A. H. *J. Chem. Res.* **1980**, (S) 203; (M) 2901.
(137) Cava, M. P.; Mitchell, M. J.; De Jongh, D. C.; Van Fossen, R. Y. *Tetrahedron Letters* **1966**, 2947.
(138) Brown, R. F. C.; Gardner, D. V.; McOmie, J. F. W.; Solly, R. K. *Aust. J. Chem.* **1967**, *20*, 139.
(139) Gardner, D. V.; McOmie, J. F. W.; Albrigtsen, P.; Harris, R. K. *J. Chem. Soc. (C)* **1969**, 1994.
(140) Berris, B. C.; Lai, V.-H.; Vollhardt, K. P. C. *J. Chem. Soc., Chem. Commun.* **1982**, 953.
(141) Barton, J. W.; Walker, R. B. *Tetrahedron Lett.* **1975**, 569.
(142) Kramer, J.; Berry, R. S. *J. Am. Chem. Soc.* **1972**, *94*, 8336.
(143) Hägele, G.; Sartori, P.; Golloch, A. Z. *Naturforsch. B* **1973**, *28*, 758.
(144) MacBride, J. A. H. *J. Chem. Soc., Chem. Commun.* **1974**, 359.
(145) MacBride, J. A. H.; Wright, P. M.; Wakefield, B. J. *Tetrahedron Lett.* **1981**, 4545.
(146) Kanokranaporn, S.; MacBride, J. A. H.; King, T. J. *J. Chem. Res.* **1980**, (S) 204; (M) 2911.
(147) Barton, J. W.; Lapham, D. J. *Tetrahedron Lett.* **1979**, 3571.
(148) Jikeli, G.; Günther, H. *Angew. Chem., Int. Ed. Engl.* **1974**, *13*, 277.
(149) Garratt, P. J.; Holmes, A. B.; Sondheimer, F.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1970**, *92*, 4492.
(150) (a) Shepherd, M. K. *J. Chem. Soc., Chem. Commun.* **1985**, 880. (b) Garratt, P. J.; Neoh, S. B. *J. Org. Chem.* **1975**, *40*, 970.
(151) Hunig, S.; Pütter, H. *Chem. Ber.* **1977**, *110*, 2532.
(152) Kanokranaporn, S.; MacBride, J. A. H. *J. Chem. Res.* **1980**, (S) 206; (M) 2941.
(153) Lombardo, L.; McCulloch, R. K.; Wege, D. *Aust. J. Chem.* **1978**, *31*, 1585.
(154) Sato, M.; Ebine, S.; Tsunetsugu, J. *J. Chem. Soc., Perkin Trans. 1* **1977**, 1282.
(155) Sato, M.; Inaba, K.; Ebine, S.; Tsunetsuga, J. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 2334.
(156) Sato, M.; Ebine, S.; Nishijima, K.; Tsunetsuga, J. *Bull. Chem. Soc. Jpn.* **1981**, *54*, 766.
(157) Baxter, C. S.; Garratt, P. J.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1969**, *91*, 7783.
(158) Garratt, P. J.; Vollhardt, K. P. C. *Angew. Chem., Int. Ed. Engl.* **1971**, *10*, 125.
(159) Cava, M. P.; Narasimhan, K.; Zeiger, W.; Radonovich, L. J.; Glick, M. D. *J. Am. Chem. Soc.* **1969**, *91*, 2378.
(160) Lombardo, L.; Wege, D. *Aust. J. Chem.* **1978**, *31*, 1569.
(161) Garratt, P. J.; Vollhardt, K. P. C. *J. Chem. Soc., Chem. Commun.* **1971**, 1143.
(162) Staley, S. W.; Heinrich, F.; Orvedal, J. *J. Am. Chem. Soc.* **1976**, *98*, 2681.
(163) Buckland, P. R.; Hacker, N. P.; McOmie, J. F. W. *J. Chem. Soc., Perkin Trans. 1* **1983**, 1443.
(164) Barton, J. W.; Shepherd, M. K.; Willis, R. J. *J. Chem. Soc., Perkin Trans. 1* **1986**, 967.
(165) Barton, J. W.; Rowe, D. J. *Tetrahedron Lett.* **1983**, *24*, 299.
(166) Diercks, R.; Vollhardt, K. P. C. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 266.
(167) Helson, H. E.; Vollhardt, K. P. C.; Yang, Z. Y. *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 114.
(168) Diercks, R.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1986**, *108*, 3150.
(169) MacBride, J. A. H. *J. Chem. Soc., Chem. Commun.* **1972**, 1219.
(170) Barton, J. W.; Rowe, D. J. *Tetrahedron* **1985**, *41*, 1323.
(171) Berris, B. C.; Hovakeemian, G. H.; Lai, Y. H.; Meadagh, H.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1985**, *107*, 5670.
(172) Barton, J. W.; Whitaker, K. E. *J. Chem. Soc. (C)* **1968**, 1663.
(173) Boulton, A. J.; Middleton, D. *J. Org. Chem.* **1974**, *39*, 2956.
(174) Friedman, L.; Lindow, D. F. *J. Am. Chem. Soc.* **1968**, *90*, 234.
(175) Friedman, L.; Rabideau, P. W. *J. Org. Chem.* **1968**, *33*, 451.
(176) Gaidis, J. M. *J. Org. Chem.* **1970**, *35*, 2811.
(177) Goldman, N. L.; Ruden, R. A. *Tetrahedron Lett.* **1968**, 3951.
(178) Hush, N. S.; Rowlands, J. R. *Mol. Phys.* **1963**, *6*, 317.
(179) Bauld, N. L.; Banks, D. *J. Am. Chem. Soc.* **1965**, *87*, 128.
(180) Waack, R.; Doran, M. A.; West, P. *J. Am. Chem. Soc.* **1965**, *87*, 5508.
(181) Cox, R. H.; Terry, H. W.; Harrison, L. W. *Tetrahedron Lett.* **1971**, 4815.
(182) Ronlán, A.; Parker, V. D. *J. Chem. Soc., Chem. Commun.* **1974**, 33.
(183) Olah, G. A.; Liang, G. *J. Am. Chem. Soc.* **1977**, *99*, 6045.
(184) Mohler, D. L.; Vollhardt, K. P. C.; Wolff, S. *Angew. Chem., Int. Ed. Engl.* **1990**, *29*, 1151.