

Positional Cloning of the Mouse Saccharin Preference (*Sac*) Locus

Alexander A. Bachmanov^{1*}, Xia Li^{1*}, Danielle R. Reed^{1*}, Jeffery D. Ohmen^{2*}, Shanru Li¹, Zhenyu Chen¹, Michael G. Tordoff¹, Pieter J. de Jong^{2,3}, Chenyan Wu², David B. West², Alu Chatterjee², David A. Ross² and Gary K. Beauchamp^{1,4}

¹Monell Chemical Senses Center, Philadelphia, PA, ²Pfizer Global Research and Development, Alameda, CA, ³Children's Hospital Oakland Research Institute, Oakland, CA and ⁴University of Pennsylvania, Philadelphia, PA, USA

Correspondence to be sent to: Gary K. Beauchamp, Monell Chemical Senses Center, 3500 Market St, Philadelphia, PA 19104, USA.
e-mail: beauchamp@monell.org

*These authors contributed equally to this article.

Abstract

Differences in sweetener intake among inbred strains of mice are partially determined by allelic variation of the saccharin preference (*Sac*) locus. Genetic and physical mapping limited a critical genomic interval containing *Sac* to a 194 kb DNA fragment. Sequencing and annotation of this region identified a gene (*Tas1r3*) encoding the third member of the T1R family of putative taste receptors, T1R3. Introgression by serial backcrossing of the 194 kb chromosomal fragment containing the *Tas1r3* allele from the high-sweetener-preferring C57BL/6ByJ strain onto the genetic background of the low-sweetener-preferring 129P3/J strain rescued its low-sweetener-preference phenotype. Polymorphisms of *Tas1r3* that are likely to have functional significance were identified using analysis of genomic sequences and sweetener-preference phenotypes of genealogically distant mouse strains. *Tas1r3* has two common haplotypes, consisting of six single nucleotide polymorphisms: one haplotype was found in mouse strains with elevated sweetener preference and the other in strains relatively indifferent to sweeteners. This study provides compelling evidence that *Tas1r3* is equivalent to the *Sac* locus and that the T1R3 receptor responds to sweeteners.

Introduction

Sweet taste transduction is thought to be initiated by the interaction of a sweetener ligand with a G-protein-coupled taste receptor on apical ends of the taste receptor cells, which evokes a sensation of pleasantness and a consummatory behavioral response (Lindemann, 1996). Many compounds that taste sweet to humans (sweeteners) are palatable to various species of animals, including mice (Beauchamp and Mason, 1991; Bachmanov *et al.*, 2001). Inbred mouse strains display marked differences in their avidity for sweet solutions (Lush, 1989; Capeless and Whitney, 1995; Lush *et al.*, 1995). Much of the difference in sweetener preferences among mouse strains is attributed to allelic variation of the saccharin preference (*Sac*) locus, on distal chromosome 4 (Phillips *et al.*, 1994; Lush *et al.*, 1995; Bachmanov *et al.*, 1997b; Blizard *et al.*, 1999; Li *et al.*, 2001b). In addition to sweetener preferences, the *Sac* genotype influences the afferent responses of gustatory nerves to sweeteners (Bachmanov *et al.*, 1997b; Li *et al.*, 2001b), suggesting that the *Sac* gene is involved in peripheral taste transduction and may encode a sweet-taste receptor.

The T1R family of putative taste receptors consists of

three genes expressed in taste receptor cells and located on the distal chromosome 4 (Hoon *et al.*, 1999; Kitagawa *et al.*, 2001; Max *et al.*, 2001; Montmayeur *et al.*, 2001; Sainz *et al.*, 2001), which makes them candidates for the *Sac* locus. The *Tas1r1* (alias *Gpr70*) gene encoding the T1R1 receptor and the *Tas1r2* (*Gpr71*, T1R2) gene have been excluded as candidates for *Sac* based on their more proximal chromosomal location (Kitagawa *et al.*, 2001; Li *et al.*, 2001b; Montmayeur *et al.*, 2001). However, the *Tas1r3* gene encoding the T1R3 receptor has been mapped to a more distal part of chromosome 4 corresponding to the *Sac* interval (Kitagawa *et al.*, 2001; Max *et al.*, 2001; Montmayeur *et al.*, 2001; Sainz *et al.*, 2001).

In this study, using high-resolution linkage analysis followed by physical mapping and sequencing, we determined the exact limits of the critical *Sac* region. Next, we identified genes within the *Sac* interval and found that *Tas1r3* is the most likely candidate for the *Sac* locus. The low-sweetener-preferring phenotype of 129P3/J strain was rescued by introgressing an allele of *Tas1r3* from a high-sweetener-preferring C57BL/6ByJ strain using serial back-

crossing during selection of a 129.B6-*Sac* congenic strain. Finally, sequence variants of *Tas1r3* that are likely to have functional significance were identified using analysis of *Tas1r3* sequences and sweetener-preference phenotypes in genealogically distant mouse strains. These *in-vivo* data provide compelling evidence that *Tas1r3* is equivalent to the *Sac* locus and that it encodes a taste receptor responding to sweeteners.

Materials and methods

Animals

C57BL/6ByJ (B6) and 129P3/J (formerly 129/J, abbreviated here as 129) mice were purchased from The Jackson Laboratory. Details of breeding F₂ hybrids (Bachmanov *et al.*, 1997b) and of marker-assisted selection of a 129.B6-*Sac* congenic strain (Li *et al.*, 2001b) are described elsewhere.

Taste preference tests

Consumption of 120 mM sucrose and 17 mM saccharin was measured in individually caged mice using 96 h, two-bottle tests, with water as the second choice, as described in detail elsewhere (Bachmanov *et al.*, 1997b; Li *et al.*, 2001b). Results are presented as sweetener solution intakes expressed per 30 g of body wt (the approximate weight of an adult mouse) per day, or as a preference score (ratio of solution intake to total fluid intake, in %). Indexes of sweetener consumption used for interval mapping were standardized within each gender and experimental group relative to the group mean and standard deviation.

Mouse genotyping and linkage analysis

Mouse genomic DNA was purified from tails by NaOH/Tris (Truett *et al.*, 2000) or phenol/chloroform extraction. Simple sequence length polymorphism (SSLP) markers were tested using a standard protocol (Dietrich *et al.*, 1992), with minor modifications (Bachmanov *et al.*, 1997b; Li *et al.*, 2001b). All other markers were tested using a single-strand conformation polymorphism (SSCP) protocol (Orita *et al.*, 1989) or by sequencing the polymerase chain reaction (PCR) products. Marker and primer details are available upon request. Polymorphic markers were mapped by genotyping F₂ and partially congenic mice. Chromosomal positions of non-polymorphic markers were confirmed using radiation hybrid mapping of the T31 mouse–hamster radiation hybrid panel (Research Genetics, Huntsville, AL) according to a standard protocol (<http://www.jax.org/resources>) and the data were submitted for analysis to the Mouse Genome Database. Markers that mapped in the *Sac* region were used to screen the RPCI-23 mouse bacterial artificial chromosome (BAC) library (Osoegawa *et al.*, 2000).

Construction of a BAC contig

The RPCI-23 female (C57BL/6J) mouse BAC library

(Osoegawa *et al.*, 2000) was screened by hybridization (Church and Gilbert, 1984) with radioactively labeled probes (Feinberg and Vogelstein, 1983). The library was screened twice, first with a probe generated from the yeast artificial chromosome (YAC) clone 178B3 (http://carbon.wi.mit.edu:8000/cgi-bin/mouse/sts_info) and, second, with pooled probes of markers in the *Sac* region. The YAC 178B3 was chosen for screening because, based on its STS content, it mapped in the *Sac* region and appeared to be non-chimeric. Positive clones identified by the initial screenings were re-arrayed and hybridized against individual probes. The secondary screening results were confirmed by PCR. BAC insert sizes were determined using pulsed-field gel electrophoresis after digestion with *NotI* (Lengeling *et al.*, 1999). The BAC contig was assembled using SEGMAP, version 3.48 (Green and Green, 1991).

Analysis of nucleic and amino acid sequences

DNA from BAC 118E21 was isolated using an alkaline lysis protocol followed by CsCl density gradient centrifugation, subcloned and sequenced. Sequence data were assembled using Phred, Phrap and Consed packages of programs. After BAC 118E21 was sequenced, the sequence-tagged site (STS) content of this BAC and overlapping BACs was confirmed by aligning the STS and BAC end sequences with the 118E21 sequence (Sequencher, Gene Codes Corporation, Ann Arbor, MI). Repeat sequences were identified using RepeatMasker (Smit and Green, 2000) and GENSCAN (Burge and Karlin, 1998) was used to predict gene content and exon–intron organization. The predicted proteins were submitted to TBLASTN searches against the *nr* and the mouse *EST* databases at NCBI or against Unigene.

To determine the intron–exon boundaries of the *Tas1r3* gene, total RNA was extracted using TRIzol Reagent (Life Technologies Inc., Rockville, MD) from enzymatically separated mouse lingual epithelium (Ruiz *et al.*, 1995; Spielman and Brand, 1995), which included fungiform, foliate and circumvallate taste papillae. The RNA was reverse transcribed (Superscript reverse transcriptase, Life Technologies). The cDNA samples were amplified using AmpliTaq DNA Polymerase with GeneAmp (Perkin Elmer Corp., Branchburg, NJ) and intron-spanning primers selected to distinguish genomic and cDNA. Single bands of expected sizes were excised from the gel, purified and sequenced.

Mouse T1R3 protein sequence was predicted from cDNA obtained from tongue epithelium by RT-PCR, and it was analyzed using the computer programs HMMTOP (Tusnády and Simon, 1998) and TOPO (Johns and Speth, 1996) to determine the hydrophobicity and the transmembrane regions. Protein motif prediction was conducted using the Motif search service on GenomeNet (<http://www.motif.genome.ad.jp>).

Analysis of variation in sweetener preference and *Tas1r3* sequences among mouse strains

Sweetener preference data were taken from previous studies for the following mouse strains: 129/Rr, 129/Sv, AKR/J, BALB/cA, BALB/cByJ, C3H/He, C57BL/6ByJ, C57BL/6Ty, C57L/Lac, CBA/Cam, DBA/2Ty, IS/Cam, SEA/GnJ, ST/bJ, SWR/J (Lush, 1989) and CAST/Ei (A. Bachmanov *et al.*, unpublished data). When preferences were available for two substrains, they were averaged and shown as 129, BALB/c and C57BL/6.

A 6.8 kb segment of genomic DNA, including ~2.6 kb upstream and ~1.2 kb downstream of *Tas1r3*, was sequenced in genealogically remote or unrelated (Beck *et al.*, 2000) mouse strains with high (C57BL/6ByJ, SWR/J and CAST/Ei) or low (129P3/J, AKR/J and DBA/2J) sweetener preferences. Comparison of the sequences from these six strains identified a haplotype of six single nucleotide polymorphisms (SNPs) associated with sweetener preference. Next, the regions contributing to this haplotype were sequenced in additional mouse strains (BALB/cByJ, C3H/HeJ, C57L/J, CBA/J, IS/CamEi, SEA/GnJ and ST/bJ). The C57BL/6J strain sequence obtained from the sequencing of RPCI-23 BAC 118E21 was identical to the C57BL/6ByJ sequence. In some cases, the exact substrain tested for sweetener preference differed from the substrain sequenced, usually because the older substrains were not available for sequencing.

Results

Linkage mapping

The initial linkage analysis was conducted using an F₂ intercross between B6 mice with high sweetener acceptance and 129 mice with the low sweetener acceptance. The F₂ mice were phenotyped using 96 h two-bottle tests with sucrose and saccharin, and genotyped with markers polymorphic between the B6 and 129 strains. Interval mapping narrowed the region containing *Sac* to ~5 cM (Figure 1a,b). This region was further reduced to 0.7 cM (flanked by *280G12-T7* and *D4Mon1* markers) during the marker-assisted selection of a 129.B6-*Sac* segregating partially congenic strain (Figure 1b,c).

Physical mapping

A contig of BAC clones representing the *Sac*-containing region was constructed by screening the RPCI-23 mouse BAC library (Figure 1d). A 246 kb BAC clone, 118E21, including most of the *Sac* interval, was sequenced. The remaining small proximal part of the *Sac* region was contained in a sequence from mouse genomic DNA (GenBank accession No. AF185591). The 0.7 cM *Sac* interval flanked by *280G12-T7* and *D4Mon1* had a physical size of 194 kb and contained twelve predicted genes (Figure 1e).

Identification of genes within the *Sac* interval

Of the twelve predicted genes, four were known (cyclin ania-6b, *Dvl*, *Ubc6p* and *Tas1r3*), two were similar to known human genes (*KIAA1716* and *SCNN1D*) and six were represented as cDNA clones with GenBank accession Nos NM_025338, AK004732, NM_024472, AK010425, AA435261 and NM_026125. Most of the genes identified within the *Sac*-containing interval are involved or potentially involved in cell division and differentiation [cyclin ania-6b and *Dvl* (MacLachlan *et al.*, 1995; Lee *et al.*, 1999)], maintenance of intracellular processes [*Ubc6p*, NM_024472 and genes similar to *KIAA1716* and *SCNN1D* (Waldmann *et al.*, 1995; Gilon *et al.*, 2000; Nagase *et al.*, 2000)], or collagen synthesis (NM_026125). The functions of the predicted genes represented by NM_025338, AK004732, AK010425 and AA435261 are unknown.

Of these twelve genes within the 194 kb *Sac* interval, only one, *Tas1r3* (taste receptor, type 1, member 3), was a G-protein-coupled receptor. A predicted *Tas1r3* protein, T1R3, has moderate sequence homology to putative G-protein-coupled taste receptors T1R1 (encoded by *Tas1r1*, alias *Gpr70*, 32%; Figure 2a), T1R2 (encoded by *Tas1r2*, alias *Gpr71*, 30%) and mGluR4 (encoded by *Gpr1d*, 23%) (Hoon *et al.*, 1999; Chaudhari *et al.*, 2000). Sequence comparison of cDNA from mouse lingual epithelium and genomic DNA showed that *Tas1r3* contains six coding exons (Figure 2b). It is translated into an 858-amino-acid protein with a predicted secondary structure that includes seven transmembrane domains and a large hydrophilic extracellular N-terminus (Figure 2c). This structure is typical of the G-protein-coupled receptor family 3, which includes the metabotropic glutamate and extracellular calcium-sensing receptors. There is substantial evidence that a G-protein-coupled mechanism is involved in sweet taste transduction (Lindemann, 1996), suggesting that *Tas1r3* is the most likely candidate for the *Sac* locus.

Sequence variants of *Tas1r3*

As a candidate for *Sac*, *Tas1r3* should have sequence variants corresponding to phenotypical *Sac* alleles. To assess this correspondence, sequences of *Tas1r3* and surrounding genomic DNA were analysed in mouse strains with known sweetener preferences (see details in Materials and methods). Two haplotypes consisting of six SNPs (Figure 2b) distinguished strains with high and low sweetener preferences (Figure 3). Two of these SNPs resulted in amino acid substitutions of threonine (found in all high-preferring strains) for alanine (found in all low-preferring strains) at position 55 (Thr55Ala) and isoleucine for threonine at position 60 (Ile60Thr; Figure 2a), both within the predicted extracellular N-terminal domain of T1R3 (Figure 2c). Two other variants occurred in the upstream regions, the fifth polymorphism was intronic and the sixth variant was a silent mutation (Ser) in exon 1.




Figure 1 Genetic and physical maps of the *Sac* region. **(a)** Interval mapping of sucrose and saccharin consumption to distal chromosome 4 using MAPMAKER software (Lander et al., 1987). Distances between markers were estimated based on data from the B6 × 129 F₂ intercross ($n = 629$). Curves trace the likelihood of odds (LOD) scores calculated using an unconstrained model [LOD threshold for significant linkage 4.3, 2 d.f. (Lander and Kruglyak, 1995)]. The horizontal lines show the 2-LOD drop confidence intervals for saccharin (dotted line, 5.3 cM) and sucrose (solid line, 4.5 cM); black triangles indicate the respective LOD score peaks (LOD 20.3 for saccharin and 23.3 for sucrose). This locus explained 18.6 and 16.2% of the variance in saccharin and sucrose intakes, respectively. Analyses of LOD scores under dominant and additive models (not shown) demonstrated that the B6 allele is dominant over the 129 allele. Analysis of preference scores showed similar results (not shown). **(b)** Average daily 17 mM saccharin consumption by mice from parental 129 and B6 strains (left), F₂ hybrids (center) and N₆, N₇, N₄F₄ and N₃F₅ segregating partially congenic 129.B6-Sac mice (right) in 96 h, two-bottle tests with water (means ± SE). Genotypes of the F₂ and congenic mice for *Tas1r3* and their numbers are indicated on the bars. Each group had approximately equal numbers of males and females. Differences between parental strains and among the F₂ and congenic genotypes were significant ($F > 39.5$, $P < 0.000001$, ANOVA). Females consumed more saccharin than males ($F > 26.5$, $P < 0.000005$) and the differences among genotypes were more pronounced in females than in males (interaction gender × strain or genotype, $F > 6.4$, $P < 0.02$). However, the main effect of genotype was the same for females and males: F₂ and congenic B6 homozygotes and heterozygotes for *Tas1r3* did not differ from each other and had higher saccharin intakes than did 129 homozygotes ($P < 0.000001$, planned comparisons). Intakes of 120 mM sucrose were 14.2 ± 0.6 ml/30 g body wt for the F₂ mice homozygous for B6 allele of *Tas1r3* ($n = 170$), 13.8 ± 0.5 ml/30 g body wt for the F₂ heterozygotes ($n = 299$) and 7.4 ± 0.4 ml/30 g body wt for the F₂ mice homozygous for 129 allele of *Tas1r3* ($n = 152$); results of statistical analyses were similar to those for saccharin. The haplotype of the donor fragment in the *Sac*-congenic mice is depicted in panel (c). **(c)** Linkage map of the *Sac*-containing region. Distances between markers were obtained from the B6 × 129 F₂ intercross [see panel (a)]. A black box depicts the donor fragment of the 129.B6-Sac partially congenic mice whose saccharin intakes are shown on panel (b), right. The location and the size of the donor fragment were determined based on the presence of B6 alleles of polymorphic markers in mice from the N₄, N₆, N₇, N₄F₄ and N₃F₅ generations. The donor fragment ends proximally between *280G12-T7* and *49O2-T7*, and distally between *350D2-T7* and *D4Mon1*. **(d)** BAC contig and physical map of distal chromosome 4 in the *Sac* region. BAC sizes (kb) are shown in parentheses. Dots indicate presence of markers within BACs detected by hybridization and confirmed by PCR and, in some cases, by sequencing. *139J18-SP6* and *D4Mon1* were found within the 118E21 sequence, but were not used for BAC screening (*D4Mon1* was used for genotyping F₂ and congenic mice). In this region, linkage distance of 1 cM corresponds to ~0.25 Mb of physical distance instead of typical 2 Mb and, therefore, the frequency of recombinations is ~8 times higher than the average throughout the genome. **(e)** Genes within the *Sac*-containing interval flanked by *280G12-T7* and *D4Mon1*. The full sequence of this region is assembled from BAC 118E21 and from *Mus musculus* cyclin ania-6b gene (GenBank accession No. AF185591). Filled areas indicate predicted genes. Arrows indicate the predicted direction of transcription.




a**b****c**

Figure 2 Structure of the *Tas1r3* gene. **(a)** Protein alignment of the novel T1R3 and the previously described T1R1 [*Tas1r1* or *Gpr70*; GenBank accession No. AF301161 (Li *et al.*, 2001b)] mouse genes. Identical amino acids are shaded in black; conservative amino acid substitutions are shaded in gray. The protein sequences of T1R3 and T1R1 were deduced from cDNA sequences of the B6 strain. Asterisks denote missense polymorphisms in T1R3. Roman numerals with solid black bars beneath indicate the transmembrane domains. **(b)** Structure of the *Tas1r3* gene based on comparison between genomic DNA and cDNA sequences from mouse tongue epithelium. The six coding exons are shown as black boxes. Exon and intron sizes are given in nucleotide base pairs (sizes of exons I and VI are partial, excluding untranslated regions). Asterisks indicate SNPs defining haplotypes of high-sweetener-preferring strains (B6-like) and low-sweetener-preferring strains (129-like; see Figure 3). The B6-like/129-like haplotypes are as follows (the haplotype nucleotides are numbered with the A in the ATG start codon as nucleotide 1): T/A at nt –2383; A/G at nt –183; A/G at nt 135 (exon I, silent at amino acid position 45); A/G at nt 163 (exon I, missense T55A); T/C at nt 179 (exon I, missense I60T); and T/C at nt 651, (intron II). **(c)** Conformation of the predicted T1R3 protein. The missense mutations (Thr55Ala and Ile60Thr) are denoted with asterisks.




Figure 3 Relationship between *Tas1r3* haplotype and sweetener preference. **(a)** Saccharin and **(b)** sucrose preferences by mice from inbred strains with two different haplotypes of the *Tas1r3* gene (see Figure 2b). The strains with the B6-like haplotype of *Tas1r3* strongly preferred saccharin ($81 \pm 4\%$) and sucrose ($86 \pm 5\%$), whereas strains with the 129-like haplotype were indifferent to these solutions (57 ± 1 and $57 \pm 3\%$, respectively, $P_s < 0.0003$, t -tests). The SEA/GnJ strain had lower sweetener preference compared with other strains with the B6-like haplotype. A mutation within the bone morphogenetic 5 protein in the SEA/GnJ strain (Kingsley et al., 1992) may cause disturbance in calcium metabolism and reduce sweetener preference (Tordoff and Rabusa, 1998). The complete strain name is shown if identical substrains were used for genotyping and phenotyping; the strain name is truncated if the substrain genotyped differed from the substrain phenotyped.

Discussion

Using a positional cloning approach, we narrowed the *Sac*-containing region to a 194 kb interval. One gene within this interval, *Tas1r3*, encodes a G-protein-coupled receptor (T1R3) that is expressed in taste receptor cells (Kitagawa et al., 2001; Max et al., 2001; Montmayeur et al., 2001; Sainz et al., 2001). Based on the effects of the *Sac* genotype on peripheral sweet-taste responsiveness (Bachmanov et al., 1997a,b; Li et al., 2001b) and on the known mechanism of sweet-taste transduction (Lindemann, 1996), *Tas1r3* is the most likely candidate for *Sac*.

If *Tas1r3* is identical to *Sac*, substitution of *Tas1r3* alleles must result in phenotypical changes attributed to the *Sac* locus. This can be tested through transgenic ('phenotype rescue' or 'knock-out') experiments. We obtained equivalent data using the 129.B6-*Sac* congenic mice. Introgression of the 194 kb chromosomal fragment containing the *Tas1r3* allele from the high-sweetener-preferring B6 strain onto the genetic background of the 129 strain fully rescued its low sweetener preference phenotype: sweetener intake of the congenic mice was as high as that of mice from the donor B6 strain. This introgression was essentially equivalent to production of BAC-transgenic mice (Antoch et al., 1997; Probst et al., 1998) because the size of the donor chromo-

somal fragment was comparable to the size of a typical BAC. These data demonstrate that substitution of *Tas1r3* alleles results in behavioral changes attributed to the *Sac* locus and therefore provide the strongest evidence to date that *Tas1r3* is identical to *Sac* and that the T1R3 receptor responds to sweeteners.

Further evidence that *Tas1r3* is equivalent to *Sac* was obtained from correspondence of *Tas1r3* sequence variants to sweetener-preference phenotypes in inbred mouse strains, which are largely attributed to allelic variation of the *Sac* locus. *Tas1r3* has multiple SNPs among the inbred strains we examined. We reasoned that variants that sort mouse strains into high- and low-sweetener preferring categories are more likely to be responsible for the *Sac* phenotype. To minimize the chances for sharing of chromosomal fragments due to identity-by-descent, we analysed mouse strains with unrelated or distant genealogies. Using this approach, we identified a haplotype of six SNPs concordant with sweetener preferences. Two of these SNPs resulted in amino acid substitutions at positions 55 (Thr55Ala) and 60 (Ile60Thr) within the predicted extracellular N-terminus domain of T1R3. The same Thr55Ala and Ile60Thr polymorphisms have been identified by others (Kitagawa et al., 2001; Max et al., 2001; Montmayeur et al., 2001; Sainz et al., 2001) and they could affect ligand binding (Rana et al., 2001) or receptor dimerization (Max et al., 2001). The SNPs in non-coding upstream and intronic regions may affect expression of the gene (Phillips, 1999; Rana et al., 2001).

What are the ligands of T1R3? The B6 and 129 strains differ in responses to several different classes of sweeteners, and to ethanol, which has a sweet-taste component (Bachmanov et al., 1996, 2001). Our genetic analysis of the B6 $\times$ 129 F₂ hybrids demonstrated that the *Sac* genotype affects behavioral and neural responses to representatives of several different sweet-tasting chemicals (sugars, non-caloric sweeteners, sweet amino acids) and also ethanol consumption (Bachmanov et al., 1997a; Tordoff et al., 1998). The effects of *Sac* genotype on taste responses to a variety of compounds suggest that they evoke sweet-taste sensation via a common receptor, T1R3.

It is controversial as to whether single or multiple sweet-taste receptors and transduction pathways exist (Bartoshuk, 1987; Bernhardt et al., 1996; Lindemann, 1996; DuBois, 1997). Our data are consistent with the existence of a single sweet-taste receptor. But it is also possible that the involvement of T1R3 in responses to diverse compounds is attributable to its dimerization with other members of the T1R family (Max et al., 2001; Montmayeur et al., 2001) or to the existence of T1R3 splice variants (Max et al., 2001) (X. Li and D. Reed, unpublished data), which may create multiple forms of sweet-taste receptors. However, the most abundant naturally occurring sweeteners, sugars, are chemically similar and may not require multiple receptors for their recognition. This may explain why the family of

putative sweet (T1R) receptors is much smaller than the family of putative bitter (T2R) receptors (Adler *et al.*, 2000).

Striking species differences in behavioral and physiological responses to sweeteners imply that the structure of sweet receptors varies among species (Beauchamp and Mason, 1991). A taste receptor for the sugar trehalose (Ishimoto *et al.*, 2000), identified in *Drosophila melanogaster*, has no homology with T1R3, suggesting the independent origin of insect and mammalian sugar taste receptors. Among mammals, cats, which are strict carnivores, apparently do not prefer carbohydrate sweeteners, but do prefer certain amino acids humans describe as sweet (Beauchamp *et al.*, 1977). Even primate species differ in the ability to recognize certain high-potency sweeteners as sweet (Glaser *et al.*, 1995). *TAS1R3*, the human ortholog of mouse *Tas1r3*, exists in a region of conserved synteny, 1p36. Preliminary sequencing of *TAS1R3* detected common SNPs (D. Reed *et al.*, unpublished data), which may be involved in genetic variation of the sensory and behavioral responses to sugars in humans (Reed *et al.*, 1997).

Finally, an understanding of sweet receptors may have practical applications. Most high-potency, low-calorie sweeteners were discovered by the accidental tasting of chemicals synthesized for other purposes (Walters, 1991). Models guided by the *Tas1r3* taste receptor structure should allow the design and development of new potent ligands (DuBois, 1997; Nofre, 2001).

To summarize, we have now defined the limits of the *Sac*-containing critical genomic interval, restricted to the 194 kb DNA fragment, and have identified 12 genes within this region. Although all these genes can be considered candidates for *Sac*, based on gene expression and predicted protein structures, *Tas1r3* is the most likely candidate. We have further demonstrated that experimental manipulation of the genomic interval containing *Tas1r3* affects the taste responses to sweeteners attributed to the *Sac* locus, which provides novel *in-vivo* evidence for the identity of *Tas1r3* and *Sac*. Finally, we have conducted the first quantitative analysis of the relationship between the *Tas1r3* sequence variants and sweetener-preference phenotypes in genealogically diverse mouse strains. In addition to *Tas1r3* missense polymorphisms reported by others, we have identified three other polymorphisms in non-coding regions that may affect *Tas1r3* function. Together, these data provide strong evidence that *Tas1r3* is identical to *Sac* and that it is associated with the taste quality of sweetness.

Acknowledgements

The data described in this paper were presented at the XXIII Annual Meeting of the Association for Chemoreception Sciences, Sarasota, FL, USA, 25–29 April 2001 (Li *et al.*, 2001a). Sequence data reported in this paper have been deposited in GenBank and assigned the accession No. AF311386. We thank Maja Bućan for guidance with physical mapping and comments on the manuscript, Tim Wiltshire for help with BAC library screening, R. Arlen Price

for initial genotyping of the F₂ hybrids and Linda M. Bartoshuk, Taufiqul Huque, John Teeter, Charles Wysocki, Nancy Rawson and Joseph Brand for providing advice and discussion. We also thank Lianchun Chen, Dulce San Juan, Joe Catanese and Al Cairo for help with physical mapping. The excellent technical support of Sarah Tobias and the Pfizer Global Research and Development–Alameda Genotyping and Sequencing Core is acknowledged. This work was supported by National Institutes of Health grants R01DC00882 (G.K.B.), R03DC03509, R01DC04188 and R01-DK55853 (D.R.R.), R01AA11028 (M.G.T.) and R03DC03853 (A.A.B.), and by a special grant from the Ambrose Monell Foundation.

References

- Adler, E., Hoon, M.A., Mueller, K.L., Chandrashekar, J., Ryba, N.J. and Zuker, C.S. (2000) A novel family of mammalian taste receptors. *Cell*, 100, 693–702.
- Antoch, M.P., Song, E.J., Chang, A.M., Vitaterna, M.H., Zhao, Y., Wilsbacher, L.D., Sangoram, A.M., King, D.P., Pinto, L.H. and Takahashi, J.S. (1997) Functional identification of the mouse circadian Clock gene by transgenic BAC rescue. *Cell*, 89, 655–667.
- Bachmanov, A.A., Tordoff, M.G. and Beauchamp, G.K. (1996) Ethanol consumption and taste preferences in C57BL/6ByJ and 129/J mice. *Alc. Clin. Exp. Res.*, 20, 201–206.
- Bachmanov, A.A., Reed, D.R., Ninomiya, Y., Inoue, M., Tordoff, M.G., Price, R.A. and Beauchamp, G.K. (1997a) Genetic locus on mouse chromosome 4 influencing taste responses to sweeteners. *Chem. Senses*, 22, 642 (Abstract).
- Bachmanov, A.A., Reed, D.R., Ninomiya, Y., Inoue, M., Tordoff, M.G., Price, R.A. and Beauchamp, G.K. (1997b) Sucrose consumption in mice: major influence of two genetic loci affecting peripheral sensory responses. *Mamm. Genome*, 8, 545–548.
- Bachmanov, A.A., Tordoff, M.G. and Beauchamp, G.K. (2001) Sweetener preference of C57BL/6ByJ and 129P3/J mice. *Chem. Senses*, BA13300.
- Bartoshuk, L.M. (1987) Is sweetness unitary? An evaluation of the evidence for multiple sweets. In Dobbing, J. (ed.), *Sweetness*. Springer, London, pp. 33–46.
- Beauchamp, G.K. and Mason, J.R. (1991) Comparative hedonics of taste. In Bolles, R.C. (ed.), *The Hedonics of Taste*. Lawrence Erlbaum, Hillsdale, NJ, pp. 159–183.
- Beauchamp, G.K., Maller, O. and Rogers, J. (1977) Flavor preferences in cats (*Felis catus* and *Panthera sp.*). *J. Comp. Physiol. A*, 91, 1118–1127.
- Beck, J.A., Lloyd, S., Hafezparast, M., Lennon-Pierce, M., Eppig, J.T., Festing, M.F. and Fisher, E.M. (2000) Genealogies of mouse inbred strains. *Nature Genet.*, 24, 23–25.
- Bernhardt, S.J., Naim, M., Zehavi, U. and Lindemann, B. (1996) Changes in IP₃ and cytosolic Ca²⁺ in response to sugars and non-sugar sweeteners in transduction of sweet taste in the rat. *J. Physiol.*, 490, 325–336.
- Blizard, D.A., Kotlus, B. and Frank, M.E. (1999) Quantitative trait loci associated with short-term intake of sucrose, saccharin and quinine solutions in laboratory mice. *Chem. Senses*, 24, 373–385.
- Burge, C.B. and Karlin, S. (1998) Finding the genes in genomic DNA. *Curr. Opin. Struct. Biol.*, 8, 346–354.
- Capeless, C.G. and Whitney, G. (1995) The genetic basis of preference for sweet substances among inbred strains of mice: preference ratio

- phenotypes and the alleles of the *Sac* and *dpa* loci. *Chem. Senses*, 20, 291–298.
- Chaudhari, N., Landin, A.M. and Roper, S.D. (2000) A metabotropic glutamate receptor variant functions as a taste receptor. *Nature Neurosci.*, 3, 113–119.
- Church, G. and Gilbert, W. (1984) *Genomic sequencing*. Proc. Natl Acad. Sci. USA, 81, 1991–1995.
- Dietrich, W., Katz, H., Lincoln, S.E., Shin, H.-S., Friedman, J., Dracopoli, N. and Lander, E.S. (1992) A genetic map of the mouse suitable for typing intraspecific crosses. *Genetics*, 131, 423–447.
- DuBois, G.E. (1997) New insights on the coding of the sweet taste message in chemical structure. In Salvadori, G. (ed.), *Olfaction and Taste: A Century of the Senses*. Allured, Carol Stream, IL, pp. 32–95.
- Feinberg, A.P. and Vogelstein, B. (1983) A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.*, 132, 6–13.
- Gilon, T., Chomsky, O. and Kulka, R.G. (2000) Degradation signals recognized by the *Ubc6p-Ubc7p* ubiquitin-conjugating enzyme pair. *Mol. Cell. Biol.*, 20, 7214–7219.
- Glaser, D., Tinti, J.M. and Nofre, C. (1995) Evolution of the sweetness receptor in primates. I. Why does alitame taste sweet in all prosimians and simians, and aspartame only in Old World simians? *Chem. Senses*, 20, 573–584.
- Green, E.D. and Green, P. (1991) Sequence-tagged site (STS) content mapping of human chromosomes: theoretical considerations and early experiences. *PCR Meth. Applic.*, 1, 77–90.
- Hoon, M.A., Adler, E., Lindemeier, J., Battey, J.F., Ryba, N.J. and Zuker, C.S. (1999) Putative mammalian taste receptors: a class of taste-specific GPCRs with distinct topographic selectivity. *Cell*, 96, 541–551.
- Ishimoto, H., Matsumoto, A. and Tanimura, T. (2000) Molecular identification of a taste receptor gene for trehalose in *Drosophila*. *Science*, 289, 116–119.
- Johns, S.J. and Speth, R.C. (1996) TOPO, transmembrane protein display software. <http://www.sacs.ucsf.edu/TOPO/topo.html>
- Kingsley, D.M., Bland, A.E., Grubber, J.M., Marker, P.C., Russell, L.B., Copeland, N.G. and Jenkins, N.A. (1992) The mouse short ear skeletal morphogenesis locus is associated with defects in a bone morphogenetic member of the TGF beta superfamily. *Cell*, 71, 399–410.
- Kitagawa, M., Kusakabe, Y., Miura, H., Ninomiya, Y. and Hino, A. (2001) Molecular genetic identification of a candidate receptor gene for sweet taste. *Biochem. Biophys. Res. Commun.*, 283, 236–242.
- Lander, E. and Kruglyak, L. (1995) Genetic dissection of complex traits: guidelines for interpreting and reporting linkage results. *Nature Genet.*, 11, 241–247.
- Lander, E., Green, P., Abrahamson, J., Barlow, A., Daley, M., Lincoln, S. and Newburg, L. (1987) MAPMAKER: an interactive complex package for constructing primary linkage maps of experimental and natural populations. *Genomics*, 1, 174–181.
- Lee, J.S., Ishimoto, A. and Yanagawa, S. (1999) Characterization of mouse dishevelled (*Dvl*) proteins in Wnt/Wingless signaling pathway. *J. Biol. Chem.*, 274, 21464–21470.
- Lengeling, A., Wiltshire, T., Otmani, C. and Bućan, M. (1999) A sequence-ready BAC contig of the GABA_A receptor gene cluster *Gabrg1-Gabra2-Gabrb1* on mouse chromosome 5. *Genome Res.*, 9, 732–738.
- Li, X., Bachmanov, A.A., Reed, D.R., Li, S., Chen, Z., Tordoff, M.G., Beauchamp, G.K., de Jong, P.J., Wu, C., West, D.B., Chatterjee, A., Ross, D.A. and Ohmen, J.D. (2001a) Molecular origins of the sweet tooth: a novel taste receptor controlling the avidity for sucrose and saccharin in mice (Abstract). 23rd AChemS Annual Meeting, p. 28.
- Li, X., Inoue, M., Reed, D.R., Huque, T., Puchalski, R.B., Tordoff, M.G., Ninomiya, Y., Beauchamp, G.K. and Bachmanov, A.A. (2001b) High-resolution genetic mapping of the saccharin preference locus (*Sac*) and the putative sweet taste receptor (*T1R1*) gene (*Gpr70*) to mouse distal Chromosome 4. *Mamm. Genome*, 12, 13–16.
- Lindemann, B. (1996) Taste reception. *Physiol. Rev.*, 76, 718–766.
- Lush, I.E. (1989) The genetics of tasting in mice. VI. Saccharin, acesulfame, dulcin and sucrose. *Genet. Res.*, 53, 95–99.
- Lush, I.E., Hornigold, N., King, P. and Stoye, J.P. (1995) The genetics of tasting in mice. VII. Glycine revisited, and the chromosomal location of *Sac* and *Soa*. *Genet. Res.*, 66, 167–174.
- MacLachlan, T.K., Sang, N. and Giordano, A. (1995) Cyclins, cyclin-dependent kinases and cdk inhibitors: implications in cell cycle control and cancer. *Crit. Rev. Euk. Gene Expr.*, 5, 127–156.
- Max, M., Shanker, Y.G., Huang, L., Rong, M., Liu, Z., Campagne, F., Weinstein, H., Damak, S. and Margolskee, R.F. (2001) *Tas1r3*, encoding a new candidate taste receptor, is allelic to the sweet responsiveness locus *Sac*. *Nat. Genet.*, 28, 58–63.
- Montmayeur, J.P., Liberles, S.D., Matsunami, H. and Buck, L.B. (2001) A candidate taste receptor gene near a sweet taste locus. *Nat. Neurosci.*, 4, 492–498.
- Nagase, T., Kikuno, R., Hattori, A., Kondo, Y., Okumura, K. and Ohara, O. (2000) Prediction of the coding sequences of unidentified human genes. XIX. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro. *DNA Res.*, 7, 347–355.
- Nofre, C. (2001) New hypotheses for the GPCR 3D arrangement based on a molecular model of the human sweet-taste receptor. *Eur. J. Med. Chem.*, 36, 101–108.
- Orita, M., Iwahana, H., Kanazawa, H., Hayashi, K. and Sekiya, T. (1989) Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphisms. *Proc. Natl Acad. Sci. USA*, 86, 2766–2770.
- Osoegawa, K., Tateno, M., Woon, P.Y., Frengen, E., Mammoser, A.G., Catanese, J.J., Hayashizaki, Y. and de Jong, P.J. (2000) Bacterial artificial chromosome libraries for mouse sequencing and functional analysis. *Genome Res.*, 10, 116–128.
- Phillips, P.C. (1999) From complex traits to complex alleles. *Trends Genet.*, 15, 6–8.
- Phillips, T.J., Crabbe, J.C., Metten, P. and Belknap, J.K. (1994) Localization of genes affecting alcohol drinking in mice. *Alc. Clin. Exp. Res.*, 18, 931–941.
- Probst, F.J., Fridell, R.A., Raphael, Y., Saunders, T.L., Wang, A., Liang, Y., Morell, R.J., Touchman, J.W., Lyons, R.H., Noben-Trauth, K., Friedman, T.B. and Camper, S.A. (1998) Correction of deafness in shaker-2 mice by an unconventional myosin in a BAC transgene. *Science*, 280, 1444–1447.
- Rana, B.K., Shiina, T. and Insel, P.A. (2001) Genetic variations and polymorphisms of G protein-coupled receptors: functional and therapeutic implications. *Annu. Rev. Pharmacol. Toxicol.*, 41, 593–624.
- Reed, D.R., Bachmanov, A.A., Beauchamp, G.K., Tordoff, M.G. and Price, R.A. (1997) Heritable variation in food preferences and their contribution to obesity. *Behav. Genet.*, 27, 373–387.
- Ruiz, C., McPheeters, M. and Kinnamon, S. (1995) Tissue culture of rat

- taste buds. In Spielman, A. and Brand, J. (eds), *Experimental and Cell Biology of Taste and Olfaction: Current Techniques and Protocols*. CRC Press, Boca Raton, FL, pp. 79–84.
- Sainz, E., Korley, J.N., Battey, J.F. and Sullivan, S.L. (2001) *Identification of a novel member of the T1R family of putative taste receptors*. *J. Neurochem.*, 77, 896–903.
- Smit, F. and Green, P. (2000) *RepeatMasker*. <http://ftp.genome.washington.edu/cgi-bin/RepeatMasker>
- Spielman, A. and Brand, J. (1995) *Collection of taste tissue from mammals*. In Spielman, A. and Brand, J.G. (eds), *Experimental Cell Biology of Taste and Olfaction: Current Techniques and Protocols*. CRC Press, Boca Raton, FL, pp. 25–32.
- Tordoff, M.G. and Rabusa, S.H. (1998) *Calcium-deprived rats avoid sweet compounds*. *J. Nutr.*, 128, 1232–1238.
- Tordoff, M.G., Bachmanov, A.A., Reed, D.R., Price, R.A. and Beauchamp, G.K. (1998) *Influence of saccharin preference (Sac) locus on ethanol preference in mice*. *Alc. Clin. Exp. Res.*, 22, 101A (Abstract).
- Truett, G.E., Heeger, P., Mynatt, R.L., Truett, A.A., Walker, J.A. and Warman, M.L. (2000) *Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris (HotSHOT)*. *Biotechniques*, 29, 52–54.
- Tusnády, G. and Simon, I. (1998) *Principles governing amino acid composition of integral membrane proteins: applications to topology prediction*. *J. Mol. Biol.*, 283, 489–506.
- Waldmann, R., Champigny, G., Bassilana, F., Voilley, N. and Lazdunski, M. (1995) *Molecular cloning and functional expression of a novel amiloride-sensitive Na⁺ channel*. *J. Biol. Chem.*, 270, 27411–27414.
- Walters, E.D. (1991) *The rational discovery of sweeteners*. In Walters, D.E., Orthoefer, F.T. and DuBois, G.E. (eds), *Sweeteners. Discovery, Molecular Design, and Chemoreception*. American Chemical Society, Washington DC, pp. 1–11.

Accepted June 18, 2001

Note added in proof

While this paper was in press, a new study [Nelson, G., Hoon, M.A., Chandrashekar, J., Zhang, Y., Ryba, N.J.P. and Zuker, C.S. (2001) *Mammalian sweet taste receptors*. *Cell*, 106, 381–390] has confirmed the identity of the *Sac* locus and *Tas1r3* gene, and provided strong evidence that the T1R3 protein functions as a sweet taste receptor.