

CORRIGENDUM

Parasitic Wasps Learn and Report Diverse Chemicals with Unique Conditionable Behaviors

D.M. Olson, G.C. Rains, T. Meiners, K. Takasu, M. Tertuliano, J.H. Tumlinson, F.L. Wäckers and W.J. Lewis

Chemical Senses, 28, 545–549

Regrettably, an error occurred in Figure 3 of this article. The x-axis in part B should have read '2,4-Dini' instead of '3-Oct'.

The corrected figure is printed below.




Figure 3 Seek and coil behaviors of female *Microplitis croceipes* conditioned with food and hosts and (A) 3-octanone and (B) 2,4-dinitrotoluene. Seek and coil responses to 3-octanone when trained with food and 3-octanone ($\chi^2_{df1} = 26.67$, $P = 0.000$, $n = 40$), hosts and 3-octanone ($\chi^2_{df1} = 15.82$, $P = 0.000$, $n = 40$) and hosts only ($\chi^2_{df1} = 1.03$, $P = 0.311$, $n = 40$). Seek and coil responses to 2,4-DNT when trained with food and 2,4-DNT ($\chi^2_{df1} = 22.57$, $P = 0.000$, $n = 40$), food only ($\chi^2_{df1} = 1.03$, $P = 0.311$, $n = 40$), hosts and 2,4-DNT ($\chi^2_{df1} = 15.17$, $P = 0.000$, $n = 40$) and hosts only ($\chi^2_{df1} = 2.11$, $P = 0.147$, $n = 40$).