


II. Brom-Bestimmungen.

Substanz	Ein- waage mg	Ver- brauch ccm $\frac{1}{100}$ -n. Lauge	Gef. % Br	Ber. % Br	Kontrollbestimm. mit Oxycyanid. Verbrauch ccm $\frac{1}{100}$ -n. Säure
Dibrom-veratrumsäure .	8.005	4.736	47.30	47.03	
	9.236	5.465	47.29		5.44
Tribrom-phenol	4.619	4.210	72.84	72.48	4.26
	6.055	5.514	72.78		5.54
	5.079	4.592	72.26		4.62
Brom-campher	6.383	2.802	35.08	34.59	2.82
	9.626	4.143	34.40		4.22
	5.451	2.353	34.50		2.36
Bromalhydrat + H ₂ O ..	6.568	6.229	75.80	75.68	
Bromalhydrat	6.746	6.693	79.29	80.24	
	3.331	3.336	80.04		3.35

Berichtigungen.

Jahrg. 65, [1932], Heft 1, S. 61, 100 mm v. o. lies:



Jahrg. 65 [1932], Heft 2, S. 148, Kopf der Tabelle: lies „E($\Sigma_7 - \Sigma_a$) in %“ statt „E Σ_7 “.

Jahrg. 65 [1932], Heft 2, S. 293, 18 mm v. o. lies „=C₁₀H₇“ statt „=C₆H₅“