



## Carbon Nanotubes

G. Q. Ning, H. Shinohara\*

### Unsynchronized Diameter Changes of Double-Wall Carbon Nanotubes during Chemical Vapour Deposition Growth

**Unsynchronized growing!** Unsynchronized diameter changes of the inner and the outer tubes are observed in the double-wall carbon nanotubes (DWNTs) prepared by CoMo/MgO catalysts. The difference of the growth surroundings for the inner and outer tubes of DWNTs can consistently explain the observed unsynchronized diameter changes.



*Chem. Asian J.*  
DOI: [10.1002/asia.200800347](https://doi.org/10.1002/asia.200800347)




## Glycobiology


K. J. Loft, P. Bojarová, K. Slámová, V. Křen,\* S. J. Williams\*

### Synthesis of Sulfated Glucosaminides for Profiling Substrate Specificities of Sulfatases and Fungal $\beta$ -N-Acetylhexosaminidases

**Systematic sulfation:** Sulfated glycoconjugates are degraded either by desulfation followed by glycoside cleavage, or by glycoside cleavage followed by desulfation. To study these processes, here we report the synthesis of four regioisomerically sulfated *p*-nitrophenyl glucosaminides from the common precursor *p*-nitrophenyl *N*-acetyl- $\beta$ -D-glucosaminide. These substrates allowed the rapid analysis of the substrate preferences of a set of four sulfatases and 24 hexosaminidases.



*ChemBioChem*  
DOI: [10.1002/cbic.200800656](https://doi.org/10.1002/cbic.200800656)




## Dendrimers

G. Bergamini, P. Ceroni,\* V. Balzani, R. Kandre, O. Lukin\*

### Dendrimers with a Pentaphenylene Core: A Photophysical Study

**A plodding dendrimer:** Intense violet luminescence both in solution and in the solid state is shown by a family of dendrimers with a *p*-pentaphenylene core and sulfonimide branches. The fourth-generation dendrimer (see image) has an extremely high steady-state fluorescence anisotropy in dichloromethane solution at 293 K.



*ChemPhysChem*  
DOI: [10.1002/cphc.200800597](https://doi.org/10.1002/cphc.200800597)




## Antibiotics


S. Petit, Y. Duroc, V. Larue, C. Giglione, C. Léon, C. Soulama, A. Denis, F. Dardel, T. Meinnel, I. Artaud\*

### Structure–Activity Relationship Analysis of the Peptide Deformylase Inhibitor 5-Bromo-1*H*-indole-3-acetohydroxamic Acid

**SAR by NMR.** A series of indole compounds derived from 5-bromo-1*H*-indole-3-acetohydroxamic acid were synthesized. Their inhibitory activities were evaluated against purified peptide deformylases (PDFs), and their antibacterial activities against *B. subtilis*, *E. coli* (wild-type and *tolC*), and a variety of pathogens were also determined. The potency of the best inhibitors was related to the NMR footprints of the respective acids with  $^{15}\text{N}$ -labeled *E. coli* Ni-PDF.



*ChemMedChem*  
DOI: [10.1002/cmdc.200800251](https://doi.org/10.1002/cmdc.200800251)





*Eur. J. Inorg. Chem.*  
DOI: 10.1002/ejic.200801044

### Dinitrogen Activation

L. Manceron,\* O. Hübner, H.-J. Himmel\*

#### Dinitrogen Activation by the $Ti_2N_2$ Molecule: A Matrix Isolation Study

$N_2$  activation by a nitride: reaction of matrix-isolated  $Ti_2(\mu-N)_2$  with  $N_2$  affords several new  $N_2$  complexes with different degrees of  $N_2$  bond activation. In neat solid  $N_2$  matrices, the complex  $[(N_2)_4Ti]_2(\mu-N)_2(\mu-\eta^2:\eta^2-N_2)$  is formed.




*Eur. J. Org. Chem.*  
DOI: 10.1002/ejoc.200801128

### Crown Ether Rotaxanes

S. Li, K. Zhu, B. Zheng, X. Wen, N. Li, F. Huang\*

#### A Bis(*m*-phenylene)-32-crown-10/Paraquat [2]Rotaxane

The first bis(*m*-phenylene)-32-crown-10/paraquat [2]rotaxane was synthesized by the threading-followed-by-stoppering strategy. The successful preparation of this [2]rotaxane showed unambiguously that pseudorotaxane-type complexation, rather than the previously reported taco-complex-type complexation, exists for complexation between bis(*m*-phenylene)-32-crown-10 and paraquat derivatives in solution.





*Angew. Chem. Int. Ed.*  
DOI: 10.1002/anie.200806070

### Protein Folding

A. Katranidis, D. Atta, R. Schlesinger, K. H. Nierhaus, T. Choli-Papadopoulou, I. Gregor, M. Gerrits, G. Büldt,\* J. Fitter\*

#### Fast Biosynthesis of GFP Molecules: A Single-Molecule Fluorescence Study

**It's not easy being green:** Real-time visualization of labeled ribosomes and de-novo-synthesized green fluorescent protein molecules using single-molecule-sensitive fluorescence microscopy demonstrates that the mutant GFPem is produced with a characteristic time of five minutes. Fluorescence of the fastest GFP molecules appears within one minute (see picture).



*ChemSusChem*  
DOI: 10.1002/cssc.200800198

### Biofuel Production

R. Kreiter, M. D. A. Rietkerk, H. L. Castricum, H. M. van Veen, J. E. ten Elshof, J. F. Vente\*

#### Stable Hybrid Silica Nanosieve Membranes for the Dehydration of Lower Alcohols

**A thirst for water:** Organic-inorganic hybrid silica nanosieve membranes with narrow pore size distributions were developed for the separation of binary (bio)alcohol/water mixtures, for example, to remove water from wet biofuels during production. These membranes dehydrate lower alcohols and show a stable performance in the presence of significant amounts of acetic acid.

