


CHEMISTRY

A EUROPEAN JOURNAL

16/16

2010



Trifluoromethylated derivative $\text{Sc}_3\text{N}@\text{C}_{80}(\text{CF}_3)_2$...

... exhibits three reversible reductions and two reversible oxidations, and affords the facile generation of the paramagnetic monocation, monoanion, and trianion in solution, which can be characterized by ESR spectroscopy. In their Communication on page 4721 ff., A. A. Popov, J. P. Phillips, S. Stevenson, S. H. Strauss, O. V. Boltalina, L. Dunsch et al. reveal a spin-density shift from the fullerene cage to the Sc_3N cluster on going from the cation to trianion, with a high spin density on one Sc atom in the latter.

A Journal of



Supported by
ACES

Inside Cover

Alexey A. Popov*, Natalia B. Shustova, Anna L. Svitova, Mary A. Mackey, Curtis E. Coumbe, J. Paige Phillips*, Steven Stevenson*, Steven H. Strauss*, Olga V. Boltalina*, and Lothar Dunsch*

Trifluoromethylated derivative $\text{Sc}_3\text{N}@\text{C}_{80}(\text{CF}_3)_2$...

... exhibits three reversible reductions and two reversible oxidations, and affords the facile generation of the paramagnetic monocation, monoanion, and trianion in solution, which can be characterized by ESR spectroscopy. In their Communication on page 4721 ff., A. A. Popov, J. P. Phillips, S. Stevenson, S. H. Strauss, O. V. Boltalina, L. Dunsch et al. reveal a spin-density shift from the fullerene cage to the Sc_3N cluster on going from the cation to trianion, with a high spin density on one Sc atom in the latter.

