## SPOTLIGHTS ...



On these pages, we feature a selection of the excellent work that has recently been published in our sister journals. If you are reading these pages on a computer, click on any of the items to read the full article. Otherwise please see the DOIs for easy online access through Wiley InterScience.




#### Structure Determination

D. Šišak, L. B. McCusker,\* G. Zandomeneghi, B. H. Meier,\* D. Bläser, R. Boese,\* W. B. Schweizer, R. Gilmour, J. D. Dunitz\*

## The Crystal Structure of D-Ribose—At Last!

Better late than never! The  $\beta$ -furanose form of D-ribose, present in countless biomolecules, does not occur in the crystalline compound. X-ray diffraction and NMR experiments show that D-ribose occurs in two crystal forms that contain  $\beta$ - and  $\alpha$ -pyranose forms in various ratios.



Angew. Chem. Int. Ed. DOI: 10.1002/anie.201001266




## Energy Transfer

Y. Zeng, Y.-Y. Li, J. Chen,\* G. Yang,\* Y. Li\*

#### **Dendrimers: A Mimic Natural Light-Harvesting System**

Passed on through the generations: Dendrimers are well-defined tree-like macromolecules having numerous chain ends all emanating from a single core, which make them mimics of natural light-harvesting systems. Recent developments of light-harvesting dendrimers will be discussed in this article, focusing on their energy transfer and electron transfer properties.



Chem. Asian J. DOI: 10.1002/asia.200900653




### Vitamins -

T. B. Fitzpatrick,\* C. Moccand, C. Roux

## Vitamin B<sub>6</sub> Biosynthesis: Charting the Mechanistic Landscape


**Enzyme of the rings**: Vitamin B<sub>6</sub> biosynthesis through the DXP-independent route is catalyzed by PLP synthase. The enzyme utilizes ribose 5-phosphate, glyceraldehyde 3-phosphate and ammonia to synthesize the cofactor form of the vitamin, pyridoxal 5'-phosphate (PLP). This review provides the emerging mechanistic details of this remarkable Pdx1:Pdx2 glutamine amidotransferase complex.



ChemBioChem
DOI: 10.1002/cbic.201000084



# ... ON OUR SISTER JOURNALS



**ChemPhysChem** DOI: 10.1002/cphc.200900966


## Electron Transfer

H. K. Ly, M. A. Marti, D. F. Martin, D. Alvarez-Paggi, W. Meister, A. Kranich, I. M. Weidinger, P. Hildebrandt,\* D. H. Murgida\*

Thermal Fluctuations Determine the Electron-Transfer Rates of Cytochrome c in Electrostatic and Covalent Complexes

Interfacial electron-transfer processes of cytochrome c covalently or electrostatically bound to electrodes coated with self-assembled monolayers (SAMs) of ω-functionalized alkanethiols are studied by time-resolved surface-enhanced resonance Raman spectroscopy and molecular dynamics simulations (see picture).





ChemMedChem DOI: 10.1002/cmdc.201000065


#### Antitumor Agents

Q. Hu, M. Negri, S. Olgen, R. W. Hartmann\*

The Role of Fluorine Substitution in Biphenyl Methylene **Imidazole-Type CYP17 Inhibitors for the Treatment of Prostate** Carcinoma

**CYP17** inhibition is a promising approach for the treatment of prostate cancer. Modification of biphenyl methylene imidazoles by fluorine substitution significantly increases the inhibitory potency of this compound class and prolongs plasma half-life. Compound 9 (ball-and-stick structure) was found to be a potent CYP17 inhibitor (IC<sub>50</sub>=131 nm) with good pharmacokinetic properties.





**ChemSusChem** DOI: 10.1002/cssc.200900234


## Biorenewables

O. Busse, K. Räuchle, W. Reschetilowski\*

Hydrocracking of Ethyl Laurate on Bifunctional Micro-/Mesoporous Zeolite Catalysts

Olefins from biomass: Metal-modified micro- and mesoporous composite materials are promising catalyst systems for the conversion of ethyl laurate, as a model compound for vegetable oils. The selectivity to light olefins can be enhanced up to 60 wt %.





**ChemCatChem** DOI: 10.1002/cctc.201000027

### Supported Enzymes

M. Shakeri, K. Engström, A. G. Sandström, J.-E. Bäckvall\*

Highly Enantioselective Resolution of β-Amino Esters by Candida antarctica Lipase A Immobilized in Mesocellular Foam: Application to Dynamic Kinetic Resolution.

Candida antarctica lipase A (CALA) immobilized in functionalized mesocellular foam (MCF) in the presence of sucrose facilitated a dramatic increase in enantioselectivity for the kinetic resolution (KR) of representative β-amino esters. The temperature of optimum activity of CALA shifted from 20-30°C to 80-90°C on immobilization in the MCF. Combination of the immobilized CALA with a ruthenium complex allowed dynamic KR at 90°C.

© 2010 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim



7363

## **SPOTLIGHTS**




## Catalyst Regeneration

#### T. Pintauer\*

### Catalyst Regeneration in Transition-Metal-Mediated Atom-Transfer Radical Addition (ATRA) and Cyclization (ATRC) Reactions

Recent advances in the area of catalyst regeneration in copperand ruthenium-mediated atom transfer radical addition (ATRA) and cyclization (ATRC) reactions in the presence of free-radical diazo initiators or magnesium as reducing agents were reviewed. Reducing agents regenerate the activator in both processes, enabling selective ATRA and ATRC reactions using very small amounts of metal catalysts.



Eur. J. Inorg. Chem. DOI: 10.1002/ejic.201000234




## Red Fluorescence

H. Langhals,\* T. Pust

## **Axially Extended Perylene Dyes**

Perylenecarboxylic anhydride imides were condensed with semicarbazide and thiosemicarbazide to obtain strongly red-fluorescent triazolinone and triazolinthione derivatives, respectively, suitable for nucleophilic labelling. A Schönberg reaction of the latter with diazoalkanes allowed the synthesis of spirothiiranes.



Eur. J. Org. Chem. DOI: 10.1002/ejoc.201000230

