

SYNTHESIS AND REACTIONS OF NEW HIGHLY STRAINED

UNSATURATED PROPELLANES, [4.2.1]PROPELLA-2,4,7-TRIENES

*


Fumio TODA, Takayuki YOSHIOKA, and Masao KAWADA

Department of Industrial Chemistry, Faculty of Engineering, Ehime University,


Matsuyama 790

The title propellane, 2,7-di-tert-butyl-11,11-dichloro-4,5,9,10-tetraaryl-tetracyclo[6.2.1.0.^{3,6}]undeca-2,4,6,9-tetraene (3), was prepared by cycloaddition of dichlorocarbene to the benzene nucleus of 2,7-di-tert-butyl-4,5,9,10-tetraaryl-tricyclo[6.2.0.0.^{3,6}]deca-1,3(6),4,7,9-pentaene (1). On Ag⁺-induced methanolysis and acetolysis of 3a, its endo-Cl reacted easily and endo-methoxy (6) and endo-acetoxy derivative (9) were obtained, respectively, probably via bishomoallyl cation of pyramidal structure (5). Hydrolysis of 3a and 9 afforded 1a.


The most highly strained unsaturated propellane which has been isolated so far is derivative of [4.2.2]propella-2,4,7-triene (2c, 2d)¹⁾ prepared by cycloaddition of tetracyanoethylene to 2,7-di-tert-butyl-4,5,9,10-tetra(p-halophenyl)tricyclo[6.2.0.0.^{3,6}]deca-1,3(6),4,7,9-pentaene (1c, 1d). We now report the isolation and some novel reactions of derivative (3) of more strained unsaturated propellane, [4.2.1]propella-2,4,7-triene (4). We also report Ag⁺-induced methanolysis and acetolysis of 3a which afford the endo-11-methoxy (6) and endo-11-acetoxy derivative (9), respectively, probably via bishomoallyl cation of pyramidal structure (5), and hydrolysis of 3a and 9 which affords 1a.




1



2



3




4

a:Ar = Ph ; b:Ar = p-Me-C₆H₄- ; c: Ar = p-Cl-C₆H₄- ; d: Ar = p-Br-C₆H₄- ; R = C(CH₃)₃

Stirring of a mixture of 1a (1 g), 50% aqueous KOH (30 ml), CHCl_3 (30 ml), and a catalytic amount of benzyltriethylammonium chloride under N_2 at room temperature for 2 h afforded 3a (0.23 g; 20%) as colorless needles, after purification by column chromatography on Al_2O_3 (CCl_4); mp 218-219 °C; $\lambda_{\text{max}}^{2)}$ 254 (ϵ 33800) and 320 sh nm (7300); $^1\text{H-NMR}$ δ 0.92 (s, tBu, 18H) and 7.08-7.80 (m, Ph, 20H); $^{13}\text{C-NMR}$ (for ^{13}C -isotope enriched sample, 20% each carbon of C_1 , C_3 , C_6 , and C_8) δ 30.4 (C_{13}), 34.0 (C_{12}), 47.3 ($\text{C}_{1,8}$), and 140.7 ppm ($\text{C}_{3,6}$). Similar treatments of 1b, 1c, and 1d afforded 3b (29%, mp 217-218 °C), 3c (24%, mp 236-237 °C), and 3d (18%, mp 205-206 °C), respectively.

Treatment of 3a with excess AgClO_4 in MeOH at room temperature for 10 min afforded the 11-methoxy derivative (6) as colorless needles in 70% yield; mp 189-191 °C; λ_{max} 260 (ϵ 31700) and 319 sh nm (10600); $^1\text{H-NMR}$ δ 0.96 (s, tBu, 18H), 3.73 (s, OMe, 3H), and 7.22 (bs, Ph, 20H). Further treatment of 6 under the same conditions for 12 h afforded the 11,11-dimethoxy derivative (7) as colorless prisms in 70% yield; mp 188-189 °C; λ_{max} 248 (ϵ 33600) and 321 sh nm (9200); $^1\text{H-NMR}$ δ 0.96 (s, tBu, 18H), 3.51 and 3.70 (each s, OMe, 3H), and 7.0-7.7 (m, Ph, 20H). On the same treatment of 6 for 1 h, however, 6 was recovered unchanged in 85% yield. Of the two MeO signals of 7, the one at relatively higher magnetic field (δ 3.51) can be ascribed to exo-methoxy group (A) which is shielded by Ph groups on the C_9 - and C_{10} -positions. The MeO groups of 6 should be at the endo-position, because its chemical shift (δ 3.73) is comparable to that of the endo-MeO (B) of 7 (δ 3.70). Similar treatments of 3a with AgClO_4 in EtOH for 10 min and 12 h afforded ethoxy-analog of 6 (67%, mp 188-190 °C) and diethoxy-analog of 7 (43%, mp 169-170 °C), respectively.


Extremely faster conversion of 3a into 6 than that of 6 into 7, can well be interpreted by assuming bishomoallyl cation of pyramidal structure (5) as an intermediate, which is formed by Ag^+ -induced elimination of the endo-Cl accompanied by participation of p-orbitals of cyclobutene ring. In the case of 6 with syn-relationship of Cl and p-orbitals, however, such the neighboring group participation would not occur efficiently. Intermediacy of bishomoallyl cation in the solvolysis of various substituted carbocycles, but not of propellane, has well been documented.³⁾ Treatments of 3a with MeMgI in ether at room temperature for 2 h and with AgOAc in AcOH under reflux for 1 h afforded its methyl-derivative (8) in 42% yield; mp 199-200 °C; λ_{max} 251 (ϵ 24200) and 323 sh nm (6500); $^1\text{H-NMR}$ δ 0.92 (s, tBu, 18H), 2.44 (s, Me, 3H), and 7.45 (bs, Ph, 20H) and acetoxy-derivative (9) in 51% yield; mp 202 °C; λ_{max} 252 (ϵ 34400) and 325 sh nm (9200); $^1\text{H-NMR}$ δ 0.96 (s, tBu, 18H), 2.29 (s, OAc, 3H), and 7.41 (s, Ph, 20H), respectively. Neither 8 nor 9 was caused further substitution reaction by prolonged reaction. Both 8 and 9 would also be produced by the reaction pathway which proceeds via the bishomoallyl cation (5), and their Me and AcO groups would also be at endo-positions.



All the above mentioned Ag^+ -induced substitution reactions of $\underline{3a}$ and $\underline{6}$ are in contrast to those of Ag^+ -induced solvolysis of saturated propellane. For example, it has been reported that Ag^+ -induced reaction of $\underline{10}$ with nucleophile (Nu^-) affords various products successively via homoallyl cation ($\underline{11}$) and bridge-head olefin intermediate ($\underline{12}$).⁴⁾

The most interesting reaction of the propellanes is hydrolysis of $\underline{3a}$ and $\underline{9}$ to $\underline{1a}$. Heating of $\underline{9}$ in acetone containing a small amount of aqueous NaOH under reflux for 2.5 h afforded $\underline{1a}$ in 72% yield. This reaction probably proceeds by decarbonylation of propellanonone ($\underline{14}$) which was initially produced by hydrolysis of $\underline{9}$ into $\underline{13}$ followed by elimination of HCl. Reaction type of the decarbonylation of $\underline{14}$ is similar to that of the previously reported thermal cycloreversion of $\underline{2}$ into $\underline{1}$ and tetracyanoethylene.¹⁾ Treatment of $\underline{3a}$ with excess $AgClO_4$ in wet tetrahydrofuran (THF) resulted in a greenish blue solution, and this color disappeared gradually and finally afforded AgCl, Ag-metal, and 2,3,5,6-tetrabenzoyl-1,4-di-tert-butylbenzene ($\underline{19}$) in 27% yield.⁵⁾ This color is probably attributed to a 1:1 complex of Ag^+ and $\underline{1a}$ which was derived from $\underline{3a}$ successively via $\underline{13}$ and $\underline{14}$, because a solution of $\underline{1a}$ and an equimolar amount of $AgClO_4$ in dry benzene showed the same color (λ_{max} 592 nm; 1H -NMR δ 1.09 and 1.24 (each s, tBu, 9H)), and because decomposition of a solution of $\underline{1a}$ and excess $AgClO_4$ in dry benzene with water gave Ag-metal and $\underline{19}$ in 66% yield. Magnetic nonequivalence of the two tBu groups

suggests that this complex is unsymmetrical as depicted (15). Although mechanism of the conversion of 15 into 19 is not clear, a plausible pathway is that proceeds via cations such as 16 and 18, because the formation of Ag metal supports production of cation by one-electron oxidation process, and because THF easily polymerized when a solution of 15 in THF was kept at room temperature. It has been reported that Lewis acid-catalyzed polymerization of THF is accelerated by cyclic compound which easily undergoes cationic ring-opening reaction.⁶⁾



References and Notes

- 1) F. Toda and T. Yoshioka, Chem. Lett., 1977, 561.
- 2) UV and NMR spectra were measured in CHCl₃ and CDCl₃, respectively. All new Compounds gave satisfactory elemental analyses and mass spectral data.
- 3) P. D. Bartlett, "Nonclassical Ions," W. A. Benjamin, Inc., New York (1965).
- 4) P. M. Warner, S-L. Lu, E. Myers, P. W. DeHaven, and R. A. Jacobson, J. Am. Chem. Soc., 99, 5102 (1977).
- 5) F. Toda, N. Dan, K. Tanaka, and Y. Takehira, J. Am. Chem. Soc., 99, 4529 (1977).
- 6) T. Saegusa and Y. Matsumoto, J. Synth. Org. Chem. Jpn., 24, 713 (1966).

(Received March 20, 1978)