

ALKYLATION OF ACTIVE HYDROGEN COMPOUNDS WITH ALLYLIC AND BENZYLIC
ALCOHOLS UNDER CoCl_2 CATALYSIS. A USEFUL SYNTHESIS OF GRIFOLIN[†]


J. Marquet and M. Moreno-Mañas*

Departamento de Química Orgánica. Universidad Autónoma
de Barcelona. Cerdanyola. Barcelona. Spain

CoCl_2 is a useful catalyst for the condensation of allylic and benzylic alcohols with active hydrogen compounds. Under these neutral conditions orcinol and farnesol react to afford the antibiotic grifolin.

The alkylation of active hydrogen compounds in neutral media is a subject of general interest since acid and alkaline conditions reduce the range of suitable starting reagents and can deteriorate the initial alkylation products. The reactions of alkaline salts of β -dicarbonyl compounds with alkyl halides offer typical examples of such limitations^{1,2,3}.


In our search for better alkylating methods of β -dicarbonyl compounds we found that bis-(pentane-2,4-dionato)cobalt(II) reacts with several reactive alkyl halides. The final products are the alkylated pentane-2,4-diones⁴. When this method was applied to benzhydrylbromide, 1, and the cobalt(II) complex of dehydroacetic acid, 2⁵, alkylation at C-3 was not observed. Instead, upon heating in the absence of solvent, 3-acetyl-5-benzhydryl-4-hydroxy-6-methyl-2-pyrone, 3⁵ (m.p. 135-7°; $\nu(\text{KBr})$: 1725, 1630(sh), 1600 cm^{-1} ; $\text{pmr}(\text{CDCl}_3)$: δ 17,6(s., 1H), 7,45-7,00(m., 10H),



5.80(s., 1H), 2.65(s., 3H), 2.00(s., 3H); mass spectrum: m/e 334(M^+ , 5), 206(24), 178(17), 168(28), 153(27), 129(36), 125(25), 115(20), 111(24), 98(21), 85(98), 83(100), 77(25), 69(42), 43(94); UV(cyclohexane): λ 322 nm ($\log \epsilon$: 4.09), was isolated. Also the pyrones 4 (m.p. 127-30°), 5⁸ (oil, never isolated in pure form), and 6^{5,8} (m.p. 161-2°) were characterized by standard methods. Product 4 must be formed by ring opening of dehydroacetic acid, 7, followed by decarboxylation and cyclization. Similarly 5 can derive from 3 although alkylation of 4 is also possible. Most probably 6 comes from alkylation of 5 since dialkylation on 2 (or 7) was never observed.


However, when a sample was taken off after refluxing for 3 h. in dioxane, only 7 and dibenzhydryl ether were characterized as organic products.

The unusual reaction at C-5 of the dehydroacetic acid ring, position which is not activated by the cobalt complex, together with the other facts above described, strongly suggested the operativity of a catalytic effect of the cobalt species, the ether 8 acting as a real alkylating agent. Indeed, when 7 and 1 were reacted under the catalytic influence of the complex 2, similar results were obtained.














As a consequence we undertook a research on the catalytic effects of some metallic complexes and salts.

The adoption of alkyl halides as alkylating agents results on the undesirable formation of one equivalent of hydrogen halide. We have observed that this has a deleterious effect upon the direct alkylation products. For instance, in one of our best experiments, treatment of 4-hydroxy-6-methyl-2-pyrone, 9, with 1-chloro-3-methyl-2-butene, under bis-(pentane-2,4-dionato)nickel(II) catalysis⁶, led to the cyclized 2,8-dioxa-3,3,9-trimethylbicyclo[4.4.0]deca-1(6),9-dien-7-one, 10 (oil, pmr (CDCl_3) δ 5.75(s., 1H), 2.45(t., J ca. 6.7 Hz, 2H), 2.20(s., 3H), 1.79(t., J ca. 6.7 Hz, 2H), 1.35(s., 6H).



Therefore, we turned upon alcohols as alkylating agents. For a variety of substrates, and in the absence of solvent, anhydrous CoCl_2 is an efficient catalyst although its conversion into a hexahydrate makes necessary its use in rather large quantities (0.2 mol unless otherwise stated). Our initial results are collected below:





Two natural products, the antibiotics grifolin, 11^{15,16,17} and its isomer neogrifolin, 12^{16,17}, were prepared from orcinol and farnesol (having ca. 30% of the 2-Z isomer). After careful chromatography on silica gel, the yields of isolated 11 and 12 were respectively 17 and 32%, which compares well with those previously reported from non neutral condensation techniques¹⁸⁻²³.

The ir of 11 was identical to that reproduced in the literature¹⁸. However, the presence of minor amounts of the 2'-Z isomers of both 11 and 12^{8,24} can not be excluded.

At present we are investigating the scope of this method.

We acknowledge financial support from the "Comisión Asesora de Investigación Científica y Técnica" of Spain, and a generous gift of farnesol from Givaudan Ibérica S.A. (Barcelona).

REFERENCES AND FOOTNOTES

- † This paper is number IV in the series "Metal Complexes in Organic Synthesis". For number III see ref. 3.
- 1.- "Some Modern Methods of Organic Synthesis". W. Carruthers; Cambridge University Press. 2nd Ed. 1978. Page 4.
 - 2.- S. Boatman, T.M. Harris, C.R. Hauser; J. Org. Chem., 30, 3321, (1965).
 - 3.- M. Boya, J. Marquet, M. Moreno-Mañas, M. Prior; Anales de Quím., 75, 920, (1979).
 - 4.- J. Marquet, M. Moreno-Mañas; Synthesis, 1979, 348.
 - 5.- Correct elemental analysis was secured for this compound.
 - 6.- M. Yamashita, Y. Takegami; Synthesis, 1977, 803.
 - 7.- Compared with an authentic sample (ref. 3).
 - 8.- Correct spectroscopic behaviour.
 - 9.- K.H. Boltze, K. Heidenbluth; Chem. Ber., 92, 982, (1959).
 - 10.- Pmr identical to that described by M.H. Palmer, G.J. McVie; J. Chem. Soc.(B), 1968, 742.
 - 11.- Z. Földi; Chem. Ber., 61, 1609, (1928).
 - 12.- E.C. Du Feu, F.J. McQuillin, R. Robinson; J. Chem. Soc., 1937, 53.
 - 13.- A.O. Fitton, G.R. Ramage; J. Chem. Soc., 1962, 4870.
 - 14.- Pmr, ir and mass spectra in agreement to those described by N. Schamp, R. Verhe, L. De Buyck; Tetrahedron, 29, 3857, (1973).
 - 15.- T. Goto, H. Kakisawa, Y. Hirata; Tetrahedron, 19, 2079, (1963).
 - 16.- J. Vrkoc, M. Budesinsky, L. Dolejs; Phytochemistry, 16, 1409, (1977).
 - 17.- H. Besl, G. Höfle, B. Jendry, E. Jägers, W. Steglich; Chem. Ber., 110, 3770, (1977).
 - 18.- M. Isobe, T. Goto; Tetrahedron, 24, 945, (1968).
 - 19.- G. Cardillo, R. Cricchio, L. Merlini, G. Nasini; Gazz. Chim. Ital., 99, 308, (1969).
 - 20.- K. Ima-ye, H. Kakisawa; J.C.S. Perkin I, 1973, 2591.
 - 21.- G.D. Manners, L. Jurd, K.L. Stevens; Chem. and Ind., 1974, 616.
 - 22.- J.H.P. Tyman, W.A. Baldwin, C.J. Strawson; Chem. and Ind., 1975, 41.
 - 23.- S. Yamada, F. Ono, T. Katagiri, J. Tanaka; Synth. Commun., 8, 241, (1978).
 - 24.- Pmr spectrum as described in ref. 22.
 - 25.- Commercial orcinol is a monohydrate.

(Received November 26, 1980)