


SYNTHESIS OF 2-AMINO-3,8-DIMETHYLIMIDAZO[4,5-*f*]QUINOXALINE (Me-IQx),  
A POTENT MUTAGEN ISOLATED FROM FRIED BEEF

Hiroshi KASAI, Tomoko SHIOMI, Takashi SUGIMURA<sup>+</sup> and Susumu NISHIMURA<sup>\*</sup>


Biology Division and <sup>+</sup>Biochemistry Division, National Cancer Center Research Institute,  
5-1-1, Tsukiji, Chuo-ku, Tokyo 104

A potent mutagen, 2-amino-3,8-dimethylimidazo[4,5-*f*]quinoxaline (Me-IQx), isolated from fried beef and its 3,7-dimethyl derivative were synthesized from 6-amino-3-methylquinoxaline and 6-amino-2-methylquinoxaline, respectively. These compounds showed strong mutagenic activity towards *Salmonella typhimurium* TA98 in the presence of S9 Mix.


We previously reported isolation of a new mutagenic compound from fried beef and its structural determination as 2-amino-3,8-dimethylimidazo[4,5-*f*]quinoxaline (Me-IQx, structure 1)<sup>1)</sup>. It is an aza-derivative of the mutagenic compounds 2-amino-3-methylimidazo[4,5-*f*]quinoline (IQ) and 2-amino-3,4-dimethylimidazo[4,5-*f*]quinoline (Me-IQ), which were previously isolated from broiled sardines<sup>2-5)</sup>. For the final assignment of the structure of Me-IQx and for further studies on its biological properties, it was necessary to synthesize chemically Me-IQx and its isomer having a methyl group in the 7-position in place of the 8-position. In this communication, we report a chemical synthesis of the mutagen Me-IQx and its isomer.



Me-IQx (1) was synthesized from 6-amino-2-methylquinoxaline (2) as shown in scheme 2 by a similar procedure to that used for the synthesis of IQ<sup>3</sup>) and Me-IQ<sup>5</sup>). The starting material for this synthesis, compound 2, was prepared by condensation of 1,2,4-triaminobenzene with methylglyoxal by the method of Klicnar and Kosek<sup>6</sup>). Contrary to their finding<sup>6</sup>) that only 6-amino-3-methylquinoxaline (2) was formed as a reaction product, in our experiment compound 2 and its isomer 3 were produced in about equal proportions. These products were separated by high pressure liquid chromatography on a TSK-GEL LS-410 column. Structural assignments of these compounds were



Scheme 1



Scheme 2

made by comparison of their properties with those of an authentic sample of compound 3, prepared separately by reduction of 2-methyl-6-nitroquinoxaline (4) which was obtained by condensation of 1,2-diamino-4-nitrobenzene with methylglyoxal (Scheme 1). In the latter condensation reaction, the more basic 2-amino-group reacts with the aldehyde group in methylglyoxal by nucleophilic addition to give a single product 4 and the structure of its reduction product 3 was established previously by unequivocal synthesis from 2,4-dinitrophenylalanine<sup>6</sup>).

Compound 2, thus obtained, was first converted to the tosyl derivative 5 and nitrated with  $\text{HNO}_3\text{-AcOH}$  mixture to afford compound 6, which was hydrolyzed with  $\text{H}_2\text{SO}_4$  to yield 6-amino-3-methyl-5-nitroquinoxaline (7). Compound 7 was then treated with  $\text{NaH}$  and methylated with  $\text{CH}_3\text{I}$  to afford compound 8. Then compound 8 was reduced to 5-amino derivative with  $\text{Fe-HCl}$  mixture and the neutralized reaction mixture was treated with  $\text{BrCN}$  to afford compound 1. UV ( $\lambda_{\text{max}}^{\text{MeOH}}$ ,  $\epsilon$ ): 214 (24,300), 274 (41,100), 240 (3,900); NMR ( $\delta_{\text{CDCl}_3}$ , J): 8.65 ppm (1H, s, 7-H), 7.75 (1H, d, 9Hz, 5-H), 7.51 (1H, d, 9Hz, 4-H), 4.88 (2H, s, broad,  $-\text{NH}_2$ ), 3.69 (3H, s, N- $\text{CH}_3$ ), 2.80 (3H, s, C- $\text{CH}_3$ ).




Fig. 1. Mutagenicity of Me-IQx to *S. typhimurium*.

a: The mutagenicity of 0.05 µg of Me-IQx was tested with various amounts of S9.

● TA98; ○ TA100.

b: Dose-response curve for mutagenicity obtained with S9 Mix containing 10 µl S9.

● TA98; ○ TA100.

In the same fashion, 2-amino-3,7-dimethylimidazo[4,5-*f*]quinoxaline (9), an isomer of Me-IQx, was synthesized from 6-amino-2-methylquinoxaline (3). UV ( $\lambda_{\text{max}}^{\text{MeOH}}$ ,  $\epsilon$ ): 214 (29,000), 274 (42,000), 237 (4,000); NMR ( $\delta_{\text{CDCl}_3}$ , J): 8.74 ppm (1H, s, 8-H), 7.73 (1H, d, 9Hz, 5-H), 7.60 (1H, d, 9Hz, 4-H), 5.10 (2H, s, broad, -NH<sub>2</sub>), 3.72 (3H, s, N-CH<sub>3</sub>), 2.77 (3H, s, C-CH<sub>3</sub>).

Synthetic compound 1 showed strong mutagenic activity towards *Salmonella typhimurium* with metabolic activation by microsomal enzymes (S9), as shown in Fig. 1a. The specific activities of Me-IQx towards TA98 and TA100 were 145,000 and 14,000 revertants/ $\mu$ g respectively under optimal conditions (Fig. 1b). The isomer of Me-IQx, compound 9, also showed strong mutagenic activity towards TA98 and TA100 (146,000 and 26,000 revertants/ $\mu$ g, respectively). The mutagenic activities of these imidazoquinoxaline derivatives are weaker than those of IQ<sup>3</sup>) and Me-IQ<sup>5</sup>). Large scale synthesis of Me-IQx for carcinogenic tests and for examination of its other biological activities is now in progress.

#### Acknowledgements

We thank Dr. S. Yokoyama, University of Tokyo, for measurements of NMR spectra and Dr. Z. Yamaizumi of our institute for measurements of mass spectra.

#### References

- 1) H. Kasai, Z. Yamaizumi, T. Shiomi, S. Yokoyama, T. Miyazawa, K. Wakabayashi, M. Nagao, T. Sugimura, and S. Nishimura, *Chem. Lett.* 1981, 485.
- 2) H. Kasai, Z. Yamaizumi, K. Wakabayashi, M. Nagao, T. Sugimura, S. Yokoyama, T. Miyazawa, N.E. Spingarn, J.H. Weisburger, and S. Nishimura, *Proc. Japan Acad.* **56B**, 278 (1980).
- 3) H. Kasai, S. Nishimura, K. Wakabayashi, M. Nagao, and T. Sugimura, *Proc. Japan Acad.* **56B**, 382 (1980).
- 4) H. Kasai, Z. Yamaizumi, S. Nishimura, K. Wakabayashi, M. Nagao, T. Sugimura, N.E. Spingarn, J.H. Weisburger, S. Yokoyama, and T. Miyazawa, *J. Chem. Soc. Perkin Transaction I*, in press.
- 5) H. Kasai, Z. Yamaizumi, K. Wakabayashi, M. Nagao, T. Sugimura, S. Yokoyama, T. Miyazawa, and S. Nishimura, *Chem. Lett.* 1980, 1391.
- 6) J. Klicnar and F. Kosek, *Collection Czechoslov. Chem. Commun.* **30**, 3102 (1965).

(Received March 31, 1981)