


SYNTHESES AND PROPERTIES OF BINAPHTHO[1,8-de]-1,3-DITHIIN-2-YLIDENE
AND ITS SELENIUM ANALOGUE


Koji YUI, Yoshio ASO, Tetsuo OTSUBO, and Fumio OGURA*
Department of Applied Chemistry, Faculty of Engineering,
Hiroshima University, Saijo, Higashi-Hiroshima 724

With a view to discovering new electron donors for low-dimensionally metallic materials, the title compounds were prepared via the corresponding 1,8-dichalcogen-bridged naphthalenes. Their donor characters were examined by cyclic voltammetry, as compared with those of the reference compounds.

The discovery of tetrathiafulvalene (TTF), forming charge-transfer salts with low-dimensionally metallic properties, has stirred considerable interest in searching new electron donors which exhibit similar conductivity.¹⁾ The hitherto unknown binaphtho[1,8-de]-1,3-dithiin-2-ylidene 1 belongs to the same tetrathia-ethylene class, but differs structurally from TTF type regarding the fused heterocyclic member. From another viewpoint, it may be regarded as an extended type with ethylene conjugation of 1,8-dichalcogen-bridged naphthalene, which has been noticeable as a novel class of donor.²⁻⁴⁾ In this context, it is very instructive to examine the properties of 1 as a potential donor. In addition, its selenium analogue 2 looks further promising in expectation of enhanced electron transfer due to introduction of the more polarizable chalcogen. We now report the syntheses and electrochemical properties of 1 and 2.




The synthetic route of 1 is shown in Scheme 1. Almost all symmetrical TTF derivatives are generally prepared by coupling of two identical heterocyclic moieties, usually 1,3-dithiole-2-thione or 1,3-dithiolium ion.⁵⁾ Compound 1 is expected to be similarly accessible from desulfurized coupling of naphtho[1,8-de]-1,3-dithiin-2-thione 6. Nakayama et al. already reported the formation of the precursor 6 as a minor product on photolysis of naphtho[1,8-de]-1,2,3-thiadiazine in carbon disulfide.⁶⁾ We have developed an alternative access via naphtho[1,8-cd]-1,2-dithiole 4, which is of choice on large-scale preparation. The intermediate 4 was previously prepared by some methods, i.e., a direct thermal reaction of naphthalene and sulfur,⁷⁾ multistage reactions starting with diazo-



Scheme 1.

tization of 1-aminonaphthalene-8-sulfonic acid,⁸⁾ and a reaction of 1,8-dilithio-naphthalene with sulfur.²⁾ Sandmann's and other groups recently reported direct substitution of unactivated aryl halides involving peri-disubstituted arenes with metal dichalcogenide in a dipolar aprotic solvent.^{4,9)} In a similar manner, sodium was allowed to react with elemental sulfur at 110 °C in hexamethylphosphoric triamide to form sodium disulfide, which was in situ treated with 1,8-dichloro-naphthalene 3¹⁰⁾ at 150 °C, giving 4 in 46% yield.¹¹⁾ Reduction of 4 with sodium borohydride at RT in tetrahydrofuran-ethanol gave quantitatively naphthalene-1,8-dithiol 5,¹²⁾ which was subsequently treated with N,N'-thiocarbonyldiimidazole at -15 °C to afford 6 in 94% yield.¹³⁾ All attempts to convert 6 into the target molecule 1 using phosphorus reagents such as triethyl phosphite, trimethyl phosphite, and triphenyl phosphine were unsuccessful. The exclusive product from heating 6 at 110 °C in triethyl phosphite was assigned to diethyl naphtho[1,8-de]-1,3-dithiin-2-yl phosphonate 7 (92% yield),¹⁴⁾ while the other phosphorus reagents led to unidentified products. The formation of 7 is most likely rationalized by a mechanism as shown in Scheme 2.




Scheme 2.

Compound 7 offers an additional approach to 1, because it may be an appropriate reagent for a Wittig modification developed by Wadsworth and Emmons.¹⁵⁾ Thus 7 was treated with lithium diisopropylamide at -78 °C in tetrahydrofuran to


generate a phosphonate carbanion, which, however, did not react with thione 6. As the real counterpart, naphtho[1,8-de]-1,3-dithiin-2-one 8 was prepared in 97% yield from a reaction of dithiol 5 and *N,N'*-carbonyldiimidazole in a similar manner as described for the synthesis of 6.¹⁶⁾ Finally, treatment of the above phosphonate carbanion with 8 gave binaphtho[1,8-de]-1,3-dithiin-2-ylidene 1 in 92% yield.¹⁷⁾

The synthesis of the selenium analogue 2 was carried out in a similar, but shorter route as shown in Scheme 3. Thus a reaction of 3 with sodium diselenide in hexamethylphosphoric triamide at 100 °C gave naphtho[1,8-cd]-1,2-diselenole 9 in 69% yield.¹⁸⁾ The reduction of 9 with lithium aluminium hydride at RT in tetrahydrofuran, followed by treatment with thiophosgene produced naphtho[1,8-de]-1,3-diselenin-2-thione 10 in 45% yield.¹⁹⁾ In contrast to the sulfur case, a reaction of 10 with trimethyl phosphite in refluxed benzene gave directly the desired binaphtho[1,8-de]-1,3-diselenin-2-ylidene 2 in 66% yield.²⁰⁾



The structures of 1 and 2 were characterized by spectroscopic and elemental analyses. Both ¹H-NMR spectra showed aromatic signals consistent with 1,8-symmetrically disubstituted naphthalene. In addition, both MS spectra showed satisfactory molecular ion peaks involving isotopic peaks due to sulfur or selenium.

The cyclic voltammetry of 1 exhibited a reversible redox wave, whose half-wave oxidation potential was situated at 1.14 V vs. a Ag/AgCl reference electrode in benzonitrile (0.1 M Bu₄NClO₄, Pt electrode, 100 mV·s⁻¹ scan rate). In contrast, the cyclic voltammetry of naphtho[1,8-de]-1,3-dithiin 11²¹⁾ and naphtho[1,8-de]-2-isopropylidene-1,3-dithiin 12²²⁾ showed irreversible oxidations with somewhat higher peak potentials. Thus the easier oxidation for 1 and the higher stability of the resulting radical cation may be attributed to extended conjugation of π-electrons through the central olefin. On the other hand, the cyclic voltammetry of 2 again showed an irreversible oxidation, though its peak potential was the almost same as that of 1. The introduction of selenium does not serve to enhance the donor character of 2 and rather prompts the resulting radical cation to decompose owing to labile C-Se bond relative to C-S bond.



This work was supported by the grant-in-aid of special research project on properties of molecular assemblies (No. 60104002) from the Ministry of Education, Science and Culture, Japan.

References

- 1) For reviews, see F. Wudl, *Acc. Chem. Res.*, 17, 227 (1984); J. M. Williams, M. A. Beno, H. H. Wang, P. C. W. Leung, T. J. Emge, U. Geiser, and K. D. Carlson, *ibid.*, 18, 261 (1985).
- 2) J. Meinwald, D. Dauplaise, F. Wudl, and J. J. Hauser, *J. Am. Chem. Soc.*, 99, 255 (1977); J. Meinwald, D. Dauplaise, and J. Clardy, *ibid.*, 99, 7743 (1977); D. Dauplaise, J. Meinwald, J. C. Scott, H. Temkin, and J. Clardy, *Ann. N. Y. Acad. Sci.*, 313, 382 (1978).
- 3) F. Wudl, D. E. Schafer, and B. Miller, *J. Am. Chem. Soc.*, 98, 252 (1976).
- 4) A. Yamahira, T. Nogami, and H. Mikawa, *J. Chem. Soc., Chem. Commun.*, 1983, 904; J. C. Stark, R. Reed, L. A. Acampora, D. J. Sandmann, S. Jansen, M. T. Jones, and B. M. Foxman, *Organometallics*, 3, 732 (1984).
- 5) M. Narita and C. U. Pittmann, Jr., *Synthesis*, 1976, 489.
- 6) J. Nakayama, T. Fukushima, E. Seki, and M. Hoshino, *J. Am. Chem. Soc.*, 101, 7684 (1979).
- 7) M. Lanfrey, *C. R. Acad. Sci.*, 152, 92 (1911); H. S. Desai and B. D. Tilak, *J. Sci. Ind. Res. (India)*, 19, 390 (1960).
- 8) W. B. Price and S. Smiles, *J. Chem. Soc.*, 1928, 2372; A. Zweig and A. K. Hoffmann, *J. Org. Chem.*, 30, 3997 (1965).
- 9) D. J. Sandmann, J. C. Stark, and B. M. Foxman, *Organometallics*, 1, 739 (1982); D. J. Sandmann, J. C. Stark, L. A. Acampora, and P. Gagne, *ibid.*, 2, 549 (1983).
- 10) G. C. Hampson and A. Weissberger, *J. Chem. Soc.*, 1936, 393.
- 11) 4: reddish brown plates from hexane, mp 123 °C (lit.,⁸) 116 °C).
- 12) 5: colorless leaflets from hexane-THF, mp 122 °C, IR(KBr) 2520, 2540 cm⁻¹(S-H).
- 13) 6: orange needles from hexane-benzene, mp 201-202 °C (lit.,⁶) 199-201 °C), IR (KBr) 1040 cm⁻¹(C=S).
- 14) 7: colorless columns from hexane-benzene, mp 120 °C, ¹H-NMR(CCl₄, 60MHz) δ 1.20 (t, J_{CH-CH}=7.4 Hz, 6H, CH₃), 4.02 (dq, J_{CH-O-P}=8.4 Hz, J_{CH-CH}=7.4 Hz, 4H, CH₂), 4.23 (d, J_{CH-P}=17 Hz, 1H, CHS), 7.2-7.7 (m, 6H, ArH).
- 15) W. S. Wadsworth, Jr., *Org. React.*, 25, 77 (1977).
- 16) 8: colorless needles from hexane-benzene, mp 140 °C, IR(KBr) 1630 cm⁻¹(C=O).
- 17) 1: pale yellow prisms from carbon disulfide, mp 297 °C, ¹H-NMR(CS₂, 360 MHz) δ 7.22-7.28 (m, 8H, ArH), 7.486 (dd, J=7.37 Hz, J'=1.92 Hz, 4H, ArH).
- 18) 9: dark violet needles from hexane, mp 124 °C (lit.,²) 127-129 °C).
- 19) 10: orange needles from hexane, mp 153-154 °C, IR(KBr) 1021 cm⁻¹(C=S).
- 20) 2: pale yellow fine crystals from toluene, mp 287-287.5 °C, ¹H-NMR(CS₂, 360 MHz) δ 7.286 (t, J=7.7 Hz, 4H, ArH), 7.476 (bd, J=7.3 Hz, 4H, ArH), 7.610 (bd, J=8.1 Hz, 4H, ArH).
- 21) Compound 11 was prepared in 26% yield by treatment of 5 with sodium hydride and diiodomethane at RT in THF; colorless plates from hexane-benzene, mp 124-125.5 °C.
- 22) Compound 12 was prepared in 95% yield from Wadsworth-Emmons reaction of 7 and acetone; colorless prisms from hexane, mp 106 °C, IR(KBr) 1595 cm⁻¹(C=C).

(Received February 7, 1986)