


A Novel Deselenation in the Reaction of Selenoamides with Organolithium Reagents

Masahito SEKIGUCHI, Akiya OGAWA,* Shin-Ichi FUJIWARA, Ilhyong RYU,
Nobuaki KAMBE, and Noboru SONODA*Department of Applied Chemistry, Faculty of Engineering,
Osaka University, Suita, Osaka 565

The reaction of selenoamides with organolithium reagents proceeds in a carbophilic manner, and more interestingly, the presence of excess lithium reagents causes a novel deselenation reaction from the carbophilic adducts.

Although much effort is being devoted to the preparation of selenocarbonyl compounds and to make a comparison with their sulfur analogues,¹⁾ reports on the fundamental reactivities of selenocarbonyl compounds are still rare. For example, thiophilic addition has been well-documented to take place in the reaction of thiocarbonyl compounds with various nucleophiles,²⁾ but in contrast, only very limited information is currently available on the reactivity of selenocarbonyl compounds toward nucleophiles. Recently, Okazaki et al. reported that the heterophilic addition of organolithium and organomagnesium reagents is much more significant in the reaction with selenoketone than with thioketone.^{3,4)} In this letter, we report that the reaction of selenoamides, as heteroatom-substituted selenocarbonyl compounds,⁵⁾ with organolithium reagents proceeds in a carbophilic manner, and more interestingly, that the presence of excess lithium reagents causes a novel deselenation reaction from the carbophilic adducts.

To a stirred THF (15 mL) solution of 1-selenobenzoylpiperidine (**1a**, 3 mmol) was added dropwise 2.2 equivalents of phenyllithium over 20-min period at 0 °C. After 100 min, the reaction mixture was quenched with water and extracted with diethyl ether (3 x 30 mL). The combined extracts were dried over MgSO₄, and the solvent was removed *in vacuo*. Flash chromatography on silica gel (eluting with hexane/Et₂O = 10/1) provided 1-benzhydrylpiperidine (**2a**) in 94% yield along with diphenyl diselenide (73%) (Eq. 1).




a: Ar = C ₆ H ₅ , NR ₂ = piperidino	94%	73%
b: Ar = <i>p</i> -MeC ₆ H ₄ , NR ₂ = piperidino	98%	91%
c: Ar = C ₆ H ₅ , NR ₂ = 1-pyrrolidiny	95%	91%
d: Ar = C ₆ H ₅ , NR ₂ = dimethylamino	82%	90%


Some other selenoamides bearing no hydrogen at α to the selenocarbonyl underwent a similar deselenation with 2.2 equivalents of PhLi. In the case using equimolar amounts of PhLi, amine **2** was not formed at all.⁶⁾

To gain insight into the reaction course of this deselenation, we attempted to follow the reaction of selenoamide **1a** with PhLi by taking the ^{13}C and ^{77}Se NMR (Fig. 1). The solution of **1a** in THF showed characteristic peaks at δ 202.9 in ^{13}C NMR and δ 745 in ^{77}Se NMR, which are assigned to the selenocarbonyl carbon and selenium, respectively. On treatment of the solution with equimolar amounts of PhLi at -50°C ,⁷⁾ the peaks immediately disappeared and new peaks appeared at δ 81.6 in ^{13}C NMR⁸⁾ and δ -7 in ^{77}Se NMR, which apparently indicates the clean formation of carbophilic adduct **3**. Addition of another one equivalent of PhLi to the solution at 0°C caused disappearance of the signals of carbophilic adduct **3**, and appearance of a peak at δ 64 in ^{77}Se NMR, which can be assigned to lithium benzeneselenolate.

1) Selenoamide **1a**^{a)}



2) Selenoamide **1a** + PhLi (1.1 equiv.)^{b)}



3) Selenoamide **1a** + PhLi (2.2 equiv.)^{c)}



4) Characteristic signals of carbophilic adducts (**3**, **3'**, **3''**) in ^{13}C NMR spectra^{d)}





Fig. 1. ^{77}Se NMR Spectra in THF at -40°C measured with MeSeMe (neat) or PhSeSePh (0.5 M CDCl_3 solution) as an external standard. a) THF solution of selenoamide **1a** (0.8 mmol, 0.53 mol dm^{-3}). b) ^{77}Se NMR spectrum obtained on addition of PhLi (0.88 mmol) to the solution of **1a** in THF. c) Another 1.1 equiv. of PhLi was added to the solution including carbophilic adduct **3**. d) Adducts were prepared *in situ* by the reaction of selenoamide, thioamide, and amide with PhLi (1.1 equiv.).

A possible reaction path is shown in Scheme 1: The reaction of selenoamide **1** with PhLi leads to formation of carbophilic adduct **3**, which undergoes deselenation⁹⁾ by the action of another one equivalent of PhLi to form benzylic lithium **4** and lithium benzeneselenolate. The formation of diphenyl diselenide is



explained by hydrolysis of PhSeLi and the subsequent air-oxidation of benzeneselenol during workups.



Scheme 1. A possible reaction path.

In sharp contrast, the corresponding amide and thioamide did not undergo deoxygenation and desulfurization by a similar treatment with excess PhLi (2.2 equiv.), and instead, benzophenone and thiobenzophenone were formed in high yields, respectively (> 80%, GLC yield).¹⁰⁾ The measurement of ¹³C NMR spectra of the solution of amide or thioamide with one equivalent of PhLi showed the formation of carbophilic adducts 3' and 3'' in analogy with selenoamide (Fig. 1). However, thus formed carbophilic adducts 3' and 3'' did not show any change on the spectrum of ¹³C NMR by treating with excess PhLi. Presumably, weaker bond energy of C-Se may partly contribute to occurrence of deselenation.¹¹⁾

The reactions of selenoamide with some alkylolithiums (2.2 equiv.) were also carried out (Eqs. 2-4). In the case using *t*-BuLi, a similar deselenation *via* carbophilic adduct took place as well as the case using PhLi. Contrary to this, the reaction with MeLi gave 1-phenyl-1-piperidinoethene (5a), the formation of which is understandable as resulting from the elimination of selenolate anion (Li₂Se or LiSeH) from the carbophilic adducts. In the reaction with *n*- and *s*-BuLi, deselenation to benzylic amines 2a'' and 2a''' was in competition with elimination to enamines 5b and 5c.



Moreover, the reaction of aliphatic selenoamide **1e** with PhLi (2.2 equiv.), followed by alkylation with methyl iodide was found to provide α -(methylseleno) enamine **6** and α -phenyl enamine **5d**; this suggests that the α -hydrogen abstraction from selenoamide took place in preference to carbophilic attack.

In conclusion, the reaction of aromatic selenoamides with organolithiums predominantly provided carbophilic adducts, which underwent novel deselenation on treatment of excess lithium reagents unlike related amides and thioamides. Further investigation on precise mechanism of the deselenation and its possible application to organic synthesis are under way.

This research was supported by Grant in Aid for Scientific Research from the Ministry of Education, Science and Culture, Japan. Thanks are due to the Instrumental Analysis Center, Faculty of Engineering, Osaka University for assistance in obtaining NMR and Mass spectra on a Bruker AM-600 and a JEOL JMS-DX303.

References

- 1) For reviews, see: F. S. Guziec, Jr., "The Chemistry of Organic Selenium and Tellurium Compounds," ed by S. Patai, John Wiley & Sons, New York (1987), Vol. 2, Chap. 4; C. Paulmier, "Selenium Reagents and Intermediates in Organic Synthesis," Pergamon Press, Oxford (1986), Chap. 3.
- 2) P. Beak and J. W. Worley, *J. Am. Chem. Soc.*, **94**, 597 (1972); M. Dagonneau and J. Vialle, *Tetrahedron*, **30**, 3119 (1974); A. Ohno, K. Nakamura, Y. Shizume, and S. Oka, *Bull. Chem. Soc. Jpn.*, **50**, 1003 (1977) and references cited therein.
- 3) Reaction of 1,1,3,3-tetramethyl-2-indanselone with some Grignard and organolithium reagents was reported: R. Okazaki, A. Ishii, and N. Inamoto, *Tetrahedron Lett.*, **25**, 5147 (1984); A. Ishii, R. Okazaki, and N. Inamoto, *Bull. Chem. Soc. Jpn.*, **61**, 861 (1988).
- 4) It was reported that di-*t*-butyl selone reacted with *n*-BuLi to give predominantly selenophilic adduct along with small amounts of reduction product: F. S. Guziec, Jr., L. J. SanFilippo, C. J. Murphy, C. A. Moustakis, and E. R. Cullen, *Tetrahedron*, **41**, 4843 (1985).
- 5) In order to prepare selenoaldehyde, reaction of selenoformate with 2,4,6-tri-*t*-butylphenyllithium was examined: R. Okazaki, A. Ishii, and N. Inamoto, *J. Chem. Soc., Chem. Commun.*, **1986**, 71; A. Ishii, R. Okazaki, and N. Inamoto, *Bull. Chem. Soc. Jpn.*, **60**, 1037 (1987).
- 6) The reaction of selenoamide **1a** with one equivalent of PhLi provided benzophenone (42%) and dibenzhydryl diselenide (40%) after aqueous workup. It was confirmed that hydrolysis of carbophilic adduct **3** led to benzophenone (40%) and dibenzhydryl diselenide (41%).
- 7) At 0 °C, the carbophilic adduct **3** gradually decomposed.
- 8) ¹³C NMR (THF-d₈) for **3**: δ 26.5 (N-CH₂-CH₂-CH₂), 27.7 (N-CH₂-CH₂), 51.4 (N-CH₂), 81.6 (Se-C-N), 124.5 (p), 126.5 (m), 130.6 (o), 154.3 (ipso).
- 9) If the carbophilic adduct **3** is to decompose to piperidinomethylithium **4** and elemental selenium, PhLi may trap the selenium to give PhSeLi.
- 10) Reaction of thioamides with some nucleophiles, see: P. Beak, J. Yamamoto, and C. J. Upton, *J. Org. Chem.*, **40**, 3052 (1975); W. Walter and H. -W. Lüke, *Angew. Chem., Int. Ed. Engl.*, **16**, 535 (1977); T. Karakasa, T. Hanzawa, and S. Motoki, *Bull. Chem. Soc. Jpn.*, **52**, 3469 (1979); Y. Tominaga, S. Kohra, and A. Hosomi, *Tetrahedron Lett.*, **28**, 1529 (1987).
- 11) Bond dissociation energies *D* [kJ mol⁻¹] for diethyl chalcogenides are as follows: *D* (Se-C) = 251; *D* (S-C) = 292; *D* (O-C) = 362, see: L. Batt, "The Chemistry of Organic Selenium and Tellurium Compounds," ed by S. Patai and Z. Rappoport, John Wiley & Sons, New York (1986), Vol. 1, Chap. 4.

(Received August 9, 1990)