


## Reactions of syn- and anti-Benzylideneaniline Radical Anions

Akito ISHIDA,\* Hiroshi ITOH, and Setsuo TAKAMUKU  
 The Institute of Scientific and Industrial Research,  
 Osaka University, Ibaraki, Osaka 567

Reactions of a benzylideneaniline radical anion have been studied by spectral and product analyses, in which successive irradiation of uv-light and ionizing radiation at low temperature have been applied to the production of the unstable syn-radical anions.

Imine is a very useful intermediate in the creation of valuable compounds. Even though a radical anion of imine is an important intermediate in electrochemical and photochemical reactions, it has not been studied by time-resolved spectroscopic methods. In this paper, we report one electron reduction of benzylideneaniline and its derivatives (**1**) studied by the combination use of photochemical and radiation chemical methods, which enables to produce very unstable syn-**1** radical anions.



Photoirradiation of 0.5 mmol dm<sup>-3</sup> anti-**1a** in a 77 K MTHF (2-methyltetrahydrofuran) matrix by use of a 150 W Xe-lamp for 5 minutes resulted in a blue shift of the UV absorption maximum of **1a** from 325 nm to 250 nm, and upon melting of the matrix the original spectrum was recovered. Other derivatives also gave a similar spectral change as shown in Table 1. This spectral change was explained in terms of a photoinduced isomerization of anti-**1** to syn-**1** and a thermal inverse reaction.<sup>1)</sup>

Radical anion of anti-**1** was produced by  $\gamma$ -ray irradiation of 0.5 mmol dm<sup>-3</sup> anti-**1** in a 77 K MTHF matrix.<sup>2)</sup> An absorption maximum of **1a**<sup>-•</sup> was observed at 455 nm (Fig. 1; curve A) and agreed with that reported by Shida et al.<sup>3)</sup> On the other hand,  $\gamma$ -ray irradiation of syn-**1a** isolated in a 77 K MTHF matrix by photoirradiation provided a different absorption spectrum. After  $\gamma$ -ray irradiation at 77 K, it has an absorption maximum at 455 nm with a shoulder around 485 nm (Fig. 1; curve B). Slight annealing of the matrix induced rapid decrease of the band at 485 nm followed by a simultaneous rise of the band at 455 nm with an isosbestic point at 475 nm (Fig. 1; curve C). A newly observed

absorption maximum around 485 nm was assigned to syn-**1a** radical anion. Other derivatives also gave absorption spectra of syn-radical anions and the absorption maxima are summarized in Table 1. A pulse radiolytic measurement was carried out by use of photoproducted syn-**1g** at 120 K, in which a transient absorption of syn-**1g** radical anion and its time dependent isomerization ( $k_i=3.5 \times 10^6 \text{ s}^{-1}$ ) were clearly observed (Fig. 2). From these results, it is clear that syn-**1** radical anion is also very unstable and rapidly isomerizes to anti-**1** radical anion.




Fig. 1. An absorption spectrum observed upon  $\gamma$ -ray irradiation of  $0.5 \text{ mmol dm}^{-3}$  **1a** in MTHF at 77 K (curve-A); absorption spectra observed upon photoirradiation and successive  $\gamma$ -ray irradiation of  $0.5 \text{ mmol dm}^{-3}$  **1a** in MTHF at 77 K (curve-B) and after annealing to 83 K (curve-C).




Fig. 2. Transient absorption spectra observed upon electron beam (28 MeV, 8 ns) irradiation of  $5 \text{ mmol dm}^{-3}$  **1g** in MTHF which had been photoirradiated at 120 K by use of a 450 W Xe lamp through a  $\text{H}_2\text{O}$  filter; insets show time dependent profiles at 465 nm and 515 nm.

Table 1. Absorption Maxima for Benzilideneanilines (**1**) and Their Radical Anions<sup>a)</sup>

| Substrate | R   | R'  | $\lambda_{\text{max}}/\text{nm}$ |                   |                                     |                   |
|-----------|-----|-----|----------------------------------|-------------------|-------------------------------------|-------------------|
|           |     |     | neutral<br>anti                  | syn <sup>c)</sup> | radical anion <sup>b)</sup><br>anti | syn <sup>d)</sup> |
| <b>1a</b> | H   | H   | 325                              | 250               | 455                                 | 485, 510          |
| <b>1b</b> | MeO | H   | 295                              | 275               | 450                                 | 480 (broad)       |
| <b>1c</b> | Me  | H   | 280                              | 260               | 455                                 | 490 (broad)       |
| <b>1d</b> | CN  | H   | 295                              | 265               | 490                                 | 530 (broad)       |
| <b>1e</b> | H   | MeO | 330                              | 250               | 455                                 | 500               |
| <b>1f</b> | H   | Me  | 330                              | 250               | 455                                 | 500               |
| <b>1g</b> | H   | CN  | 295                              | 275               | 470                                 | 520               |

a) Measurement conditions;  $[\mathbf{1}]=0.5 \text{ mmol dm}^{-3}$ , MTHF, 77 K.

b)  $\gamma$ -Ray irradiation was carried out at 77 K by a  $^{60}\text{Co}$  source for 15 min.

c) The syn isomers were produced by photoirradiation of the anti isomers at 77 K by a 150 W Xe lamp through a  $\text{H}_2\text{O}$  filter for 5 min.

d) The syn-radical anions were produced by successive  $\gamma$ -ray irradiation after the photoirradiation.




Fig. 3. Transient absorption spectra observed upon electron beam (28 MeV, 8 ns) irradiation of 5 mmol dm<sup>-3</sup> anti-**1a** in THF at room temperature; insets show time dependent profiles at 370 nm and 455 nm.





Fig. 4. Relationships between  $k_{\text{EtOH}}$ 's observed for a series of anti-**1** (4-substituted;  $\square$ , 4'-substituted;  $\bullet$ ) and Hammett  $\sigma_p$  values for the substituents.

Reactions of anti-**1** radical anions were studied by pulse radiolysis. Fig. 3 shows transient absorption spectra obtained by pulse radiolysis of 5 mmol dm<sup>-3</sup> anti-**1a** in THF at room temperature. The initially observed absorption band with a maximum at 455 nm is the radical anion of anti-**1a**, decrease of which leads to another absorption band with a maximum at 370 nm. In EtOH, this absorption band at 370 nm was observed even immediately after pulse irradiation (<20 ns). Rapid protonation of anti-**1a** radical anion was supposed to induce the formation of a neutral radical, which was confirmed by kinetic measurements. Pseud-first-order decay rates of the radical anion were observed in a series of THF-EtOH mixture solvents, which depended on the concentration of EtOH; the second-order plot gave a straight line, from which a reaction rate constant of the radical anion and EtOH,  $k_{\text{EtOH}}$  was obtained. The rate constant strongly depends on the substituent. A Hammett plot of  $k_{\text{EtOH}}$  against  $\sigma_p$  of R' gives almost linear relationship with a negative slope as shown in Fig. 4. This result suggests that a protonation site of the radical anion is the N atom, the electron density of which depends on polar effect of R'. A MNDOC calculation also supports this estimation.

In order to identify the structure of the radical, a product analysis was carried out.  $\gamma$ -Ray irradiation of anti-**1a** in THF at room temperature afforded a THF adduct, N-[(2-tetrahydrofuranyl)-benzyl]-aniline (**2a**) in a yield of 47% (conv. 57%,  $G=2.54$ ),<sup>5)</sup> which is supposed to be derived from the benzyl-type radical (**3a**).<sup>6)</sup> In THF, the proton source is supposed to be a radical cation of THF. A total reaction scheme of the syn- and anti-radical anions is summarized as follows.



This work was supported by a grant from the Research Program on "Creation of New Materials through Intelligent Design" of ISIR, Osaka University.

#### References

- 1) M. Kobayashi, M. Yoshida, and H. Minato, *Chem. Lett.*, **1976**, 185; D. G. Anderson and G. Wettermark, *J. Am. Chem. Soc.*, **87**, 1433 (1965); E. Fisher and Y. Frei, *J. Chem. Phys.*, **27**, 808 (1957).
- 2) Benzyldeneaniline (**1a**) was commercially available and purified by column chromatography. Other derivatives were synthesized by coupling of respective aniline and benzaldehyde by a reported procedure.<sup>7)</sup> MTHF was stirred with NaOH pellets for removing stabilizer and distilled over  $\text{LiAlH}_4$ . For spectral measurement, a sample solution was degassed by freeze-pump-thaw cycles and sealed in a Suprasil cell. Steady state  $\gamma$ -ray irradiation was carried out with a  $^{60}\text{Co}$  source ( $2.6 \times 10^{14}$  Bq, dose rate  $6.3 \times 10^2$  Gy/h). Absorption spectra of low temperature matrices were measured with a multichannel spectrophotometer (Otsuka Electronix, MCPD-1100) and a quartz Dewar vessel. Pulse radiolysis was carried out by electron beam (28 MeV, 8 ns) from ISIR linac. Low temperature irradiation was carried out in a cryostat (Oxford D-1704).
- 3) T. Shida, "Electronic Absorption Spectra of Radical Ions," Elsevier, New York (1988), p. 192.
- 4) G value is a molecular amount at energy absorption of 100 eV.
- 5) **2a**; colorless oil; MS  $m/e$  (relative intensity); 253 ( $\text{M}^+$ , 7), 182 (100), 104 (11), 91 (3), 77 (15), 51 (3);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ =1.7-1.9 (m, 4H), 3.90 (ddd, 1H,  $J=7.0$  Hz), 4.05 (ddd, 1H,  $J=6.3$  Hz,  $J=7.0$  Hz), 4.23 (ddd, 1H,  $J=3.9$  Hz), 4.42 (d,  $J=3.9$  Hz, 1H), 6.5-7.5 (m, 10H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$ =25.45, 25.63, 25.78 (t,  $-\text{CH}_2-$ ), 27.18, 28.73 (t,  $-\text{CH}_2-$ ), 60.72, 61.96, 62.02 (d,  $\text{Ph}-\text{CH}$ ), 68.50, 68.68 (t,  $-\text{CH}_2-\text{O}-$ ), 81.97, 82.93 (d,  $-\text{CH}-\text{O}$ ), 113.77-114.08 (d, Ph), 117.34-117.46 (d, Ph), 127.19-127.68 (d, Ph), 127.92-129.18 (d, Ph), 139.99 (s, Ph), 141.62 (s, Ph), 147.34 (s, Ph), 147.34 (s, Ph), 147.77 (s, Ph).
- 6) G. Porter and E. Strachan, *Trans. Faraday Soc.*, **54**, 1595 (1958).
- 7) J. A. Castellano, J. E. Goldmacher, L. A. Barton, and J. S. Kane, *J. Org. Chem.*, **33**, 3501 (1968).

(Received June 6, 1991)