

The Catalytic Beckmann Rearrangement of
Ketoxime Trimethylsilyl Ethers Using an Antimony(V) Salt

Teruaki MUKAIYAMA and Tsunehiro HARADA

Department of Applied Chemistry, Faculty of Science,
Science University of Tokyo, Kagurazaka, Shinjuku-ku, Tokyo 162


In the presence of a catalytic amount of an antimony(V) salt generated from antimony(v) chloride and silver antimonate, the Beckmann rearrangement of several ketoxime trimethylsilyl ethers proceeds smoothly to give the corresponding amides or lactams in good yields.

The Beckmann rearrangement is a fundamental and useful reaction, long recognized as an extremely valuable and versatile method for the preparation of amides or lactams, and often employed even in the industrial processes.¹⁾ However, the reaction generally requires at least an equimolar amount of activators such as strong acids including Lewis acids.^{1,2)} In the vapor phase process, there have been reported a few examples that were promoted by a catalytic amount of activators such as boria-hydroxyapatite³⁾ under rather drastic conditions. Therefore, it was strongly desired to explore an efficient catalyst which could promote the Beckmann rearrangement under mild conditions. Recently, we have found that a catalytic amount of tin(IV) species, easily prepared in situ from SnCl_4 and AgClO_4 , realizes the highly stereoselective glycosylation reaction between a simple glycosyl donor, 1-O-acetyl glucose, and silyl alkoxides, and also gadolinium(III) species for the catalytic Friedel-Crafts acylation.⁴⁾ In this communication, we would like to describe the Beckmann rearrangement performed under mild condition using a catalytic amount of such active species as an antimony(v) salt.

First, the Beckmann rearrangement of benzophenone oxime derivatives was tried by taking tin(IV) species generated from SnCl_4 and AgSbF_6 as a

catalyst. As shown in Table 1, only in the case of trimethylsilyl ether, the reaction did proceed smoothly in CH_2Cl_2 at refluxing temperature to give the corresponding amide in good yield.


Table 1. The Beckmann Rearrangement of Benzophenone Oxime Derivatives



Entry	R	Temperature	Time/h	Yield/%
1	H	reflux	7	trace
2	Ac	reflux	7	48
3	TMS	rt	overnight	47
4	TMS	reflux	7	71
5	Me	reflux	7	trace


Next, several Lewis acids and solvents were examined employing the above mentioned reaction as a model. As shown in Table 2, neither TiCl_4 nor SnCl_4 alone did catalyze the reaction, and nor AgSbF_6 did either. On the contrary, the reaction proceeded smoothly to give the corresponding amide in 71-78% yields when active species generated from SnCl_4 or SbCl_5 and AgSbF_6 was employed respectively. A better result was obtained in the case of the combined use of SbCl_5 and AgSbF_6 .

Table 2. Effect of Catalyst



Entry	Catalyst(20 mol%)	Yield/%
1	$\text{SnCl}_4 + \text{AgSbF}_6$	71
2	$\text{TiCl}_4 + \text{AgSbF}_6$	24
3	$\text{AlCl}_3 + \text{AgSbF}_6$	49
4	$\text{SbCl}_5 + \text{AgSbF}_6$	78
5	TiCl_4	2
6	SnCl_4	no reaction
7	AgSbF_6	19

Table 3. Effect of Solvent



Entry	Solvent	Reaction time/h	Yield/%
1	CH_2Cl_2	7	78
2	Et_2O	7	53
3	$\text{CH}_2\text{ClCH}_2\text{Cl}$	3	98
4	CH_3CN	3	91
5	C_6H_6	3	87

Several examples of the present Beckmann rearrangement are demonstrated in Table 4. In every case, the reaction proceeds smoothly at refluxing

temperature in CH_3CN to give the corresponding amides or lactams in good yields. In the case of benzophenone oxime trimethylsilyl ether, 1,2-dichloroethane was shown to be an effective solvent, while, in the case of other oxime trimethylsilyl ethers, CH_3CN was the best solvent.

Table 4. The Beckmann Rearrangement of Several Ketoxime Trimethylsilyl Ethers^{a)}


Entry	Substrate	Product	Solvent	Temperature	Time/h	Yield/%
1		PhNHCOPh	$\text{CH}_2\text{ClCH}_2\text{Cl}$	reflux	3	98
2		PhNHCOMe	CH_3CN	reflux	3	83
3		$\text{PhNHCOCH}_2\text{Ph}$	CH_3CN	reflux	3	83
4		$\text{PhCH}_2\text{NHCOCH}_2\text{Ph}$	CH_3CN	reflux	3	74
5		EtNHCOEt	CH_3CN	reflux	3	50
6			CH_3CN	40 °C	6	65
7			CH_3CN	reflux	3	77

a) Catalyst: $\text{SbCl}_5 + \text{AgSbF}_6$ (20 mol%).

The catalytic cycle of this rearrangement is postulated as shown in Scheme 1. The antimony(V) species activates a trimethylsilyloxy group of oxime moiety, followed by rearrangement of R anti to the trimethylsilyloxy function to form the cation stabilized by antimonate anion (**3**) and CH_3CN . Then the ion pair (**3**) reacts with initially formed

tetrachloro(trimethylsilyloxy)antimony(**2**) to give the desired imino trimethylsilyl ether(**4**) along with a regeneration of the catalyst(**1**).

A typical experimental procedure for the Beckmann rearrangement of benzophenone oxime trimethylsilyl ether is as follows; a CH_2Cl_2 solution of 0.5 M antimony(v) chloride(0.12 ml) was added to a silver hexafluoroantimonate(20.6 mg, 0.06 mmol) solution of 1,2-dichloroethane under ice cooling with the protection from the light. After stirring for 1 h, a 1,2-dichloroethane(1.5 ml) solution of benzophenone oxime trimethylsilyl ether(80.8 mg, 0.3 mmol) was added at rt. After refluxing for 3 h, the mixture was quenched with aqueous NaHCO_3 . The organic layer was dried over Na_2SO_4 . After the evaporation of the solvents, the residue was purified by preparative TLC to afford 58.2 mg(98%) of benzanilide, mp 163-165 °C.



Scheme 1. The catalytic cycle.

References

- 1) R. E. Gawley, *Org. React.*, **35**, 1(1988), and references cited therein.
- 2) J. C. Jochims, S. Hehl, and S. Herzberger, *Synthesis*, **1990**, 1128, and references cited therein.
- 3) Y. Izumi, S. Sato, and K. Urabe, *Chem. Lett.*, **1983**, 1649, and references cited therein.
- 4) T. Mukaiyama, M. Katsurada, and T. Takashima, *Chem. Lett.*, **1991**, 985; T. Mukaiyama, T. Ohno, T. Nishimura, S. Suda, and S. Kobayashi, *ibid.*, **1991**, 1059, and references cited therein.

(Received July 5, 1991)