


BuSnCl₃ Mediated *Z*-Selective 2-Butenylation and *erythro*-Selective 1-Methyl-2-propenylation
of Aldehydes by 1-(Tributylstannyl)-2-butene

Hideyoshi MIYAKE* and Kimiaki YAMAMURA

Department of Chemistry, College of General Education, Kobe University, Nada, Kobe 657

Butyltin trichloride mediated reaction of 1-(tributylstannyl)-2-butene with aldehydes is described. This reaction is useful as a stereoselective method for (*Z*)-2-butenylation and *erythro*-1-methy-2-propenylation of aldehydes.


The utility of 1-(tributylstannyl)-2-butene(**1**) is well-established.¹⁾ Lewis acid catalyzed reaction and thermal reaction of **1** with aldehydes give 1-methyl-2-propenylated products.²⁾ On the other hand, 2-butenylation by **1** can be accomplished by a transmetalation method.^{3,4)} For example, Bu₂SnCl₂ mediated reaction of **1** with aldehydes proceeds regio- and stereoselectively to give (*Z*)-2-butenylated products predominantly.⁴⁾ This reaction contains 3-(dibutylchlorostannyl)-1-butene(**2**) as an intermediate. Although 3-(butyldichlorostannyl)-1-butene(**3**), prepared from **1** and BuSnCl₃ (**4**), is more reactive than **2**,⁵⁾ the utility of **3** in organic synthesis has not been studied extensively. In this paper, we wish to report the **4** mediated stereoselective (*Z*)-2-butenylation and *erythro*-1-methyl-2-propenylation of aldehydes by **1**.





Regio- and stereoselectivity of **4** mediated reaction of **1** with aldehydes depends on the stereochemistry of **1** and the reaction conditions. Especially, the order of the addition of **1**, **4**, and aldehydes is important. When (*Z*)-1-(tributylstannyl)-2-butene(**(Z)-1**)⁶⁾ was used and **4** was added last, (*Z*)-2-butenylated product(**6**) was obtained predominantly (Method A).⁷⁾ In this method, the initially formed **3** reacts immediately with aldehydes without isomerization to 1-(butyldichlorostannyl)-2-butene(**5**) to give **6**.

When **4** was added to a solution of **1**, the transmetalation of Bu₃Sn to BuSnCl₂ proceeded immediately to give **3** and **5**, and the former slowly isomerized to **5**. When aldehyde was added after sufficient time elapsed for the isomerization of **3**, 1-methy-2-propenylated product(**7**) was obtained predominantly (Method B,⁸⁾ Runs 8-17 in Table 1).

When *E,Z* mixture of **1** was used in Method A, a considerable amount of **7** was obtained (Runs 2 and 5). Similar results were obtained in Method B with short reaction time at low temperature (Runs 6 and 7). When **4** was added to the solution of **1**, transmetalation of Bu_3Sn to BuSnCl_2 proceeded immediately even at -78°C . These results suggest that the transmetalation of (*Z*)-**1** proceeds with migration of the double bond, and that of (*E*)-**1** proceeds without it (Scheme 1). The reasons for these results are under investigation.




The *Z*-selectivity of the reaction of **3** with aldehydes can be explained by the well-established chairlike cyclic transition state.⁴⁾ (Scheme 2)



In Method B, the ratios of **6**, **7**(*erythro*), and **7**(*threo*) depend on the reaction time and temperature. When CH_2Cl_2 was used as solvent, the reaction at 0°C for 8 hs gave the best *erythro* selectivity. Longer reaction time and higher temperature increase the ratio of *threo* isomer. These results can be explained as follows. When the initially formed **3** slowly isomerizes to **5**, isomerization to (*Z*)-**5** is faster than that to (*E*)-**5**, because of the steric effect shown in Scheme 3.⁹⁾ However, (*E*)-**5** is thermodynamically more stable than (*Z*)-**5**. The ratio of (*E*)-**5** slowly increases at higher temperature. (*Z*)-**5** is the kinetically controlled product, and (*E*)-**5** is the thermodynamically controlled product. (*Z*)-**5** and (*E*)-**5** form the chairlike transition state shown in Scheme 4 by the reaction with benzaldehyde to give **7**(*erythro*) and **7**(*threo*), respectively.¹⁰⁾

Table 1. BuSnCl₃ Mediated Reaction of **1** with Aldehydes


Run	<i>E/Z</i> ratio of 1	R	Method	Solvent	Temp/°C, Time	Yield/%	6	: 7 (<i>erythro</i>)	: 7 (<i>threo</i>)
1	0 / 10	Ph	A	CH ₂ Cl ₂	-78	96	96	: 3	: 1
2	6 / 4	Ph	A	CH ₂ Cl ₂	-78	96	44	: 34	: 22
3	0 / 10	Ph	A	CH ₂ Cl ₂	0	95	80	: 16	: 4
4	0 / 10	Hex	A	CH ₂ Cl ₂	-78	94	95	: 5 (<i>threo</i> + <i>erythro</i>)	
5	6 / 4	Hex	A	CH ₂ Cl ₂	-78	90	69	: 31 (<i>threo</i> + <i>erythro</i>)	
6	0 / 10	Ph	B	CH ₂ Cl ₂	-10, 1 min	92	96	: 3	: 1
7	6 / 4	Ph	B	CH ₂ Cl ₂	-10, 1 min	95	54	: 23	: 23
8	0 / 10	Ph	B	CH ₂ Cl ₂	0, 8 h	93	6	: 80	: 14
9	6 / 4	Ph	B	CH ₂ Cl ₂	0, 8 h	68	trace	: 69	: 31
10	0 / 10	Ph	B	CH ₂ Cl ₂	0, 15 h	75	4	: 74	: 22
11	6 / 4	Ph	B	CH ₂ Cl ₂	0, 15 h	80	0	: 69	: 31
12	0 / 10	Ph	B	CH ₂ Cl ₂	reflux, 30 min	93	trace	: 75	: 25
13	0 / 10	Ph	B	CHCl ₃	0, 16 h	86	0	: 82	: 18
14	0 / 10	Ph	B	CHCl ₃	30, 3 h	90	4	: 65	: 31
15	0 / 10	Ph	B	CHCl ₃	reflux, 1 h	95	trace	: 50	: 50
16	0 / 10	Ph	B	toluene	reflux, 1 h	65	trace	: 48	: 52
17	6 / 4	Ph	B	CHCl ₃	0, 14 h	80	trace	: 77	: 23



Scheme 4.

References

- 1) M. Peryre, J. -P. Quinterd, and A. Rahm, "Tin in Organic Synthesis," Butterworth, London (1987); Y. Yamamoto, *Acc. Chem. Res.*, **20**, 243 (1987).
- 2) Y. Naruta, S. Ushida, and K. Maruyama, *Chem. Lett.*, **1979**, 919; H. Yatagai, Y. Yamamoto, and K. Maruyama, *J. Am. Chem. Soc.*, **102**, 4548 (1980); Y. Yamamoto, H. Yatagai, Y. Naruta, and K. Maruyama, *ibid.*, **102**, 7107 (1980); B. W. Gung and D.T. Smith, *Tetrahedron Lett.*, **32**, 13 (1991); B. W. Gung, A. J. Peat, B. M. Snook, and D. T. Smith, *ibid.*, **32**, 453 (1991).
- 3) Y. Yamamoto, N. Maeda, and K. Maruyama, *J. Chem. Soc., Chem. Commun.*, **1983**, 742; J. Iqbal, and S. P. Joseph, *Tetrahedron Lett.*, **30**, 2421 (1989); S. Matsubara, K. Wakamatsu, Y. Morizawa, N. Tsuboniwa, K. Oshima, and H. Nozaki, *Bull. Chem. Soc. Jpn.*, **58**, 1196 (1985).
- 4) A. Gambaro, P. Ganis, D. Marton, V. Peruzzo, and G. Tagliavini, *J. Organomet. Chem.*, **231**, 307 (1982).
- 5) A. Gambaro, V. Peruzzo, G. Plazzogna, and G. Tagliavini, *J. Organomet. Chem.*, **197**, 45 (1980).
- 6) Typical procedure of Method A (Run 1 in Table 1) is as follows: To a solution of **1** (0.690 g, 2.0 mmol) and benzaldehyde (0.233 g, 2.2 mmol) in CH_2Cl_2 (10 ml) under nitrogen atmosphere at -78°C , was added **4** (0.564 g, 2.0 mmol) in CH_2Cl_2 (5 ml) slowly. The mixture was allowed to warm to 0°C , and stirred at that temperature for 30 min. The reaction mixture was quenched with H_2O and extracted with ether. The organic phase was dried, condensed, and purified by column chromatography on silica gel to give the mixture of **6a**, **7a** (*erythro*), and **7a** (*threo*) (total 0.302 g, 1.84 mmol, 92% yield). The ratio of **6a** : **7a** (*erythro*) : **7b** (*threo*) was determined by ^1H NMR.
- 7) Typical procedure of Method B (Run 8 in Table 1) is as follows: To a solution of **1** (0.690 g, 2.0 mmol) in CH_2Cl_2 (10 ml) at -10°C under nitrogen atmosphere, was added **4** (0.564 g, 2.0 mmol) in CH_2Cl_2 (5 ml) slowly. After stirring at 0°C for 8 h, benzaldehyde (0.254 g, 2.4 mmol) in CH_2Cl_2 (2 ml) was added slowly to the mixture. After stirring at 0°C for 30 minutes, the reaction mixture was quenched and purified in a similar manner to Method A. The total yield of **6a**, **7a** (*erythro*), and **7a** (*threo*) was 95% (0.301g, 0.186 mmol).
- 8) Stereoselective synthesis of (*Z*)-(1); H. Miyake and K. Yamamura, *Chem. Lett.*, **1992**, 507.
- 9) ^{13}C NMR data of (*E*)-3-(tributylstannyl)-2-pentene suggest its most stable conformation to be **8**. This result supports the most stable conformation of **3** to be **9**: H. Miyake and K. Yamamura, *Chem. Lett.*, **1992**, in press.



- 10) Y. Masuyama, J. P. Takahara, and Y. Kurusu, *Tetrahedron Lett.*, **30**, 3437 (1989); A. J. Pratt and E. J. Thomas, *J. Chem. Soc., Perkin Trans. 1*, **1989**, 1521; Y. Yamamoto and K-i. Saito, *J. Chem. Soc., Chem. Commun.*, **1989**, 1676.

(Received March 18, 1992)