

The Simple Preparation of Aldehydes and Ketones by the Photo-Oxygen
Rearrangement of Naphtho[1,8-*de*]dithiin Monooxides

Naomichi FURUKAWA,* Takayoshi FUJII, Takeshi KIMURA, and Hisashi FUJIHARA
Department of Chemistry, University of Tsukuba, Tsukuba, Ibaraki 305

Naphthalene-1,8-dithiol reacted with aldehydes to give dithioacetals which were oxidized to 2-substituted naphtho[1,8-*de*]1,3-dithiin-1-oxides (**4**). 2, 2-Disubstituted naphtho[1,8-*de*]1,3-dithiin-1-oxides (**5**) were obtained on treatment of **4** with NaH/electrophiles. Photolysis of **4** and **5** undergoes intramolecular oxygen rearrangement to generate aldehydes and ketones quantitatively together with naphthalene-1,8-dithiole.

In general, 1,3-dithianes and related derivatives have been utilized as carbonyl synthones¹⁾ in organic synthesis since they readily provide aldehydes and ketones by desulfurization with many efficient methods such as metal-induced,²⁾ oxidative,³⁾ alkylative,⁴⁾ photo-induced,⁵⁾ and electrolytic⁶⁾ decompositions or by hydrolysis of the corresponding monooxides with strong acids.⁷⁾

1,8-Dithia or -diselena naphthalene derivatives are presumably the candidates for generating various active species or creating new reactions via the formation of reactive dithia- or diselena dications.⁸⁾ In fact, 1,8-dithia or -diselena naphtho[1,8-*bc*]dithiocins undergo unusual Pummerer and thio-Claisen rearrangement reactions via the formation of dications.⁹⁾ Furthermore, quinodimethane is generated from photolysis of 8,13-dihydrobenzo[*g*]naphtho[1,8-*bc*][1,5]diselenonin.¹⁰⁾ As a further extension of these studies, we prepared naphtho[1,8-*de*]dithiins and found that photolysis of the corresponding monooxides became a convenient procedure to yield the carbonyl compounds. This paper reports the results on the photo-oxygen rearrangement reaction of naphtho[1,8-*de*]dithiin monooxides to the aldehydes and ketones.

1,8-Naphthalene dithiol (**1**) was treated with aldehydes in the presence of SiCl₄ in CH₂Cl₂ affording the corresponding 2-substituted naphtho[1,8-*de*]1,3-dithiins (**2**) in high yields.¹¹⁾ However, 2, 2-disubstituted naphtho[1,8-*de*]1,3-dithiins (**3**) were obtained in poor yields on analogous treatment with ketones and **1** and hence dithioketals **3** were prepared from the reaction of 2-lithiated **2** with several electrophiles. 2-Substituted naphtho[1,8-*de*]1,3-dithiin-1-oxides (**4**) and 2, 2-disubstituted naphtho[1,8-*de*]1,3-dithiin-1-oxides (**5**) were obtained in high yields with high diastereo-selectivities (for instance, diastereomeric ratios of **4a**, **4b**, and **4d** determined by ¹H NMR spectroscopy were found to be 5:1, 7:1, and 3:1 respectively) by oxidation using *m*-chloroperbenzoic acid (*m*CPBA) in CH₂Cl₂ at -20 °C.¹¹⁾ Furthermore, compounds **4** were also converted to dithioacetal monooxides **5** by the reaction with NaH in THF at 50 °C and then with several electrophiles in moderate yields.¹¹⁾ Although compounds **2** and **3** were thermally and photochemically stable molecules, monooxides **4** and **5** were found to decompose to the corresponding aldehydes and ketones quantitatively with complete recovery of naphthalene-1, 8-dithiole (**8**) on exposure to high pressure mercury lamp (400 W) in benzene for 20 h (Scheme 1). After evaporation of benzene and work-up, the residue was chromatographed to

give **7** and **8** quite readily (Table 1). As shown in Table 1, the results reveal that the reactions are clean and aldehydes or ketones were obtained quantitatively with the recovered naphthalene-1, 8-dithiole (**8**). Other five- and six-membered 1,3-dithia derivatives **9** and **10** were unreactive under the present photolysis conditions. Although 1-naphthylthioacetal monooxide (**11**) underwent photodecomposition under identical conditions, the products obtained were a mixture of intractable compounds which were unable to separate by a simple procedure. Therefore, the mechanism for the decomposition of **4** and **5** can be explained by the initial photo-excitation of the sulfoxides to form the reactive intermediates **6** probably by the intramolecular migration of the sulfinyl oxygen atom to the 2-carbon atom *via* the S-S through space interaction. The photochemical oxygen migration of sulfoxides has been reported in the literature.¹²⁾ Finally, the intermediate sulfenic esters **6** should be converted to the corresponding aldehydes or ketones and naphthalene-1, 8-dithiole (**8**).





Table 1. Photolysis of Dithioacetal Monooxides **4** and Dithioacetal Monooxides **5** in Benzene^{a)}

	R	R'	Yield of 7 /(% ^{b)})	Yield of 8 /(% ^{b)})
4a	Ph	H	>99	>99
4b	CH ₃ (CH ₂) ₅	H	>99	>99
4c	PhCH=CH	H	>99 (97) ^{c)}	>99 (98) ^{c)}
4d		H	>99	>99
5a	Ph	D	>99	>99
5b	Ph	CH ₃	>99 (93) ^{c)}	>99 (100) ^{c)}
5c	Ph	CH ₃ CH ₂	>99 (95) ^{c)}	>99 (99) ^{c)}
5d	Ph		>99 (96) ^{c)}	>99 (98) ^{c)}
5e	Ph		>99 (98) ^{c)}	>99 (100) ^{c)}
5f	Ph	PhCH(OH)-	- (82) ^{c)}	- (91) ^{c)}
5g	Ph	PhCH(OCH ₃)-	>99 (97) ^{c)}	>99 (98) ^{c)}

a) 400 W high pressure Hg lamp, $\lambda > 300$ nm, Substrates (0.1 mmol), Benzene (5 ml). b) Yields were determined by gas chromatography and ¹H-NMR spectroscopy. c) Isolated yields.

In order to understand the mechanism for the oxygen migration ^{18}O labelled compound **4a**- ^{18}O (^{18}O content=97 %) was prepared and subjected to photolysis under irradiation with a high pressure Hg lamp. The contents of ^{18}O were determined with mass spectrometric measurement. The sulfinyl ^{18}O labelled in **4a** was found to migrate completely to the oxygen atom of benzaldehyde (^{18}O content=94 %) as shown in Scheme 2.

Furthermore, 2-deuterated 2-phenyl-naphtho[1,8-*de*]1,3-dithiin-1-oxide (**5a**) was also photolyzed under identical conditions as described above to result in the formation of benzaldehyde-1D ($\text{D}>99\%$).¹³⁾ Thus, this photolysis of the compounds **4a** and **5a** are a convenient process to prepare the ^{18}O and D labelled aldehydes.



Recently, Foote et al. and Clennan et al. have reported the photooxidation of 1,5-dithiacyclooctane with oxygen in the presence of a sensitizer, in which they have described that the transannular S-S interaction determines the product formation.¹⁴⁾ Furthermore photolysis of simple dithianes or dithiolanes in the presence of dyes as sensitizers has also been reported to give aldehydes or ketones in high yields.¹⁵⁾ The "environmentally benign" nature of the procedure has been stressed to be advantageous in synthetic organic chemistry.

The advantage of our present procedure is a promised method to prepare aldehydes and ketones in high yields without use of either unpleasant-smelling alkyl dithiols or toxic heavy metals or strong acids for deprotection. Furthermore, the starting material naphthalene-1,8-dithiol (**8**) can be recovered completely and recycled after the reactions.

References

- 1) E. Block, "Reactions of Organosulfur Compounds," ed by A. T. Blomquist and H. H. Wasserman, Academic Press, New York (1978), Vol. 37, pp. 56.
- 2) E. J. Corey and B. W. Erickson, *J. Org. Chem.*, **36**, 3553 (1971); D. Seebach and H. Neumann, *Chem. Ber.*, **107**, 847 (1974); R. Ballini and M. Petrini, *Synthesis*, **1990**, 336; T. Taguchi, H. Okamura, and H. Takei, *Chem. Lett.*, **1975**, 853.
- 3) J. B. Chattopadhyaya and A. V. Rama Rao, *Tetrahedron Lett.*, **1973**, 3735; W. F. J. Huurdeman, H. Wynberg, and D. W. Emerson, *ibid.*, **1971**, 3449; M. Hojo and R. Masuda, *Synthesis*, **1976**, 678; K. Nishida, K. Yokota, D. Nakayama, T. Sumiya, M. Node, and K. Fuji, *Tetrahedron Lett.*, **34**, 345 (1993).
- 4) M. Fetizon and M. Jurion, *J. Chem. Soc., Chem. Commun.*, **1972**, 382; H. L. Wang-Chang, *Tetrahedron Lett.*, **1972**, 1989; T. Oishi, H. Takechi, K. Kamemoto, and Y. Ban, *ibid.*, **1972**, 1085; I. Stahl, M. Hetschko, and J. Gosselck, *ibid.*, **1971**, 4077; T. L. Ho and C. M. Wong, *Synthesis*, **1972**, 561.

- 5) T. T. Takahashi, C. Y. Nakamura, and J. Y. Satoh, *J. Chem. Soc., Chem. Commun.*, **1977**, 680; O. Hoshino, S. Sawai, and B. Umezawa, *Chem. Pharm. Bull.*, **27**, 538 (1979); M. Kamata, Y. Kato, and E. Hasegawa, *Tetrahedron Lett.*, **32**, 4349 (1991).
- 6) A. M. Martre, B. Mousset, R. B. Rhlid, and H. Veschambre, *Tetrahedron Lett.*, **31**, 2599 (1990); R. S. Glass, A. Petsom, and G. S. Wilson, *J. Org. Chem.*, **51**, 4337 (1986); J. Gourcy, P. Martigny, J. Simonet, and G. Jeminet, *Tetrahedron*, **37**, 1495 (1981); Q. N. Porter and J. H. P. Utley, *J. Chem. Soc., Chem. Commun.*, **1978**, 255; M. Planten and E. Steckhan, *Chem. Ber.*, **117**, 1679 (1984).
- 7) T. Fukuyama and T. Kishi, *J. Am. Chem. Soc.*, **98**, 6723 (1976); K. Ogura, M. Yamashita, M. Suzuki, S. Furukawa, and G. Tsuchihashi, *Bull. Chem. Soc. Jpn.*, **57**, 1637 (1984).
- 8) R. S. Glass, W. Andreuski, J. L. Broeker, H. Firouzabadi, L. K. Steffen, and G. S. Wilson, *J. Am. Chem. Soc.*, **111**, 4036 (1989); R. S. Glass, L. Adamowicz and J. L. Broeker, *ibid.*, **113**, 1065 (1991); H. Fujihara, J. -J. Chiu, and N. Furukawa, *Chem. Lett.*, **1990**, 2217; H. Fujihara, M. Yabe, J. -J. Chiu, and N. Furukawa, *Tetrahedron Lett.*, **32**, 4345 (1991).
- 9) H. Fujihara, R. Saito, M. Yabe, and N. Furukawa, *Chem. Lett.*, **1992**, 1437; N. Furukawa, H. Shima, and T. Kimura, *J. Chem. Soc., Chem. Commun.*, **1993**, 1762.
- 10) H. Fujihara, M. Yabe, and N. Furukawa, *J. Org. Chem.*, **58**, 5291 (1993).
- 11) The compounds **2**, **3**, **4**, and **5** were prepared and their structures were determined by ^1H -, ^{13}C -NMR, and mass spectrometries. They gave satisfied elemental analysis.
- 12) I. W. J. Still, "The chemistry of sulphones and sulfoxides," ed by S. Patai, Z. Rappoport, and C. J. M. Stirling, John Wiley & Sons, Chichester (1988), p. 873; D. C. Dittmer and G. C. Kuhlman, *J. Org. Chem.*, **35**, 3676 (1970); A. G. Schultz and R. H. Schlessinger, *J. Chem. Soc., Chem. Commun.*, **1970**, 1294; A. G. Schultz and R. H. Schlessinger, *Tetrahedron Lett.*, **1973**, 3605; K. Kobayashi and K. Mutai, *ibid.*, **22**, 5201 (1981); I. W. J. Still, P. C. Arora, M. S. Chauhan, M. H. Kwan, and M. T. Thomas, *Can. J. Chem.*, **54**, 455 (1976).
- 13) D. Seebach, B. W. Erickson, and G. Singh, *J. Org. Chem.*, **31**, 4303 (1969); F. Mutterer and J. P. Fluery, *ibid.*, **39**, 640 (1974).
- 14) C. Sheu, C. S. Foote, and C. -L. Gu, *J. Am. Chem. Soc.*, **114**, 3015 (1992); E. L. Clennan, D. -X. Wang, K. Yang, D. J. Hodgson, and A. R. Oki, *ibid.*, **114**, 3021 (1992).
- 15) D. L. Illman, *Chemical and Engineering News*, **1993**, Sept. 6, 26.

(Received February 28, 1994)