

Synthesis, Structure and Properties of [7]Heterohelices Possessing Phenolic Hydroxy Functions

Kazuhiko Tanaka,* Takuji Kume, Tadashi Takimoto, Yoshinori Kitahara, Hitomi Suzuki, Hideji Osuga, and Yasushi Kawai[†]
 Department of Chemistry, Graduate School of Science, Kyoto University, Kyoto 606-01
[†]Institute for Chemical Research, Kyoto University, Uji, Kyoto 611


(Received February 19, 1997; CL-970120)

Photocyclization of 2,7-bis[3-(*t*-butyldimethylsilyloxy)-phenylvinylene]benzo[1,2-b:4,3-b']dithiophene affords three regioisomers of the bis(silyloxy)[7]thiaheterohelices together with the unexpected 4-phenyl-substituted [7]thiaheterohelicene. Cleavage of the silyl ethers gave the helical quinone, which includes ethanol as guest molecules to form a 1:1 inclusion complex.


Atropisomeric 1,1'-bi-2-naphthol has been used in a wide range of asymmetric synthesis¹ as well as in optical resolution,^{2,3} where the phenol functions are essential for high stereo-differentiating reactions. In contrast, the asymmetric synthesis based on helicity is relatively unexplored,⁴⁻⁸ probably due to the difficulty in the preparation of optically active functionalized helical molecules with high stability. Among helices such as helicenes,⁹⁻¹¹ helicates^{12,13} and helixanes,¹⁴ helicene skeletons have a rigid framework and possess high optical stability. The synthesis and properties of functionalized carbohelicenes such as helical bis(quinones) have been extensively studied by Katz and co-workers, who utilize a Diels-Alder reaction for the construction of the quinones.¹⁵ However, the chemistry of helical phenol derivatives has not been fully investigated. We report here the photocyclization of 2,7-bis[3-(*t*-butyldimethylsilyloxy)phenylvinylene]benzo[1,2-b:4,3-b']dithiophene (**3**), and the structure and properties of the resulting [7]thiaheterohelices containing phenolic hydroxy functions.

The olefin **3** was obtained in 80% yield from 2,7-diformylbenzodithiophene (**1**) and phosphonium salt (**2**) derived from 3-hydroxybenzaldehyde (Scheme 1). Irradiation of **3** with a high-pressure mercury lamp under Katz's conditions¹⁶ gave a mixture of three possible regioisomers, the *para,para*-coupled product (**4a**), the *ortho,para*-coupled product (**5a**) and the *ortho,ortho*-coupled product (**6a**) in 63% yield with a ratio of 1:3:2. The structures of **4a**, **5a** and **6a** were determined by ¹H-NMR and MS,¹⁷ and in the case of **6a** by X-ray crystallographic analysis (Figure 1). The stereoview of **6a** clearly indicates that the two *t*-butyldimethylsilyl groups located on opposite sides of the helicene framework are directed *anti* in order to minimize the steric interaction and the whole molecule conforms to the idealized C₂ symmetry. We were surprised to isolate a minor, additional product (3% yield) in this photocyclization reaction because the crystal structure analysis shows that the compound **7a**¹⁷ possesses both helical and biphenyl frameworks (Figure 2). The dihedral angle between the terminal benzene units of the helicene part is 15.9°, and the phenyl ring of the biphenyl part is tilted relative to the terminal benzene by 49.7°, which is very close to that of 2,2'-biphenylcarboxamide (49.6°).¹⁸

Treatment of **4a** with (*n*-Bu)₄NF at 0 °C in THF gave 3,14-(dihydroxy)dinaphtho[1,2-d:1',2'-d']benzo[1,2-b:4,3-b']dithiophene (**4b**) as a yellow solid in 90% yield. In contrast, the reaction of **6a** with (*n*-Bu)₄NF produced a dark-red solution. Stirring the solution overnight under aerobic conditions afforded a black



Scheme 1. Reagents and conditions: i, *t*-BuOK, THF; ii, hv, I₂, benzene.



Scheme 2. Reagent and conditions: i, (*n*-Bu)₄NF, THF.




Figure 1. Stereoview of 6a.




Figure 2. Stereoview of 7a.




Figure 3. Molecular structure of [8][EtOH].

solid after chromatography on silica gel. X-ray analysis indicates that the molecule possesses a helical quinone structure **8** and contains both ethanol and a disordered guest in its cavity, although it was not possible to determine unambiguously the identity of the second guest¹⁷ (Figure 3). The oxygen atom of the guest molecule is hydrogen-bonded to the hydroxy group of one enantiomer of **8** with $O\cdots H = 1.86 \text{ \AA}$ and $O\cdots H-O = 170^\circ$, while the hydrogen atom interacts with the $C=O$ group of its antipode through a hydrogen bond with $H\cdots O = 1.86 \text{ \AA}$ and $O-H\cdots O = 164^\circ$. The dihedral angle between the quinone plane and the phenol plane of **8** is 26.2° . This is the first example of isolation of the supramolecular helical inclusion complex where the guest molecules are embedded between (*M*)- and (*P*)-quinones.

We gratefully acknowledge Grants-in-Aid from the Ministry of Education, Science, Sports and Culture (Nos. 07558089 and 08236221) and JSPS Fellowships for Young Scientists (No. 3089).

References and Notes

- For a recent review, see: R. Noyori, *Tetrahedron*, **50**, 4259 (1994) and references cited therein.
- K. Mori and F. Toda, *Chem. Lett.*, **1988**, 1997.
- K. Tanaka, A. Moriyama, and F. Toda, *J. Chem. Soc., Perkin Trans. 1*, **1996**, 603.
- T. Mizutani, S. Yagi, A. Honmaru, and H. Ogoshi, *J. Am. Chem. Soc.*, **118**, 5318 (1996).
- G. S. Hanan, J.-M. Lehn, N. Kyritsakas, and J. Fischer, *J. Chem. Soc., Chem. Commun.*, **1995**, 765.
- K. Ohkubo, T. Hamada, and H. Ishida, *J. Chem. Soc., Chem. Commun.*, **1993**, 1423.
- J. Jurczak, P. Lipkowski, T. Stankiewicz, and Z. Urbanczyk-Lipkowska, *Supramol. Chem.*, **6**, 87 (1995).
- F. Toda, K. Tanaka, Z. Stein, and I. Goldberg, *Acta Cryst.*, **B51**, 856 (1995).
- R. H. Martin, *Angew. Chem., Int. Ed. Engl.*, **13**, 646 (1974).
- H. Wynberg, *Acc. Chem. Res.*, **4**, 65 (1971).
- W. H. Laarhoven and J. C. Prinsen, *Top. Curr. Chem.*, **125**, 63 (1984).
- W. Zarges, J. Hall, and J.-M. Lehn, *Helv. Chim. Acta*, **74**, 1843 (1991).
- G. Bernardinelli, C. Piguet, and A. F. Williams, *Angew. Chem., Int. Ed. Engl.*, **31**, 1622 (1992).
- D. Gange, P. Magnus, L. Bass, E. V. Arnold, and J. Clardy, *J. Am. Chem. Soc.*, **102**, 2134 (1980).
- C. Nuckolls, T. J. Katz, and L. Castellanos, *J. Am. Chem. Soc.*, **118**, 3767 (1996).
- L. Liu, B. Yang, T. J. Katz, and M. K. Poindexter, *J. Org. Chem.*, **56**, 3769 (1991).
- Satisfactory analyses have been obtained for the new compounds **4a**, **5a,6a**, **4b**, **7a**, and **8**.
- F. Toda, A. Kai, Y. Tagami, and T. C. W. Mak, *Chem. Lett.*, **1987**, 1393.