

Photoinduced Electron Transfer Reaction of 3,3-Dimethyl-3*H*-pyrazoles: The Formation of Solvent Adducts through Cyclopropene Derivatives

Nobukazu Miyagawa,[†] Takashi Karatsu, and Akihide Kitamura*

Department of Materials Science, Faculty of Engineering, Chiba University, 1-33, Yayoi-cho, Inage-ku, Chiba 263

[†]Graduate School of Science and Technology, Chiba University, 1-33, Yayoi-cho, Inage-ku, Chiba 263

(Received June 19, 1997; CL-970472)

3,3-Dimethyl-3*H*-pyrazoles were photolyzed, and in the presence of 2,4,6-triphenylpyrylium tetrafluoroborate (TPP⁺) as a sensitizer in acetonitrile; in addition to cyclopropenes, 2*H*-pyrroles were obtained as solvent adducts to 1,3-radical cation intermediate.

Diradicals and radical-cations, which are generated by photolysis¹ and the photoinduced electron transfer (PET) reaction² of cyclic azoalkanes, give cyclization products and/or rearrangement products. On the other hand, 3*H*-pyrazoles, which have an unsaturated carbon-carbon double bond with an azo chromophore in its skeleton, have been as a vinyl carbene precursor and the mechanism of carbene generation has been studied.³ In the present paper, we report the photochemistry of 3,3-dimethyl-3*H*-pyrazoles **1** in the presence of 2,4,6-triphenylpyrylium tetrafluoroborate (TPP⁺) as an electron accepting sensitizer, and the comparison of the two radical-cations (**6** and **9**) generated in the reaction.

3*H*-pyrazoles **1a-d** were prepared by the cycloaddition of 2-diazopropane to the corresponding alkynes.^{2g} The structure of **1a** was identified by an X-ray crystallographic analysis.⁴

In order to examine the interaction between singlet excited sensitizers and **1**, fluorescence quenching experiments were performed. Fluorescences of TPP⁺ and 9,10-dicyanoanthracene (DCA) were efficiently quenched by **1a** with the quenching rate constants (k_q) of 1.1×10^{10} and 1.3×10^9 l mol⁻¹ s⁻¹, respectively. For other pyrazoles, the k_q s were close to the diffusion-controlled limit. The differences in free energy (ΔG) for single electron transfer (SET) are exergonic or slightly endergonic (ΔG is calculated as -11.8 and 1.15 kcal mol⁻¹ for **1a** and TPP or DCA, respectively) as calculated from the oxidation potentials of the pyrazoles (for example E_{ox} of **1a** is 2.02 V vs. SCE) using the Rehm-Weller equation.⁵ These results suggest that these reactions proceed via an electron transfer.

3*H*-pyrazoles **1** in the presence of TPP⁺ were irradiated in acetonitrile using a 405 nm light under deaerated conditions. The pyrazoles **1** lost nitrogen to give the corresponding cyclopropenes **2**, which were the same products obtained in the direct photolysis of **1**, and the product **3** having acetonitrile unit. The product distributions are summarized in Table 1. In the photo-reaction of **1** in the presence of DCA, only **2** was obtained. The electron transfer reaction of cyclopropene derivatives sensitized by DCA is known to cause their dimerization,^{6,7} but the corresponding dimers were not obtained in each reactions.

The structure of **3** was determined by ¹H NMR and its NOE measurement.⁸ The ¹H NMR spectrum was simple⁹ and the NOE signal was only observed between the aromatic proton and the methyl proton originated from the acetonitrile. **3** is not the adduct of the nitrogen-eliminated intermediate from **1** to

Table 1. Conversion yields of **2** and **3** for photolysis of **1** with TPP⁺ ^a

Pyrazole	Decomposition (%)	Yield (%) ^b	
		2	3
1a	10	85	trace
	51	35	15
1b	9	49	28
1c	30	20	22
1d ^c	25	82	trace ^d

^a3*H*-pyrazole **1** (5-8 mM) with TPP⁺ (ca. 1.5 mM) in acetonitrile was irradiated using a 405 nm light (1 kW Hg lamp) under deaerated conditions. ^bConversion yields based on the decomposition obtained by GC analysis. ^cThe concentration of **1d** was 1 mM. When the concentration was 6 mM, only precipitation was obtained. ^dThese yields were estimated by GC and GC-MS analysis. See ref. 8.





Figure 1. Decomposition of 3*H*-pyrazole **1a** and formation of products. 5.26 mM of **1a** with TPP⁺ (1.4 mM) in MeCN was irradiated using 405 nm light (1 kW Hg lamp) under degassed conditions. **1a** (---▲---); **2a** (—●—); **3a** (---■---).

acetonitrile. It is impossible to explain the formation of **3** with the simple PET mechanism, because the structure of the solvent adducts was not **8** which forms from nitrogen eliminated intermediate of **6** (Scheme 1). We investigated the time correlation of product yields. Figure 1 shows the decrease of **1a**, and the yields of **2a** and the solvent adduct **3a**. The product yield of **2a** plateaus after an hour of irradiation, and the yield of **3a** increased with an induction time. The irradiation of isolated **2a** in the presence of TPP⁺ under the same conditions showed the rapid decomposition of **2a** and the production of **3a** at 30% conversion yield. Furthermore, the TPP⁺ fluorescence was quenched by cyclopropene **2a** with k_q value of 1.7×10^{10} , and the calculated ΔG between **2a** and TPP⁺ is -11.8 kcal mol⁻¹ (E_{ox} =2.02 V for **2a**). From these results, it is apparent that the formation of **3a** from the TPP⁺ sensitized decomposition of **1a**



Scheme 1. Reaction mechanism for photolysis of 3*H*-pyrazole **1** in the presence of TPP^+ as a sensitizer.

proceeds via **2a**. Moreover, the structures of **3** support this mechanism. The addition of acetonitrile to radical cation intermediates is rarely reported,^{2b,f,10} and none reported the addition of acetonitrile to cyclopropenes to form 2*H*-pyrrole derivatives.

In the reactions in which the yields of **2** are low, their mass balances are also low (Table 1). This tendency can be explained by the following reasons. Some of **2** were thermally unstable and polymerized,^{3a,11} thus the yields of **3** from PET reactions of **2** were low. In fact, precipitates regarded as a polymer were observed in some reactions.¹¹

We postulate that this reaction mechanism using TPP^+ is as follows (Scheme 1). SET occurs between the 3*H*-pyrazole **1** and an excited sensitizer to form the initial radical cation $1^{\bullet+}$ and the neutral radical of the sensitizer ($\text{TPP}^{\bullet}$). $1^{\bullet+}$ immediately loses a nitrogen to form the radical cation **6**. At this step, C-N bond cleavage between the olefin and azo group is unfavorable since conjugation between olefin and azo group is interrupted. After elimination of nitrogen, cation site must be at σ -orbital, however, MOPAC calculation shows that the stable configuration of the cation radical is **6** (π -type cation), therefore electron in **5** might be redistributed to more stable configuration **6** quickly. The back electron transfer (BET) from the $\text{TPP}^{\bullet}$ to **6** and cyclization (CYC) or CYC and BET produce cyclopropene **2**. If **2** is unstable, its polymerization could occur. Here, if **2** can undergo PET to an excited sensitizer, the radical cation $2^{\bullet+}$ is formed. The C-C bond of $2^{\bullet+}$ is then cleaved to form **9**, and acetonitrile, which is a weak nucleophile, adds to **9**, and resulting in the formation of 2*H*-Pyrrole **3**. The reason why only **9** gives **3** but **6** does not give **8** might be explained as follows. Acetonitrile adds to the cation radical **6** to give **7** or **9** to give **10**. However, spin density of R_2 substituted carbon of **7** must be low by a substitution of electron-withdrawing ester group and it prevents further cyclization (CYC) to **8**, thus **7** might return to **6**. To examine the possibility of generation of **6** by SET of **2** in the presence of TPP^+ , a trapping experiment by alcohol was performed, but it was unsuccessful at this moment.

In the case of DCA, the acetonitrile adduct was not formed because BET from $\text{DCA}^{\bullet-}$ to **6** or **9** can occur more efficiently than that from $\text{TPP}^{\bullet}$.¹²

This work was supported by a Grant-in-Aid for Scientific Research (C) (2) 08640671 from the Ministry of Education, Science and Culture of Japan and Futaba Denshi Foundation.

References and Notes

- a) P. S. Engel, *Chem. Rev.*, **80**, 99 (1980). b) P. S. Engel and L. Shen, *Can. J. Chem.*, **52**, 4040 (1974).
- a) T. Karatsu, H. Hotta, and A. Kitamura, *J. Chem. Soc., Chem. Commun.*, **1991**, 1451. b) T. A. Zona and J. L. Goodman, *J. Am. Chem. Soc.*, **115**, 4925 (1993). c) W. Adam and J. Sendelbach, *J. Org. Chem.*, **58**, 5310 and 5316 (1993). d) W. Adam, C. Sahin, J. Sendelbach, H. Walter, G.-F. Chen, and F. Williams, *J. Am. Chem. Soc.*, **116**, 2576 (1994). e) H. Ikeda, T. Minegishi, and T. Miyashi, *J. Chem. Soc., Chem. Commun.*, **1994**, 297. f) T. A. Zona and J. L. Goodman, *J. Am. Chem. Soc.*, **117**, 5879 (1995). g) T. Karatsu, H. Itoh, T. Kikunaga, Y. Ebashi, H. Hotta, and A. Kitamura, *J. Org. Chem.*, **60**, 8270 (1995). h) H. Ikeda, T. Minegishi, Y. Takahashi, and T. Miyashi, *Tetrahedron Lett.*, **39**, 4377 (1996).
- For example; a) G. L. Closs, W. A. Boll, H. Heyn, and V. Dev, *J. Am. Chem. Soc.*, **90**, 173 (1968). b) L. Schrader, *Chem. Ber.*, **104**, 941 (1971). c) M. F.-Neumann and J.-J. Lohmann, *Angew. Chem. Int. Ed. Engl.*, **16**, 323 (1977). d) H. Heydt and M. Regitz, *J. Chem. Res., (S)* **1978**, 326. e) C. L. Leach, Jr., and J. W. Wilson, *J. Org. Chem.*, **43**, 4880 (1978). f) M. F.-Neumann and M. Miesch, *Tetrahedron Lett.*, **25**, 2909 (1984).
- Colorless crystals of **1a** are prismatic, the space group is $P2_1/a$ (#14) with $a = 8.523(1) \text{ \AA}$, $b = 18.485(23) \text{ \AA}$, $c = 8.544(1) \text{ \AA}$, $V = 1262.6(3) \text{ \AA}^3$, $Z = 4$, and $D_c = 1.211 \text{ g}\cdot\text{cm}^{-3}$. The final R value was 0.078 for 1371 reflections with $I > 3.5\sigma(I)$.
- a) D. Rehm and A. Weller, *Ber. Bunsenges. Phys. Chem.*, **73**, 834 (1969). b) D. Rehm and A. Weller, *Isr. J. Chem.*, **8**, 259 (1970).
- A. Padwa, S. I. Goldstein, and R. J. Rosenthal, *J. Org. Chem.*, **52**, 3278 (1987).
- a) C. D. DeBoer, D. H. Wadsworth, and W. C. Perkins, *J. Am. Chem. Soc.*, **95**, 861 (1973). b) K. A. Brown-Wensley, S. L. Mattes, and S. Farid, *J. Am. Chem. Soc.*, **100**, 4162 (1978).
- 3d** was only detected on GC and GC-MS, because of its low yield.
- For example, **1a**: 270 MHz, CDCl_3 ; δ 7.24-7.38 (m, 2H, Ar), 7.25-7.19 (m, 3H, Ar), 3.61 (s, 3H, CO_2Me), 2.12 (s, 3H, $\text{MeC}=\text{N}$), 1.45 (s, 6H, Me).
- a) T. A. Zona and J. L. Goodman, *Tetrahedron Lett.*, **33**, 6093 (1992). b) K. Mizuno, K. Nire, H. Sugita, and Y. Otsuji, *Tetrahedron Lett.*, **34**, 6563 (1993).
- 2a-d** are not stable. Especially **2d** was polymerized in a few hours at rt; cf. ref 3(a).
- The $2^{\bullet+}/\text{TPP}^{\bullet}$ or $10/\text{TPP}^{\bullet}$ pair may make solvent separated pairs more efficiently than the $2^{\bullet+}/\text{DCA}^{\bullet-}$ or $10/\text{DCA}^{\bullet-}$ pair. cf. T. Karatsu, H. Kobayashi, E. Shinkai, and A. Kitamura, *Chem. Lett.*, 2131 (1992).