


Dicyanodiaryl-*p*-quinodimethanes : An Efficient Synthesis Using a New Dilithium Reagent and Their Solvent-Dependent Properties

Takeshi Kawase, Masaki Wakabayashi, Chizuko Takahashi, and Masaji Oda*
Department of Chemistry, Graduate School of Science, Osaka University, Toyonaka, Osaka 560

(Received July 4, 1997; CL-970520)

A series of 7,7-dicyano-8,8-diaryl-*p*-quinodimethane are readily prepared by the reaction of 4-lithiophenyldicyanomethyl-lithium with diaryl ketones; the 8-(4-dimethylaminophenyl) compounds thus obtained show solvent-dependent absorption spectra and bond rotation of the exocyclic double bonds and form a head-to-tail dimeric structure in the crystal.

Highly dipolar π -electron systems have attracted considerable attention in relation to development of new materials.¹ Dicyanoquinodimethanes **1** having electron-donor(s) on the other side of methylene carbon are a class of molecules with highly dipolar property and a number of such molecules have been reported.²⁻⁴ Gompper and co-workers reported the synthesis of a number of dicyanoquinodimethanes including 7,7-dicyano-8,8-diphenyl-*p*-quinodimethane (**2a**).² Introduction of substituent(s) in the phenyl group(s) of **2a** would allow tuning of polarity and provide valuable information for design of novel quinodimethanes of physicochemical interest. Gompper's synthetic method for **2a** seems, however, not effectively applicable to the synthesis of its derivatives. Here we report a new and efficient synthesis of dicyanodiarylquinodimethanes **2a-2e**, X-ray structure of **2d**, and their properties involving substituent effects.



It was found that 4-lithiophenyldicyanomethyl-lithium **5** is a new versatile synthon for dicyanoquinodimethanes.

When (4-bromophenyl)malononitrile (**3**)⁵ was treated with 2.2 equiv. of *n*-BuLi in THF at -78 °C, the bromine-lithium exchange forming dilithio compound **5** was considerably slow probably due to electronic effect of the negative charge in monoanion **4** formed initially; however, 6 hours' reaction (1.0 mmol scale; 15 ml of THF) was found enough for the dilithiation as judged by dimethylation to **6** (85% yield; MeI quenching). Shorter time or higher temperature for the lithiation resulted in poorer yield of **6**. The cyano groups remain intact under the lithiation condition, although intramolecular electronic repulsion should increase nucleophilic reactivity of the aryllithium part of **5**.

Reactions of **5** with benzophenone and its derivatives followed by chromatography on silica gel, where dehydration mostly takes place, afforded **2a-e** in good yields (Scheme 1).^{6,7} Adamantanone, an unenolizable dialkyl ketone, also yielded dicyanoquinodimethane **7**.⁷ However, enolizable ketones such as

cyclohexanone and camphor were not successful enough, because their reactions produced mixtures mainly owing to aromatizing tendency of the corresponding quinodimethanes.





Table 1 lists the selected physical data of **2a-2e** and **7**. ¹³C NMR spectra clearly show the large dipolar property of **2a-2e** as well as the strong substituent effects: the chemical shift difference between C7 and C8 [$\Delta\delta$ (C8-C7)] increases sharply from 64.8 ppm of **2a** to 115.4 ppm of **2e**. Charge separation of **7** ($\Delta\delta$ = 61.0 ppm) is smaller than **2a**. It has been reported that there is a linear correlation between the nitrile stretching frequency and degree of charge transfer in TCNQ CT-complexes.⁸ As far as **2a-2e** are concerned, there seems also present a linear correlation between the nitrile stretching frequencies and the C7 chemical shifts of **2a-2e** (Figure 1).⁹ This correlation indicates clear and strong substituent effects on the resonance contribution of the dipolar structures such as **2d-B**.¹⁰

Quinodimethanes **2a-2e**, in particular **2d** and **2e**, show large solvent effects on absorption spectra; for example, **2d** exhibits a visible absorption at 611 nm in benzene, whereas much stronger absorptions at 696 nm in methanol. The solvent effects, which are attributable mainly to twisted intramolecular charge transfer (TICT¹¹), are more prominent than those observed for the corresponding *p*-quinone methides.¹²




Temperature-dependent ¹H-NMR spectra of unsymmetrical **2d** agree with the solvent-dependent TICT. In bromobenzene-*d*₅, **2d** shows broadening of the four non-equivalent protons in the quinodimethane moiety (Ha-Hd, see **2d-A**) only above 110 °C ($\Delta G^\ddagger > 18$ kcal·mol⁻¹ for the rotation of C4-C8 bond). However, in dimethylformamide (DMF)-*d*₇, Hc and Hd are observed to be

Table 1. Selected spectral and electrochemical data of **2a-e**

Compound	CN stretching ^a (cm ⁻¹)	δ C7 ^b (ppm)	δ C8 ^b (ppm)	$\Delta\delta$ (C8-C7) (ppm)	Longest absorption maxima $\lambda_{\max}$ nm(log ϵ)/CH ₂ Cl ₂	Redox potentials (V) ^c		
						E ¹ _{ox} ^d	E ¹ _{red}	E ² _{red} ^d
2a	2209	80.22	145.03	64.8	495 (4.59)	1.38	-0.44	-1.46
2b	2204	78.18	152.62	74.4	514 (4.56)	1.29	-0.47	-1.46
2c	2199	66.09	155.61	89.5	558 (4.65)	1.15	-0.51	-1.47
2d	2196	63.00	166.65	103.7	648 (4.61), 690sh (4.55) 611 (4.51)/C ₆ H ₆ 696 (4.72)/MeOH	0.83	-0.56	-1.45
2e	2186	55.13	170.48	115.4	640sh (4.50), 698 (4.82) 646 (4.69), 690sh (4.60)/C ₆ H ₆	0.76	-0.72	-1.60
7	2211	96.14	157.14	43.6	436 (4.63), 457 (4.59)	0.33	-1.73 ^d	—

^a KBr disk. ^b In CDCl₃. ^c V vs Ag/AgCl, in 0.1 M nBu₄NClO₄/DMF, sweep rate 100 mVsec⁻¹ at 25 °C, Fc⁺/Fc = 0.52 V. ^d Peak potential.

**Figure 1.** Correlation of nitrile stretching frequencies and dicyanomethylene carbon chemical shifts of **2a-e**.

equivalent at 100 °C, appreciably broadened at 30 °C, almost disappearing at 10 °C, and two doublets (δ 7.23 and 7.61) below -50 °C [$\Delta G^\ddagger = 11.5 \pm 0.5$ kcal·mol⁻¹ ($T_c = -10 \pm 5$ °C)]. The energy barrier for the rotation is intermediate in less polar acetone-*d*₆ ($\Delta G^\ddagger = 13.0 \pm 0.5$ kcal·mol⁻¹).¹³

The cyclic voltammograms of **2a-2e** are composed of one irreversible oxidation wave and the first reversible and second irreversible reduction waves. Adamantylidene compound **7** behaves differently showing only one irreversible reduction wave at higher potential to suggest instability of its anion radical. Although much weaker than TCNQ ($E^1_{\text{red}} = 0.06$ V, $E^2_{\text{red}} = -0.33$ V), the electron affinity of **2a** is comparable to that of *p*-benzoquinone ($E^1_{\text{red}} = -0.45$ V, $E^2_{\text{red}} = -1.33$ V).

X-ray structure analysis of **2d**¹⁴ (Figure 2) revealed two interesting features, although the accuracy is lowered due to high


disorder of the included solvent molecules. First, while diamino-phenyl group is twisted by only 16° from the plane of quinodimethane, the phenyl group largely by 50° to indicate effective conjugation between the quinodimethane and diamino-phenyl groups. Second, the molecules are faced each other in a head-to-tail manner in crystal probably owing to large intermolecular dipole interaction.

Further studies on dicyanoquinodimethanes and synthetic application of **5** are in progress.

This work was supported by Grant-in-Aid for Scientific Research (No. 08454201 and 08640683) from Ministry of Education, Science and Culture (Japan).

References and notes

- R. Gompper and H.-U. Wagner, *Angew. Chem. Int. Ed. Engl.*, **27**, 1437 (1988); D. J. Williams, *Angew. Chem. Int. Ed. Engl.*, **23**, 690 (1984), and references cited therein.
- R. Gompper, H.-U. Wagner, and E. Kutter, *Chem. Ber.*, **101**, 4123 and 4144 (1968).
- K. Takahashi, N. Hirata, and K. Takase, *Tetrahedron Lett.*, **1970**, 1285.
- W. R. Hertler, H. D. Hartrizer, D. S. Acker, and R. E. Benson, *J. Am. Chem. Soc.*, **84**, 3387 (1962); O. A. El Seoud, F. P. Ribeiro, A. Martins, and P. P. Brotero, *J. Org. Chem.*, **50**, 5099 (1985) and references cited therein.
- M. Uno, K. Seto, M. Masuda, W. Ueda, and S. Takahashi, *Tetrahedron Lett.*, **26**, 1553 (1985).
- Treatment of the intermediate crude alcohols with phosphoryl chloride or methanesulfonyl chloride in pyridine also effect the dehydration.
- All new products gave satisfactory analytical and spectral data other than those given in Table 1. **2a**: dark red needles; mp 162.5-163.5 °C (lit. 2: 162 °C). **2b**: dark red needles; mp 218-219 °C. **2c**: dark purple needles; mp >300 °C. **2d**: metallic green needles; mp 106-107 °C. **2e**: metallic green needles; mp >290 °C dec. **6**: colorless oil; IR (oil) $\nu = 2252$ cm⁻¹ (CN); ¹H-NMR (270 MHz, CDCl₃) $\delta = 2.09$ (s, 3H), 2.39 (s, 3H), 7.29 (2H, AA'BB', $J = 1.6, 8.6$ Hz), 7.45 (2H, AA'BB', $J = 1.6, 8.6$ Hz). **7**: orange crystals; mp 283 °C dec.
- J. S. Chappell, A. N. Bloch, W. A. Bryden, M. Maxfield, T. O. Poehler, and D. O. Cowan, *J. Am. Chem. Soc.*, **103**, 2442 (1981).
- Simple extrapolation of the correlation to the nitrile frequency of TCNQ (2227 cm⁻¹) gives an estimated value of ca. δ 100 for the yet unknown chemical shift of C7(8) of TCNQ.
- The nitrile stretching frequency of **2e** (2186 cm⁻¹) corresponds to that of TCNQ complexes with about 0.9 unit charge transfer.⁹
- W. Schlenk, *Justus Liebig's Ann. Chem.*, **368**, 294 (1909); F. C. Adam and W. T. Simpson, *J. Mol. Spectroscopy*, **3**, 363 (1959).
- S. Hünig, H. Schwegberg, and H. Schwarz, *Justus Liebig's Ann. Chem.*, **587**, 132 (1954); S. Hünig and H. Schwarz, *ibid.*, **599**, 131 (1957).
- Similar solvent dependent easy C=C bond rotation was observed for 6-aminofulvene derivatives; J. H. Crabtree and D. J. Bertelli, *J. Am. Chem. Soc.*, **89**, 5384 (1967).
- Crystal data for **2d**·C₆H₆·0.12(CH₂Cl₂) (the composition was determined by elemental analysis): crystallized from benzene-dichloromethane; monoclinic, space group *C2/c*(#15), $a = 9.434(4)$, $b = 24.609(8)$, $c = 20.880(5)$ Å, $\beta = 90.09(2)^\circ$, $V = 2153(1)$ Å³, $Z = 8$, $\rho_{\text{calc}} = 1.215$ gcm⁻³, $m = 2.80$ cm⁻¹, 1713 independent observed reflections with $[F > 3\sigma(F)]$, $3^\circ \leq 2\theta \leq 54.9^\circ$, refined to $R(R_w) = 0.090(0.092)$.

**Figure 2.** Left: ORTEP drawing of **2d** (50% probability). Right: Head-to-tail arrangement of **2d** in the crystal.