

Copper-mediated Coupling Reaction of Kinetically Stabilized 1-Methyl-2-phosphaethenyllithiums Involving an Isomerization to a 3-Phospha-2-propenyllithium Derivative¹


Shigekazu Ito, Shigeo Kimura, and Masaaki Yoshifuji*

Department of Chemistry, Graduate School of Science, Tohoku University, Aoba, Sendai 980-8578

(Received April 11, 2002; CL-020307)

Copper-mediated coupling reaction of 1-methyl-2-(2,4,6-tri-*t*-butylphenyl)-2-phosphaethenyllithiums afforded 1,4-diphosphabuta-1,3-diene or 1,6-diphosphahex-1,5-diene derivatives, depending upon non-coordinated or coordinated to $W(CO)_5$, which facilitates an isomerization to a 3-phospha-2-propenyllithium.

We and others have reported various types of stable multiple bonds of phosphorus by utilizing the kinetic stabilization method.² Phosphaethenyllithiums **1** bearing the 2,4,6-tri-*t*-butylphenyl (hereafter abbreviated to Mes^{*}) group are suitable reagents for synthesis of some low-coordinated phosphorus compounds, and we have studied copper-mediated coupling reactions of **1**, where X=Cl, Br, SPh, and STol, i. e. lithium phosphanylidene carbenoids, affording 1,4-diphosphabuta-1,3-diene derivatives **2** as a conjugated system involving the P=C moieties.^{2,3} Recently, a considerable attention has been paid to coordination chemistry of multiple-bonded phosphorus compounds to develop a novel class of synthetic catalyst,⁴ and we expected to utilize **2** for a ligand of metal complexes as an analog of 3,4-diphosphanylidene cyclobutene derivatives **3**.^{4c-f,5} Although **2** displayed quite low reactivity to the transition metal such as tungsten or chromium, or undesirable reactivity to some palladium or platinum reagents, an alternative method was employed to synthesize phosphaethenyllithiums in a transition-metal complex form exemplified such as **4**.⁶ In this paper we report copper-mediated coupling reaction of a bulky 1-methyl-2-phosphaethenyllithium in a free form (**5**) or a coordinated form on the tungsten metal (**5w**).



(*Z*)-2-Bromo-1-(2,4,6-tri-*t*-butylphenyl)-1-phosphapropene (**6**) was allowed to react with butyllithium to afford 2-lithio-1-phosphapropene **5**,⁷ which was treated with copper(II) chloride and oxygen at $-78^\circ C$ to give 2,3-dimethyl-1,4-diphosphabuta-1,3-diene **7** as a mixture of three isomers (*EE*-, *ZZ*-, and *EZ*-**7**) almost quantitatively (Scheme 1). The ratio of the isomers was *EE* : *ZZ* : *EZ* = 1 : 1 : 3, which was determined by ^{31}P NMR spectroscopy.⁸ The mixture **7** was irradiated by a medium-pressure mercury lamp through a Pyrex filter ($\lambda > 300$ nm) or heated in refluxing toluene to afford the single isomer (*EE*-**7**), whereas (*EZ*)-1,4-bis(2,4,6-tri-*t*-butylphenyl)-1,4-diphosphabuta-1,3-diene was reported to afford the *ZZ*-isomer by irradiation or heating by Appel et al.⁹ The structure of *EE*-**7** was confirmed by X-ray crystallography revealing a planar *s*-trans conformation

with a torsion angle of $174.6(1)^\circ$ in the P=C-C=P system (Figure 1).⁸ The P=C bond lengths [1.703(2) and 1.699(2) Å] and the (P)C-C(P) bond length [1.465(3) Å] are slightly longer than the corresponding distances for **2** (X=Cl) [P=C 1.691(4) Å; C-C 1.445(7) Å],^{3a} probably due to an electron donating effect of the methyl groups.





Figure 1. Molecular structure of *EE*-7.

Coordination chemistry of **7** is of interest, because compound **3**, including the 1,4-diphosphabuta-1,3-diene skeleton, has been utilized as ligands for synthetic catalysts.⁴ But the coordinating property of **7** on transition metals turned out to be very poor by the fact that **7** did not coordinate on the transition-metals such as group 6 or group 10 metals. Thereupon we attempted to couple **5** on a transition metal. Compound **6** was allowed to react with pentacarbonyl(tetrahydrofuran)tungsten(0) to afford the corresponding complex **6w** in a moderate yield (60%).¹⁰ In an attempt to obtain complex **7w**, **6w** was allowed to react with butyllithium and copper(II) chloride, but the product was not **7w** but **8** in 30% yield as purple crystals,¹¹ indicating a [1,2]-hydrogen rearrangement to leading to the 1,6-diphosphahex-1,5-diene skeleton through an intermediate (**5'w**) (Scheme 1). Oxygen was not used in this coupling reaction to prevent decomplexation of **6w**. The structure of **8** was confirmed by X-ray crystallography as shown in Figure 2,¹¹ displaying a *ZZ* configuration. The P=C length



Scheme 1. Copper-mediated coupling reaction of 1-methyl-2-phosphaethenyllithiums. Reagents and conditions: a) *n*-BuLi, THF, $-78^\circ C$; b) $CuCl_2$, O_2 , $-78^\circ C$; c) Medium pressure Hg lamp (100 W), benzene, $0^\circ C$; d) toluene, reflux; e) $W(CO)_5$ (thf), rt; f) $CuCl_2$.





Figure 2. Molecular structure of 8.

[1.660(7) Å] is comparable to the corresponding value of **9** [1.635(4) Å],¹² and the P–W bond length [2.539(2) Å] is similar to that of the pentacarbonyltungsten(0) complex of 3,3-diphenyl-1-(2,4,6-tri-*t*-butylphenyl)-1-phosphaallene [2.531(2) Å].¹³ The C1–C2 and C2–C2* distances are 1.478(9) and 1.54(1) Å, respectively.

To verify the [1,2]-hydrogen shift of **5w**, we examined the reaction of **6w** as follows: **6w** was allowed to react with 1 eq of butyllithium and quenched with methanol-*d*₄ at -78 °C to give (Z)-3-deutero-1-phosphapropene complex **Z-10** in an 80%-D ratio, (E)-1-phosphapropene complex **E-11**, and 1-phosphaallene complex **12** in a 6 : 3 : 1 ratio almost quantitatively.¹⁰ Formation of **Z-10** indicates the [1,2]-rearrangement leading to a 3-phospha-2-propenyllithium intermediate (**5'w**). Compound **E-11** might have been generated by proton abstraction of **5w** from **6w**, indicating slow rate of the halogen-metal exchange to generate **5w**, and **12** might have been formed by an elimination of hydrogen bromide from **6w**. It indicates a facile [1,2]-hydrogen shift of **5w** proceeds to afford the 3-phospha-2-propenyllithium derivative (**Z-5'w**).



As for **5w**, coordination of the pentacarbonyltungsten(0) group might increase the acidity of the methyl group, which resembles in the property of (*η*⁶-arene)transition metal complexes.¹⁴ Moreover, we assume that steric congestion around the “solvated” lithium atom might facilitate the [1,2]-rearrangement. Although the detailed reaction mechanism for the facile [1,2]-rearrangement of **5w** has not been clarified, it is interesting to note that the reactivity of **5** changes upon coordination.

This work was supported in part by a Grant-in-Aid for Scientific Research (No. 13304049) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References and Notes

- This paper is dedicated to Prof. Dr. Dieter Seebach at Eidgenössische Technische Hochschule Zürich on the occasion of his 65th birthday.
- a) M. Yoshifuji, *J. Chem. Soc. Dalton Trans.*, **1999**, 3343. b) M. Yoshifuji, *J. Organomet. Chem.*, **611**, 210 (2000). c) M. Regitz and O. J. Scherer, “Multiple Bonds and Low Coordination in Phosphorus Chemistry,” Georg Thieme Verlag, Stuttgart (1990). d) K. B. Dillon, F. Mathey, and J. F. Nixon, “Phosphorus: The Carbon Copy,” Wiley, Chichester (1998).
- a) S. Ito, K. Toyota, and M. Yoshifuji, *Chem. Lett.*, **1995**, 747. b) S. Ito, K. Toyota, and M. Yoshifuji, *J. Organomet. Chem.*, **553**, 135 (1998). c) S. Ito and M. Yoshifuji, *Chem. Lett.*, **1998**, 651. d) S. Ito and M. Yoshifuji, *Chem. Lett.*, **2000**, 1390.
- a) B. Breit, *J. Mol. Catal. A: Chem.*, **143**, 143 (1999). b) R. Shintani, M. M.-C. Lo, and G. C. Fu, *Org. Lett.*, **2**, 3695 (2000). c) K. Toyota, K. Masaki, T. Abe, and M. Yoshifuji, *Chem. Lett.*, **1995**, 221. d) S. Ikeda, F. Ohhata, M. Miyoshi, R. Tanaka, T. Minami, F. Ozawa, and M. Yoshifuji, *Angew. Chem. Int. Ed.*, **39**, 4512 (2000). e) F. Ozawa, S. Yamamoto, S. Kawagishi, M. Hiraoka, S. Ikeda, T. Minami, S. Ito, and M. Yoshifuji, *Chem. Lett.*, **2001**, 972. f) T. Minami, H. Okamoto, S. Ikeda, R. Tanaka, F. Ozawa, and M. Yoshifuji, *Angew. Chem. Int. Ed.*, **40**, 4501 (2001). g) L. Weber, *Angew. Chem. Int. Ed.*, **41**, 563 (2002).
- N. Yamada, K. Abe, K. Toyota, and M. Yoshifuji, *Org. Lett.*, **4**, 569 (2002).
- a) P. Le Floch, D. Carmichael, and F. Mathey, *Organometallics*, **10**, 2432 (1991). b) H. Teunissen and F. Bickelhaupt, *Organometallics*, **15**, 794 (1996).
- a) S. Ito, K. Toyota, and M. Yoshifuji, *Chem. Commun.*, **1997**, 1637. b) G. Märkl and W. Bauer, *Angew. Chem. Int. Ed. Engl.*, **28**, 1695 (1989).
- a) To a solution of **6** (0.13 mmol) in THF (6 mL) was added butyllithium (0.15 mmol), and in 5 min copper(II) chloride (0.25 mmol) was added at -78 °C. The reaction mixture was stirred at -78 °C for 1 h, and oxygen was bubbled for 5 min (ca. 32 mmol). After treatment with aqueous sodium sulfite and warming up to room temperature, the mixture was treated with ammonia (10% NH₃ in saturated aqueous NH₄Cl) and extracted with ether. The organic layer was concentrated in vacuo, and the residue was purified by silica-gel column chromatography (hexane/toluene 5 : 1) to afford 37 mg of **7** (97% yield) as yellow solids. b) **EE-7**: Yellow crystals, mp 233–234 °C; ¹H NMR (200 MHz, CDCl₃) δ = 1.35 (18H, s, *p*-*t*Bu), 1.50 (36H, s, *o*-*t*Bu), 1.55 (m, 6H, Me), 7.41 (4H, brs, arom); ¹³C{¹H} NMR (50 MHz, CDCl₃) δ = 183.9 (dd, ¹J_{PC} = 13 Hz, ²J_{PC} = 3 Hz, P=C); ³¹P{¹H} NMR (81 MHz, CDCl₃) δ = 232; HRMS calcd for C₄₀H₆₄P₂, 606.4483; found, 606.4488. c) **ZZ-7**: δ_P = 237; **EZ-7**: δ_P = 240, 288 (AB_q, ³J_{PP} = 154 Hz). d) Crystal data for **EE-7**: C₄₀H₆₄P₂ *M* = 606.89, triclinic, *P*1 (no. 2), *a* = 18.524(3), *b* = 18.564(2), *c* = 6.237(6) Å, *α* = 98.23(2), *β* = 93.72(2), *γ* = 64.40(1)°, *V* = 1914(1) Å³, *Z* = 2, *ρ* = 1.053 g cm⁻³, *μ*(Mo Kα) = 0.138 mm⁻¹, *T* = 150 K, 5428 reflections measured (*2θ*_{max} = 50.0°), 5426 observed [*I* > 1.0σ(*I*)], *R*₁ = 0.047 [*I* > 2.0σ(*I*)]. *R*_w = 0.108 (all data). CCDC-181946.
- R. Appel, J. Hünerbein, and N. Siabalis, *Angew. Chem. Int. Ed. Engl.*, **26**, 779 (1987).
- ³¹P{¹H} NMR (162 MHz, CDCl₃) **6w**: δ = 210 (¹J_{PW} = 270 Hz); **Z-10**: δ = 208 (¹J_{PW} = 262 Hz); **E-11**: δ = 199 (¹J_{PW} = 259 Hz); **12**: δ = 42 (¹J_{PW} = 259 Hz).
- a) To a solution of **6w** (0.14 mmol) in THF (10 mL) at -78 °C was added butyllithium (0.15 mmol) and in 10 min copper(II) chloride (0.29 mmol). The reaction mixture was stirred for 1 h and warmed to room temperature. The reaction mixture was treated with ammonia as described in the case of **7**, and extracted with ether. The organic layer was concentrated in vacuo, and the residue was purified by silica gel column chromatography (hexane/toluene 5 : 1) from recrystallization in hexane to afford 26 mg of **8** (30% yield) together with *E/Z-11* and their decomplexed products. **8**: mp 228–231 °C; ¹H NMR (400 MHz, CDCl₃) δ = 1.03 (4H, m, CH₂), 1.33 (18H, s, *p*-*t*Bu), 1.58 (36H, s, *o*-*t*Bu), 6.80 (2H, dt, ²J_{PH} = 20 Hz, ³J_{HH} = 7 Hz, P=CH), 7.42 (4H, d, ⁴J_{PH} = 3 Hz, arom); ³¹P NMR (162 MHz, CDCl₃) δ = 202 (dt, ²J_{PH} = 20 Hz, ³J_{PH} = 20 Hz; ¹J_{PW} = 259 Hz); IR (KBr) ν = 2073, 1990, 1932 cm⁻¹; FAB-MS *m/z* 1254 (M⁺). Anal. Calcd for C₅₀H₆₄O₁₀P₂W₂: C, 47.84; H, 5.14. Found: C, 48.11; H, 5.14. b) Crystal data for **8**·1/2C₇H₈: C_{53.5}H₆₈O₁₀P₂W₂, *M* = 1300.73, monoclinic, *P*2₁/c (no. 14), *a* = 13.320(8), *b* = 15.838(2), *c* = 14.112(3) Å, *β* = 102.65(2), *V* = 2904(1) Å³, *Z* = 2, *ρ* = 1.540 g cm⁻³, *μ*(Mo Kα) = 4.070 mm⁻¹, *T* = 150 K, 4936 reflections measured (*2θ*_{max} = 50.0°), 4714 observed [*I* > 2.0σ(*I*)], *R*₁ = 0.050 [*I* > 2.0σ(*I*)]. *R*_w = 0.064 (all data). CCDC-181947.
- R. Appel, V. Winkhaus, and F. Knoch, *Chem. Ber.*, **120**, 125 (1987).
- M. Yoshifuji, K. Toyota, T. Sato, N. Inamoto, and K. Hirotsu, *Heteroat. Chem.*, **1**, 339 (1990).
- J. P. Collmann, L. S. Hegedus, J. R. Norton, and R. G. Finke, “Principles and Applications of Organotransition Metal Chemistry,” University Science Books, New York (1987).