

# Environmentally Benign and Stereoselective Construction of 2-Deoxy and 2,6-Dideoxy- $\beta$ -glycosidic Linkages Employing 2-Deoxy and 2,6-Dideoxyglycosyl Phosphites and Montmorillonite K-10

Hideyuki Nagai, Shuichi Matsumura, and Kazunobu Toshima\*

Department of Applied Chemistry, Faculty of Science and Technology, Keio University,  
3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522


(Received June 19, 2002; CL-020511)

The  $\beta$ -stereoselective glycosidations of 2-deoxy- and 2,6-dideoxyglycosyl diethyl phosphites with alcohols using a heterogeneous and environmentally benign solid acid, montmorillonite K-10, have been developed.

Highly effective, simple and environmentally benign glycosidations have attracted considerable attention in current synthetic organic chemistry related to both biomolecules and functional materials.<sup>1</sup> Some of the challenges for the greening of chemical glycosidations may include the use of a heterogeneous and reusable solid acid as the activator. On the other hand, deoxy sugars frequently appear in the glycosidic components of the bioactive substances.<sup>2,3</sup> Among them, 2-deoxyglycosides including 2,6-dideoxyglycosides are some of the most common and important, and found in many biologically attractive natural products. However, the direct and stereoselective  $\beta$ -glycosidation of a 2-deoxy sugar is quite difficult due to the anomeric effect and the lack of the stereodirecting anchimeric assistance from the C-2 position.<sup>1c,4,5</sup> The most extensively developed strategy for the synthesis of 2-deoxy- $\beta$ -glycosides utilizes donors with equatorial C-2 heteroatom substituents that are reductively removed after the glycosidation event. Furthermore, most of these methods use a homogeneous Lewis acid as the activator, which contaminates the reaction solvent and can not be reused. Therefore, the development of direct and stereoselective  $\beta$ -glycosidations of 2-deoxy sugar in an environmentally compatible manner is of particular interest. Herein, we report the direct and stereoselective glycosidations of the 2-deoxy- and 2,6-dideoxyglycosyl diethyl phosphites<sup>6,7</sup> with alcohols using a heterogeneous and environmentally friendly solid acid, montmorillonite K-10, for the stereoselective synthesis of the 2-deoxy- and 2,6-dideoxy- $\beta$ -glycosides.

In our previous studies, we have demonstrated the novel and stereoselective  $\beta$ -glycosidations of a benzyl-protected glucopyranosyl diethyl phosphite and alcohols using montmorillonite K-10 with a non-participating group.<sup>8</sup> Therefore, based on these results, we first examined the glycosidations of the totally benzylated 2-deoxyglucopyranosyl diethyl phosphite **1** and an alcohol **2** using montmorillonite K-10<sup>9</sup> under several conditions. These results are summarized in Table 1. It was unfortunately found that the glycosidation of **1** ( $\alpha/\beta = 80/20$ ) and **2** under the conditions similar to the optimized conditions<sup>8</sup> for the benzylated glucopyranosyl diethyl phosphite gave the 2-deoxyglucopyranoside **3** in low yield with low stereoselectivity (entry 1 in Table 1). However, when the glycosidation was performed at a lower temperature,  $-78^\circ\text{C}$ , a better yield and  $\beta$ -stereoselectivity were obtained (entry 2 in Table 1). In addition, it was found that the

**Table 1.** Glycosidations of **1** and **2** using montmorillonite K-10 under several conditions




Entry	K-10 /wt%	Solvent	Temp. / $^\circ\text{C}$	Yield /%	$\alpha/\beta$ Ratio <sup>a</sup>
1	100	$\text{CH}_2\text{Cl}_2/\text{MeCN}=10:1$	$-20$	58	43/57
2	100	$\text{CH}_2\text{Cl}_2/\text{MeCN}=10:1$	$-78$	78	26/74
3	100	$\text{CH}_2\text{Cl}_2$	$-78$	87	23/77
4	100	EtCN	$-78$	81	32/68
5	100	PhMe	$-78$	69	25/75
6	100	THF	$-78$	84	47/53
7	100	$\text{Et}_2\text{O}$	$-78$	87	19/81
8	200	$\text{Et}_2\text{O}$	$-78$	82	24/76
9	50	$\text{Et}_2\text{O}$	$-78$	80	21/79
10	20	$\text{Et}_2\text{O}$	$-78$	63	21/79
11	100	$\text{Et}_2\text{O}$	$-50$	80	31/69
12	100	$\text{Et}_2\text{O}$	$-20$	77	44/56


<sup>a</sup>  $\alpha/\beta$  Ratios were determined by HPLC analysis.

glycosidations of **1** and **2** in  $\text{Et}_2\text{O}$  at  $-78^\circ\text{C}$  effectively proceeded to afford **3** in high yield with good  $\beta$ -stereoselectivity (entry 7 in Table 1).  $\text{Et}_2\text{O}$  was shown to be superior to the other solvents (entries 3–7 in Table 1) with respect to both the chemical yield and unusual  $\beta$ -stereoselectivity. Moreover, it was confirmed that the chemical yield and  $\beta$ -stereoselectivity were highly dependent on the amount of montmorillonite K-10 and the reaction temperature. Thus, the highest chemical yield and  $\beta$ -stereoselectivity were realized when 100 wt% montmorillonite K-10 was used in  $\text{Et}_2\text{O}$  at  $-78^\circ\text{C}$  for 1 h (entries 7–12 in Table 1). The optimized conditions for selectively giving the 2-deoxy- $\beta$ -glucopyranoside significantly differed from that for the previously reported  $\beta$ -stereoselective glucosylation.


With these favorable new results, next our attention turned to the glycosidations of a typical 2,6-dideoxy sugar, olivose, because  $\beta$ -olivoside is a very common and important 2,6-dideoxyglycoside.<sup>2,3</sup> Therefore, we examined the glycosidations of the benzylated olivosyl diethyl phosphites **4** ( $\alpha/\beta = 86/14$ ), **5** ( $\alpha/\beta = 72/28$ ), **6** ( $\alpha/\beta = 89/11$ ) and **7** ( $\alpha/\beta = 79/21$ ) with **2**. These results are summarized in Table 2. It was found, in this

**Table 2.** Glycosidations of **4–7** and **2** using montmorillonite K-10


Entry	Donor	R <sup>1</sup>	R <sup>2</sup>	Yield /%	$\alpha/\beta$ Ratio <sup>a</sup>
1	<b>4</b>	Bn	Bn	76	20/80
2	<b>5</b>	Bn	Bz	92	10/90
3	<b>6</b>	Bz	Bn	72	22/78
4	<b>7</b>	Bz	Bz	68	16/84

<sup>a</sup>  $\alpha/\beta$  Ratios were determined by HPLC analysis.**Table 3.** Glycosidations of **1** and **5** with **2, 8–13** using montmorillonite K-10


Entry	Donor	Acceptor	Yield /%	$\alpha/\beta$ Ratio <sup>a</sup>
1	<b>1</b>	<b>2</b>	87	19/81
2	<b>1</b>	<b>8</b>	93	17/83
3	<b>1</b>	<b>9</b>	94	14/86
4	<b>1</b>	<b>10</b>	97	22/78
5	<b>1</b>	<b>11</b>	96	23/77
6	<b>1</b>	<b>12</b>	88	21/79
7	<b>1</b>	<b>13</b>	70	29/71
8	<b>5</b>	<b>2</b>	92	10/90
9	<b>5</b>	<b>8</b>	86	14/86
10	<b>5</b>	<b>9</b>	90	10/90
11	<b>5</b>	<b>10</b>	95	19/81
12	<b>5</b>	<b>11</b>	86	17/83
13	<b>5</b>	<b>12</b>	89	17/83
14	<b>5</b>	<b>13</b>	82	15/85

<sup>a</sup>  $\alpha/\beta$  Ratios were determined by HPLC analysis.

case, that the chemical yield and stereoselectivity were significantly dependent on the protecting groups of the C-3 and 4 hydroxy groups of the donors, and the C-4 acyl group significantly assisted the  $\beta$ -stereoselectivity by the participating

effect.<sup>10</sup> Thus, the glycosidations of **5** and **2** using 100 wt% montmorillonite K-10 in Et<sub>2</sub>O at  $-78^{\circ}\text{C}$  for 1 h effectively proceeded to furnish the corresponding oliviosides in high yield with high  $\beta$ -stereoselectivity (entry 2 in Table 2).

Based on these results, the glycosidations of **1** and **2** using other primary and secondary alcohols **8–13** were next examined. From the results summarized in Table 3, it was found that all the glycosidations of **1** and **5** with **8–13** using 100 wt% montmorillonite K-10 in Et<sub>2</sub>O at  $-78^{\circ}\text{C}$  for 1 h, as well as that of **2**, effectively proceeded to give the corresponding 2-deoxy- $\beta$ -glucopyranosides and  $\beta$ -oliviosides, respectively, in high yields with good stereoselectivities.

Finally, we tested the solid acid recycling in the glycosidations of **1** and **8**. After filtration, washing with chloroform and methanol, and heating at  $100^{\circ}\text{C}/1\text{ mmHg}$  for 12 h, the montmorillonite K-10 was reused for at least three times and showed high yields and stereoselectivities; 1st (93%, 17/83), 2nd (89%, 17/83), 3rd (89%, 17/83), and 4th (87%, 17/83).

A general experimental procedure: To a stirred solution of the glycosyl phosphite **1** ( $\alpha/\beta = 80/20$ , 0.1 mmol) or **5** ( $\alpha/\beta = 72/28$ , 0.1 mmol) and an alcohol (0.2 mmol) in dry Et<sub>2</sub>O (1 ml) was added montmorillonite K-10 (100 wt% to the glycosyl donor **1** or **5**). After stirring for 1 h at  $-78^{\circ}\text{C}$ , the mixture was filtered and the filtrate was concentrated in vacuo. Purification of the residue by flash column chromatography gave the 2-deoxyglucopyranosides or the oliviosides.

## References and Notes

- Review: K. Toshima and K. Tatsuta, *Chem. Rev.*, **93**, 1503 (1993); G.-J. Boons, *Tetrahedron*, **52**, 1095 (1996); B. G. Davis, *J. Chem. Soc., Perkin Trans. 1*, **2000**, 2137.
- P. Jütten and R. Greven, in "Polysaccharides in Medicinal Applications," ed. by S. Dumitriu, Marcel Dekker, New York (1996), p 339.
- A. Kirschning, A. F.-W. Bechtold, and J. Rohr, *Top. Curr. Chem.*, **188**, 1 (1997).
- J. Thiem and W. Klaffke, *Top. Curr. Chem.*, **154**, 285 (1990).
- C. H. Marzabadi and R. W. Frank, *Tetrahedron*, **56**, 8385 (2000).
- An efficient  $\beta$ -stereoselective glycosidation of 2-deoxyglycopyranosyl diethyl phosphites and trimethylsilyl triflate: S. Hashimoto, A. Sano, H. Sakamoto, M. Nakajima, Y. Yanagiya, and S. Ikegami, *Synlett*, **1995**, 1271.
- Y. Guo and G. A. Sulikowski, *J. Am. Chem. Soc.*, **120**, 1392 (1998); R. Pongdee, B. Wu, and G. A. Sulikowski, *Org. Lett.*, **3**, 3523 (2001).
- H. Nagai, S. Matsumura, and K. Toshima, *Tetrahedron Lett.*, **43**, 847 (2002).
- Montmorillonite K-10 was purchased from Aldrich Chemical Company, Inc. and dried at  $200^{\circ}\text{C}/1\text{ mmHg}$  for 12 h before use.
- P. J. Garegg and S. Oscarson, *Carbohydr. Res.*, **50**, C12 (1976); V. Pozsgay, *Carbohydr. Res.*, **69**, 284 (1979); H. Paulsen and R. Leubhn, *Justus Liebigs Ann. Chem.*, **1983**, 1047; C. A. A. van Boeckel, T. Beetz, and S. F. van Aelst, *Tetrahedron Lett.*, **40**, 4097 (1984); T. Jyojima, N. Miyamoto, Y. Ogawa, S. Matsumura, and K. Toshima, *Tetrahedron Lett.*, **40**, 5023 (1999).