

Anomalous ^{13}C NMR Chemical Shifts of High-spin Saddle Shaped Manganese(III) Octaethyltetraphenylporphyrin Complexes

Akira Ikezaki,¹ Mikio Nakamura,^{*1,2,3} and Ru-Jen Cheng⁴¹Department of Chemistry, School of Medicine, Toho University, Ohta-ku, Tokyo 143-8540²Division of Chemistry, Graduate School of Science, Toho University, Funabashi 274-8510³Research Center for Materials with Integrated Properties, Toho University, Funabashi 274-8510⁴Department of Chemistry, National Chung-Hsing University, Taichung, Taiwan 402, Taiwan


(Received October 11, 2005; CL-051294; E-mail: mnakamu@med.toho-u.ac.jp)

NMR spectra of highly saddled Mn(III) porphyrinates such as $\text{Mn}(\text{OETPP})(\text{CD}_3\text{OD})_2^+$ have been measured. The extraordinarily upfield shifted *meso*- ^{13}C signals, which reach as much as -492 ppm at 298 K, are ascribed to the $d_{x^2-y^2}-a_{2u}$, $d_{xy}-a_{1u}$, and $d_{\pi}-3e_g$ interactions, the former two of which occur only when the porphyrin ring is saddled.

Electronic structures of metal porphyrinates are controlled not only by the number and nature of the axial ligands but also by the nonplanarity of porphyrin ring.¹ This is because some metal-porphyrin interactions, which are symmetry forbidden in planar complexes, become possible by the ring deformation.²⁻⁴ In the process of studying the physicochemical properties of deformed Mn(III) porphyrinates, we found that highly saddled $\text{Mn}(\text{OETPP})\text{Cl}$ (**1a**) and $\text{Mn}(\text{OETPP})(\text{CD}_3\text{OD})_2^+$ (**1b**) exhibit quite unusual ^{13}C NMR spectra. In this paper, we report the anomalous ^{13}C NMR chemical shifts of **1** and explain the reasons for the anomaly.

Five-coordinate **1a** was prepared from $(\text{OETPP})\text{H}_2$ and MnCl_2 in refluxed DMF solution,^{5,6} which was converted to six-coordinate **1b** in CD_3OD solution.⁷ These complexes are high-spin ($S = 2$) with $(d_{xy})^1(d_{yz})^1(d_{xz})^1(d_{z^2})^1$ in solution as revealed by the effective magnetic moments, $5.1 \mu_B$ for both complexes, determined by the Evans method.⁸ The ^1H and ^{13}C NMR chemical shifts of the analogous OEP (**2**), TPP (**3**), T'PrP (**4**), and T'BuP (**5**) were also examined for comparison (Scheme 1).⁹⁻¹²

Figures 1 and 2 show the ^1H , ^2H , and ^{13}C NMR spectra. Table 1 lists the chemical shifts at 298 K. The four and two methylene signals in **1a** and **1b**, respectively, suggest that the saddled structure is maintained in solution. Several spectroscopic characteristics are clearly seen in the data of Table 1 which can be explained in terms of the Mn(III)-porphyrin orbital interactions. In



Scheme 1. Mn(III) porphyrinates examined in this study. **1a–5a** are five-coordinate complexes where axial ligand is Cl^- (**1a–4a**) or CH_3CO_2^- (**5a**). **1b–4b** are six-coordinate complexes where axial ligand is CD_3OD .





Figure 1. ^1H NMR spectra of (a) **1a** and (b) phenyl-deuterated **1a-d**₂₀ taken in CD_2Cl_2 at 298 K. (c) ^2H NMR spectrum of **1a-d**₂₀ determined in CHCl_3 at 298 K.

the high-spin five-coordinate complexes (**1a–5a**), the unpaired electrons delocalize through the $d_{\pi}-3e_g$ and $d_{z^2}-a_{2u}$ interactions regardless of the porphyrin structures.^{2,13} The former induces the upfield shift of the Py-H and the downfield shifts of the CH_2 signals while the latter causes the downfield shift of the *meso*-C and *m*-H signals; note that the $3e_g$ orbital has relatively large coefficient at the β -C while the a_{2u} orbital has large coefficient at the *meso*-C as shown in Scheme 2. In the six-coordinate complexes (**1b–4b**), the Mn(III) ion is in the center of the N_4 plane. Thus, the $d_{\pi}-3e_g$ interaction is strengthened while the $d_{z^2}-a_{2u}$ interaction is weakened; note that the a_{2u} and d_{z^2} orbitals are orthogonal in the complexes with D_{2d} symmetry. As a result, both the Py-H and the *meso*-C signals exhibit a large upfield shift by ca. 10 and 100 ppm, respectively.

The data in Table 1 also suggest that the chemical shifts of **1** are quite different from those of **2–5**. The major differences are i) the upfield shift of the *meta*-H, ii) the downfield shifts of the *o*- and *p*-H, and iii) the presence of an extremely upfield shifted *meso*-C signal. These NMR characteristics, which suggest the presence of a large negative spin at the *meso*-carbon atoms,¹⁴⁻¹⁶



Scheme 2. Frontier orbitals of porphyrin.




Figure 2. ^{13}C NMR spectra of *meso*- ^{13}C enriched (a) **1a**, (b) **1b**, (c) **3a**, and (d) **3b** taken at 298 K.

Table 1. ^1H and ^{13}C NMR chemical shifts^a (298 K, δ ppm)

Complexes	^1H NMR							^{13}C NMR
	Py-H	CH ₂	CH ₃	<i>meso</i>	<i>o</i> (CH _{α)}	<i>m</i> (CH _{β)}	<i>p</i>	<i>meso</i>
1a	—	38.4 10.2	13.7 -2.2	1.3	—	13.1 12.3	5.3 4.7	10.5 -406
1b	—	30.3	3.3	1.1	—	13.4	4.3	11.2 -492
2a	—	23.1	0.5	2.7	52.6	—	—	80
2b	—	14.5	2.3	76.4	—	—	—	-24
3a^b	-21.9	—	—	—	— ^d	8.3	7.3	110
3b^b	-30.8	—	—	—	9.2	7.3	7.9	-8
4a^b	-19.4	—	—	—	(9.5)	(3.2)	—	216
4b^b	-31.6	—	—	—	— ^d	(2.1)	—	87
5a^c	-2.9, -28.3	—	—	—	(—)	(3.2)	—	— ^d

^a**1a–5a** were taken in CD_2Cl_2 while **1b–4b** were taken in CD_3OD . ^bref 11. ^c $\text{L} = \text{CH}_3\text{CO}_2^-$, Ref. 12. The pyrrole protons give two signals due to the slow inversion rate of the ruffled porphyrin ring on the ^1H NMR timescale. ^dSignals were too broad.

should be ascribed to the specific orbital interactions such as the $d_{x^2-y^2}-a_{2u}$ and $d_{xy}-a_{1u}$ that can occur only when the planar porphyrin ring is saddled.^{2–4} Cheng and co-workers recently proposed on the basis of the DFT calculation that the interaction between the empty $d_{x^2-y^2}$ orbital and the nitrogen lone pairs in high-spin Mn(III) porphyrinate causes anomalous spin polarization along the bonding skeleton, inducing the negative and positive spin at the pyrrole nitrogen and α carbon atoms, respectively.¹⁷ Thus, it is possible that the interaction between the empty $d_{x^2-y^2}$ and the doubly occupied a_{2u} orbital in saddled complexes induces a large negative spin at the *meso*-carbon and shifts the *meso* signal upfield with a considerable broadening. In addition, the $d_{xy}-a_{1u}$ and $d_{\pi}-3e_g$ interactions could further contribute to the upfield shift since both the a_{1u} and $3e_g$ orbitals have zero coefficient at the *meso*-carbon atoms.¹⁸ We have reported that the structurally analogous intermediate-spin Fe(III) complexes such as $\text{Fe}(\text{OETPP})(4\text{-CNPpy})_2^+$ and $\text{Fe}(\text{OETPP})-$

$(\text{THF})_2^+$ also exhibit the *meso* signals rather upfield.^{16,19,20} Thus, the upfield shift of the *meso* signal must be the phenomenon commonly observed in highly saddled $S = 2$ Mn(III) and $S = 3/2$ Fe(III) complexes.

This work was supported by the Grant-in-Aid for Scientific Research (No. 16550061) from Ministry of Education, Culture, Sports, Science and Technology, Japan. A. I. thanks Futaba Electronics Memorial Foundation and the Sasakawa Scientific Research Grant from The Japan Science Society for financial support.

References and Notes

- W. R. Scheidt in *The Porphyrin Handbook*, ed. by K. M. Kadish, K. M. Smith, R. Guilard, Academic Press, San Diego, **2000**, Vol. 3, Chap. 16, p. 49.
- R.-J. Cheng, P.-Y. Chen, T. Lovell, T. Liu, L. Noodleman, D. A. Case, *J. Am. Chem. Soc.* **2003**, *125*, 6774.
- R.-J. Cheng, Y.-K. Wang, P.-Y. Chen, Y.-P. Han, C.-C. Chang, *Chem. Commun.* **2005**, 1312.
- J. Conradie, A. Ghosh, *J. Phys. Chem. B* **2003**, *107*, 6486.
- A. D. Adler, F. R. Longo, F. Kampas, J. Kim, *J. Inorg. Nucl. Chem.* **1970**, *32*, 2443.
- K. M. Barkigia, M. D. Berber, J. Fajer, C. J. Medforth, M. W. Renner, K. M. Smith, *J. Am. Chem. Soc.* **1990**, *112*, 8851.
- UV-Vis($\lambda_{\text{max}}/\text{nm}$): **1a**(CH_2Cl_2) 375, 405, 496, 595, 635; **1b**(CH_3OH) 391, 410, 484, 579, 616.
- D. F. Evans, T. A. James, *J. Chem. Soc., Dalton Trans.* **1979**, 723.
- H. M. Goff, A. P. Hansen, *Inorg. Chem.* **1984**, *23*, 321.
- P. Turner, M. J. Gunter, *Inorg. Chem.* **1994**, *33*, 1406.
- A. Ikezaki, M. Nakamura, *Inorg. Chem.* **2003**, *42*, 2301.
- A. Ikezaki, M. Nakamura, *Chem. Lett.* **2005**, *34*, 1046.
- J. Mao, Y. Zhang, E. Oldfield, *J. Am. Chem. Soc.* **2002**, *124*, 13911.
- Presence of a negative spin at the *meso* carbon was first proposed by Yatsunyk and Walker¹⁵ to explain the ^1H NMR chemical shifts of the $S = 3/2$ $\text{Fe}(\text{OETPP})(4\text{-CNPpy})_2^+$ reported by Nakamura and co-workers.¹⁶
- L. A. Yatsunyk, F. A. Walker, *Inorg. Chem.* **2004**, *43*, 757.
- T. Ikeue, Y. Ohgo, T. Yamaguchi, M. Takahashi, M. Takeda, M. Nakamura, *Angew. Chem., Int. Ed.* **2001**, *40*, 2617.
- R.-J. Cheng, S.-H. Chang, K.-C. Hung, *Chem. Commun.*, submitted.
- The $d_{\pi}-3e_g$ interaction in saddle shaped low-spin ($S = 1/2$) Fe(III) porphyrinate, $\text{Fe}(\text{OETPP})(\text{imidazole})_2^+$, induces the upfield shift of the *meso* signal by 110 ppm at 298 K. The upfield shift of the *meso* signal due to the $d_{\pi}-3e_g$ interaction in **1b** could be much larger because the complex has two unpaired electrons in the d_{π} orbitals.
- T. Sakai, Y. Ohgo, T. Ikeue, M. Takahashi, M. Takeda, M. Nakamura, *J. Am. Chem. Soc.* **2003**, *125*, 13028.
- A. Hoshino, Y. Ohgo, M. Nakamura, *Inorg. Chem.* **2005**, *44*, 7333.