

## ALKALOIDS OF CORYDALIS SEVERTZOVII

M. S. Yunusov and S. Yu. Yunusov

*Khimiya Prirodnkh Soedinenii*, Vol. 4, No. 1, pp. 61-62, 1968

From the epigeal part of C. severtzovii Rgl., collected in the flowering stage near Tashkent (2.4 kg of air-dry plant material) by ordinary chloroform extraction we have isolated 20 g (0.83%) of total alkaloids. The separation of the latter yielded protopine (0.2 g), cryptopine (0.15 g), and corlumine (4 g). The authenticity of the first two was shown by their comparison with authentic samples and that of the third by the identification of the products of oxidative decomposition under the action of nitric acid.

After the extraction of the alkaloids with chloroform, the plant was treated with methanol. The methanolic extract was evaporated and the residue was diluted with 5% sulfuric acid. The acid solution was washed with ether and made alkaline with ammonia, and the alkaloids were extracted with ether. This gave 0.7 g of total alkaloids, from which a further 0.2 g of corlumine was isolated. When the ether used for washing was evaporated, fumaric acid (0.7 g) crystallized out.

From C. severtzovii (180 g) gathered on the northern slopes of the Turkestan range (upper reaches of the R. Zaa-min) we isolated protopine (0.3 g) and a base (0.15 g) with mp 193-195°C (methanol-acetone);  $[\alpha]_D^{33} -110^\circ$  (c 0.27; chloroform); UV spectrum,  $\lambda_{\max}$  285 m $\mu$  (log  $\epsilon$  3.86). The IR spectrum of the base was identical with that of d-bicuculline. A mixture of the two samples melted at 187-210°C. These results permitted us to consider that the base isolated is the levorotary form of the known alkaloid d-bicuculline, and this is apparently the first time it has been found in a plant.

In addition to protopine and l-bicuculline, we isolated small amounts (10 mg) of a base with mp 200-202°C, IR spectrum: 920 cm<sup>-1</sup>, 940, 1470, 1485, 1505, 2770, 2790, 3400 cm<sup>-1</sup>. The base contains a methylenedioxide group (reaction with gallic acid).

Protopine, sanguinarine, and  $\alpha$ -allocryptopine have previously been isolated from the tubers of C. severtzovii [1].

Thus, the two cases cited show that different alkaloids (with the exception of protopine, found in both plants but in different amounts) have been found in one and the same species. This fact is a striking example of the dependence of the qualitative and quantitative composition of the alkaloids on the growth site of the plant [2].

### REFERENCES

1. B. K. Rostotskii and I. L. L'vova, *Med. prom SSSR*, no. 2, 17, 1964.
2. S. Yu. Yunusov, *Jubilee Symposium Devoted to the 25th Anniversary of the UzSSR* [in Russian], p. 223, Tashkent, 1950.

11 November 1967

Institute of the Chemistry of Plant  
Substances, AS UzSSR

UDC 577.1:547.965

## AMMONOLYSIS OF PHENYLTHIOHYDANTOIN (PTH) DERIVATIVES OF AMINO ACIDS

G. Ya. Lamm, A. R. Rakhimov, K. G. Ioffe, and I. A. Asatov

*Khimiya Prirodnkh Soedinenii*, Vol. 4, No. 1, pp. 62-63, 1968

In the determination of N-terminal amino acids in proteins and peptides by the phenyl isothiocyanate method, the amino acids are identified by chromatographing both the PTHs [1] and also the free amino acids obtained by regeneration from the PTHs [2-5]. We have attempted to develop a method of identifying the amino acids by the hydrolysis with ammonia of the corresponding PTHs.

To five 100- $\gamma$  samples of the PTHs of serine, alanine, aspartic acid, tyrosine, and valine we added 300  $\mu$ l of 28% ammonia solution. The mixture was heated in a sealed capillary in the boiling water bath for 2, 4, 6, 9, and 12 hr (figure). One-dimensional chromatography was carried out in the water-saturated phenol system; the chromatograms were revealed with 0.25% ninhydrin in butanol.